

This is a preprint of an revised article accepted for publication in ChemPlusChem:

Nitrocatechols as Tractable Surface Release Systems

Robin Wehlauch, Johannes Hoecker, Karl Gademann

Department of Chemistry, University of Basel
National Centre of Competence in Research "Chemical Biology"
St. Johanns-Ring 19, 4056 Basel (Switzerland)

E-mail: karl.gademann@unibas.ch
Homepage: <http://chemie.unibas.ch/~gademann>

DOI: 10.1002/cplu.201200251

Please cite this article as:

R. Wehlauch, J. Hoecker, K. Gademann *ChemPlusChem* **2012**, *77*, doi:10.1002/cplu.201200251

DOI: 10.1002/cplu.201200251

Nitrocatechols as Tractable Surface Release Systems**

Robin Wehlauch,[✧] Johannes Hoecker[✧] and Karl Gademann^{*}

We report the development of a molecular surface modification platform based on nitrocatechol derivatives that allows for small molecule functionalization of TiO₂ under mild aqueous conditions and efficient release triggered by light therefore uncaging a small molecule cargo on demand. Surface modifications by using molecular approaches have found applications in a wide variety of fields, as they allow combining the bulk properties of a material with the molecular features of the coating, which can additionally be tailored by synthetic chemistry.¹ A central challenge remains the nature of the molecular anchor that links coating and surface. In this respect, catechols display unique properties due to their ability to strongly bind to metal oxides in aqueous media. Originally found in mussel adhesive proteins,² dihydroxyphenylalanine (DOPA) or dopamine derived catechols were successfully used in surface modifications despite their shortcomings of weak binding and sensitivity to oxidation. We have introduced catechols with electron-withdrawing substituents based on the iron chelator anachelin that result in stable coatings with non-polymeric architectures.³ A second-generation design introduced nitrocatechols such as nitrodopamine due to the ease of preparation, improved adhesion properties and stability towards oxidation.⁴ All these anchors have found widespread applications,^{1,5} for example, we have shown that such catechols can be used to functionalize surfaces with complex antibiotics by an operationally simple dip-and-rinse procedure to generate antimicrobial surfaces.⁶

In addition to immobilization, the release of small molecules from a modified surface triggered by an external stimulus would be highly desirable. Potential applications would range from drug delivery, small molecule microarrays to selective probes in chemical biology. In particular in the latter area, such an approach would combine the well-known strategy of temporarily disabling the biological activity of a small molecule ('caging') with the possibility of controlled release on demand. This would allow for both high spatial control (through immobilization) and temporal control (through caging) of the biological activity of a small molecule, two key elements in chemical biology approaches.^{7,8} In

this paper, we report on the development of such a molecular platform that allows for the spatio-temporal control of small molecule release. In particular, we propose a bio-inspired approach that leverages the presence of a catechol (easily tunable by a variety of parameters such as pH, ion strength or temperature) with the versatility of the nitrophenyl system (Figure 1).




Figure 1. Concept of the light induced surface release of carrier molecules for the photolabile NPP protecting group.


Among the most commonly used photocleavable protecting groups are several nitrophenyl derivatives, including (2-nitrophenyl)ethyl (NPE) and (2-nitrophenyl)propyl (NPP) derivatives.^{9,10} However, upon inspection of the literature, we were surprised to find that no free nitrocatechols have been utilized as photocleavable groups and the surface functionalization and release properties of such systems have not been investigated.¹¹ We have targeted both NPE and NPP systems **1** and **2**, respectively, and demonstrate in this work that controlled bonding and release is readily achieved for the latter derivatives.

The first target compound constituted the 2-nitrophenyl ethyl derivative **1**. Its synthesis started with the preparation of nitrocatechol **4** by nitration of commercially available 3,4-(methylenedioxy)acetophenone (**3**) using HNO₃ in acetic acid (Scheme 1). Treatment with AlCl₃ at low temperature and subsequent hydrolysis with conc. HBr gave nitrocatechol **4** in good yield. Catechol **4** was then further protected as MOM ether and the keto group was reduced by using NaBH₄ leading to alcohol **5**. In the context of this synthesis, it was found necessary to protect the catecholate OH groups, in order to prevent attachment of this compound to silica during chromatographic purification. The alcohol **5** was then attached to the fluorophore¹² **6** via Mitsunobu reaction in moderate yields. When alcohol **5** was coupled, reactions were performed under exclusion of light to prevent decomposition of the formed product. During isolation of

R. Wehlauch, J. Hoecker, Prof. K. Gademann
Department of Chemistry, University of Basel
National Centre of Competence in Research "Chemical Biology"
St. Johannis-Ring 19, 4056 Basel (Switzerland)
E-mail: karl.gademann@unibas.ch
Homepage: <http://chemie.unibas.ch/~gademann>


[✧] Both authors contributed equally to this work.

Supporting information for this article is available on the WWW
under <http://dx.doi.org/10.1002/cplu.200xxxxx>.



Scheme 1. Preparation of NPE conjugate **1** for surface modification: a) HNO_3 , AcOH , 0°C to rt, 2.5 h, 59%; b) AlCl_3 , DCE , -5°C , 1 h, then 48% HBr , rt, 24 h, 84%; c) MOMCl , K_2CO_3 , MeCN , 0°C to rt, 3 h, 87%; d) NaBH_4 , MeOH , 0°C to rt, 3.5 h, 99%; e) **6**, PPH_3 , DIAD , THF , 0°C to rt, over night, 41%; f) TFA , H_2O .

the photolabile compound from the reaction mixtures, it became already apparent that the uncaging process could be induced, when TLC plates were exposed to UV light at 366 nm (see video, supporting information). The MOM protecting group could be finally removed using aqueous TFA , leading to the NPE derivative **1**.




Scheme 2. Deprotection studies of fluorinated and unsubstituted catechols.

With the NPE derivative in hand, we sought to prepare the related unsubstituted compound **7** ($X = \text{H}$) as well as the fluorinated compound **8** ($X = \text{F}$, for detailed synthesis see supporting information) to investigate the role of the nitro substituent on the cleavage mechanism. There are at least two fundamentally different pathways for cleavage of such catechols from titania possible, either via (1) photocatalytic oxidation to the quinone⁷ or (2) via nitroaryl mediated bond cleavage.^{8,9} In order to investigate these two options and to delineate the role of the nitro substituent, we have prepared control compounds **7** and **8** lacking this substituent, where only photocatalytic oxidation to the quinone would be possible. We were surprised to discover that the deprotection under a variety of conditions of these compounds proved to be impossible, as only decomposition of the starting material and the free fluorophore could be detected. This could be explained by the formation of reactive quinone methides by cleavage of the fluorophore at the benzylic position.¹³

In order to circumvent the decomposition *via* the quinone methide pathway, we wanted to investigate the corresponding *homologated* propyl substituted nitrophenols **2**. Therefore, acetonide **10** was successfully prepared from catechol **9** by using 2,2-dimethoxypropane and catalytic amounts of $p\text{TsOH}$ in refluxing benzene (Scheme 3). Subsequent nitration of the acetonide with half-concentrated HNO_3 afforded **10** after 1.5 h in very good yield. To our delight, these strongly acidic, aqueous conditions did not affect the acetonide protection group. A crystal structure of **10** could be obtained, which confirmed the correct installment of the nitro group (see supporting information). The reverse synthetic order, first nitration followed by protection, failed as no acetonide formation could be detected. The crucial elongation reaction of **10** with paraformaldehyde and Triton B gave the desired product **11**,¹⁴ although only moderate conversion was achieved (with 40% starting material **10** recovered). With the catechol protected as the acetonide, carbonate formation using triphosgene and triethylamine in THF led to the desired chloro-

formate in quantitative yield within 25 min at 0°C .¹⁵ To prevent decomposition, the chloroformate was coupled immediately to coumarin **12** to afford the desired carbonate, which was deprotected to the target catechol **2** using neat TFA . In total the



Scheme 3. Preparation of NPP conjugate **2** for surface modification: a) 2,2-dimethoxypropane, cat. $p\text{TsOH}$, benzene, reflux, overnight, b) HNO_3 , H_2O , 0°C to rt, 1.5 h, 86% (over 2 steps); c) CH_2O , Triton B in MeOH , 85°C , 65 h, 80% (*brsm*); d) i) triphosgene, Et_3N , THF , 0°C , 25 min, ii) **3b**, pyridine, DCM , 0°C to rt, 1.5 h; e) TFA , H_2O , rt, over night, 68% (over 2 steps). DIAD = diisopropyl diazodicarboxylate; *brsm* = based on recovery of starting material

preparation of **2** was achieved in five steps using only two purifications rendering **2** an easily accessible and attractive candidate for further investigations. In particular, we expect that the alcohol **11** could serve as an ideal starting point for the attachment of various cargo compounds by coupling chemistry.

The stability of the different caged compounds was then investigated using a variety of methods and assays. Gratifyingly, the NPP linker **2** was fully stable to hydrolysis in aqueous medium (MOPS buffer) at pH 5.5 in the dark for at least 72 hours. This is contrast to the ethyl derivative **1**, which displays only limited stability. Photocleavage of catechol **2** was investigated by detection of the evolving fluorescence during the release of coumarin **12** to determine the half-life time of **2** in aqueous solution (MOPS buffer) under near UV-irradiation.¹⁶ As expected, a rapid increase in fluorescence was detected during the first minutes of irradiation and the fluorescence maximum was





Figure 2. A) Fluorescence spectra of **12** and **2** after certain irradiation times; B) Fluorescence emission intensity at 454 nm depending on the irradiation time at 366 nm of **2** as a $5 \cdot 10^{-5}$ M solution in MOPS buffer; C) UV spectra of **2** and **12**; D) Decay of **2** upon UV irradiation at 366 nm, determined by HPLC-MS.

identified at 454 nm (figure 2A). However, after approximately 15 min, the detected fluorescence intensity was decreasing, indicating that photobleaching of coumarin **12** occurred, which could be confirmed by control experiments (fig. 2B, and supporting information). Quantification by HPLC-MS determined the half-life time of **2** to be around 12 min at a concentration of $5 \cdot 10^{-5}$ M in buffer (figure 2D). This is an attractively short period for cleavage, since the light intensity of a simple laboratory UV-lamp is rather low ($\sim 20 \text{ mW cm}^{-2}$).



Scheme 4. Immobilization of nitrocatechol **2** on TiO₂ microparticles to deliver **Ti-2** and release of the cargo upon irradiation.

To evaluate the surface release properties, TiO₂ particles (1.0-2.0 μm) were functionalized with nitrocatechol **2** in MOPS buffer at 55°C according to previously developed conditions (Scheme 4).¹⁷ After the incubation, the functionalized particles **Ti-2** were washed three times with CH₃CN, and HPLC analysis of the washing solutions determined only small amounts of fluorophore **12** and no free **2** present.

The release of the fluorescent cargo from the functionalized beads was investigated next. The functionalized TiO₂ beads **Ti-2** were suspended in MOPS buffer and the mixture was irradiated. Fluorescence became immediately visible (figure 3B). Aliquots were taken after certain irradiation time intervals and analyzed by




Figure 3. A) TLC plate coated with nitrocatechol **2** (**Si-2**) and irradiated for 60 sec at 366 nm under a printed foil, after 10 sec exposure to UV light without foil (1), after 60 sec (2) and after 180 sec (3) UV irradiation at 366 nm (see videos, supporting information); B) **Ti-2** (left) and negative control (uncoated TiO₂ particles, right) in MOPS buffer upon UV irradiation; C) Release kinetics of **12** from **Ti-2** upon UV irradiation at 366 nm in MOPS buffer.

HPLC at 366 nm. The obtained data demonstrates that no catechol **2** is present and the amount of **12** was increasing over the irradiation time (figure 3C), providing evidence that photocleavage of surface-adsorbed nitrocatechol **2** and concomitant release of coumarin **12** has occurred. In this reaction setup, a half-life time of ca. 19 min was determined for the release of **12**. Qualitatively, these observations could be corroborated by the immobilization of **2** on silica (TLC plate) and controlled release under UV light (figure 3A; see supporting information for videos). In this simple application, a printed foil was used as template to generate the desired surface pattern, which could be unmasked by release of the fluorophore via irradiation.

In conclusion, we report in this communication the development of a bio-inspired surface modification platform based on nitrocatechols that allows for the controlled release of small molecules under an external stimulus. Salient features of this method involve (1) ease of functionalization of TiO₂ particles under an operationally simple dip-and-rinse procedure, (2) stability of the resulting functionalized particles to repeated washing and (3) rapid release of the small molecule cargo under an external stimulus, *i.e.* UV light. We think that this method displays advantages with regard to controlled release. This surface modification platform might find applications in drug delivery, as caged probes in chemical biology or for direct assays 'on chip'. The utilization of this platform for the immobilization and release of biologically active small molecules is currently under investigation in our laboratories.

Experimental Section

Experimental details for the cleavage, along with characterization and spectral data of all compounds are provided in the supporting information.

Acknowledgements

K. G. is a European Young Investigator (EURYI). We thank Dr M. Neuburger for X-ray analyses and J. Gomes for useful discussions. We gratefully acknowledge financial support by the SNF (PE002-117136/1).


Keywords: release on demand • adhesion • biomimetic • photochemistry • synthetic design

- [1] Reviews: a) Q. Ye, F. Zhou, W. Liu, *Chem. Soc. Rev.* **2011**, *40*, 4244-4258; b) J. L. Dalsin, P. B. Messersmith, *Mater. Today* **2005**, *8*, 38-46.
- [2] a) H. Zhao, J. H. Waite, *J. Biol. Chem.* **2006**, *281*, 26150-26158; b) H. Lee, N. F. Scherer, P. B. Messersmith, *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 12999-13003; c) J. H. Waite, *Nature Mater.* **2008**, *7*, 8-9; d) H. Zeng, D. S. Hwang, J. N. Israelachvili, J. H. Waite, *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 12850-12853; e) H. Lee, S. M. Dellatore, W. M. Miller, P. B. Messersmith, *Science* **2007**, *318*, 426-430.
- [3] S. Zürcher, D. Wäckerlin, Y. Bethuel, B. Malisova, M. Textor, S. Tosatti, K. Gademann, *J. Am. Chem. Soc.* **2006**, *128*, 1064-1065.
- [4] a) B. Malisova, S. Tosatti, M. Textor, K. Gademann, S. Zürcher, *Langmuir* **2010**, *26*, 4018-4026; b) S. Saxer, C. Portmann, S. Tosatti, K. Gademann, S. Zürcher, M. Textor, *Macromolecules* **2010**, *43*, 1050-1060; c) J.-Y. Wach, B. Malisova, S. Bonazzi, S. Tosatti, M. Textor, S. Zürcher, K. Gademann, *Chem. Eur. J.* **2008**, *14*, 10579-10584; d)

- Review: K. Gademann, J. Kobylinska, J.-Y. Wach, T. M. Woods, *Biometals* **2009**, *22*, 595-604.
- [5] For recent examples: a) B. Geiseler, L. Fruk, *J. Mater. Chem.* **2012**, *22*, 735-741; b) A. S. Goldmann, C. Schödel, A. Walther, J. Yuan, K. Loos, A. H. E. Müller, *Macromol. Rapid Commun.* **2010**, *31*, 1608-1615.
- [6] J.-Y. Wach, S. Bonazzi, K. Gademann, *Angew. Chem.* **2008**, *120*, 7232-7235; *Angew. Chem. Int. Ed.* **2008**, *47*, 7123-7126.
- [7] T. Tachikawa, Y. Asanoi, K. Kawai, S. Tojo, A. Sugimoto, M. Fujitsuka, T. Majima, *Chem. Eur. J.* **2008**, *14*, 1492-1498.
- [8] X. Wang, S. Werner, T. Weiß, K. Liefelth, C. Hoffmann, *RSC Adv.* **2012**, *2*, 156-160.
- [9] Reviews: a) G. Mayer, A. Heckel, *Angew. Chem.* **2006**, *118*, 5020-5042; *Angew. Chem. Int. Ed.* **2006**, *45*, 4900-4921; b) D. Puliti, D. Warther, C. Orange, A. Specht, M. Goeldner, *Biorg. Med. Chem.* **2011**, *19*, 1023-1029; c) D. Warther, S. Gug, A. Specht, F. Bolze, J.-F. Nicoud, A. Mourot, M. Goeldner, *Biorg. Med. Chem.* **2010**, *18*, 7753-7758; d) C. G. Bochet, *J. Chem. Soc., Perkin Trans. 1* **2002**, 125-142.
- [10] a) I. Aujard, C. Benbrahim, M. Gouget, O. Ruel, J.-B. Baudin, P. Neveu, L. Jullien, *Chem. Eur. J.* **2006**, *12*, 6865-6879; b) D. Warther, F. Bolze, J. Léonard, S. Gug, A. Specht, D. Puliti, X.-H. Sun, P. Kessler, Y. Lutz, J.-L. Vonesch, B. Winsor, J.-F. Nicoud, M. Goeldner, *J. Am. Chem. Soc.* **2010**, *132*, 2585-2990.
- [11] During the preparation of this manuscript for publication, an article describing a similar system appeared: Z. Shafiq, J. Cui, L. Pastor-Pérez, V. San Miguel, R. A. Gropeanu, C. Serrano, A. del Campo, *Angew. Chem.* **2012**, *124*, 4408-4411; *Angew. Chem. Int. Ed.* **2012**, *51*, 4332-4335.
- [12] F. Fringuelli, O. Piermatti, F. Pizzo, *Synlett* **2003**, 2331-2334.
- [13] a) M. Sugumaran, V. Semensi, *J. Biol. Chem.* **1990**, *266*, 6073-6078; b) M. Sugumaran, V. Semensi, S. J. Saul, *Arch. Inst. Biochem. Phys.* **1988**, *9*, 269-281; c) M. Sugumaran in *Biological Oxidation Systems*, Vol 1, (Eds: C. C. Reddy, G. A. Hamilton, K. M. Madyastha), Academic-Press, New York, **1990**, pp. 347-363.
- [14] K. R. Bhushan, *Org. Biomol. Chem.* **2006**, *4*, 1857-1859.
- [15] a) G. H. McGall, A. D. Barone, M. Diggelmann, S. P. A. Fodor, E. Gentalen, N. Ngo, *J. Am. Chem. Soc.* **1997**, *119*, 5081-5090; b) A. Gautier, D. P. Nguyen, H. Lusic, W. An, A. Deiters, J. W. Chin, *J. Am. Chem. Soc.* **2010**, *132*, 4086-4088.
- [16] S. Walbert, W. Pfeleiderer, U. E. Steiner, *Helv. Chim. Acta* **2001**, *84*, 1601-1611.
- [17] J. Gomes, K. Gademann, *unpublished results*.
-
- Received: ((will be filled in by the editorial staff))
Published online: ((will be filled in by the editorial staff))

Entry for the Table of Contents

COMMUNICATION



X = fluorophore, bioactive small molecule, polymer, proteins, DNA, etc.

R. Wehlauch, J. Hoecker, K. Gademann*

Page No. – Page No.

Nitrocatechols as Tractable Surface Release System

Anchor aweigh! The synthesis and evaluation of new nitrochatechols for surface modification and tractable release on TiO_2 are reported. The properties of catecholate anchoring and nitrobased photocleavable release were merged to create a new tool for surface immobilization and controlled release of small molecules, which was investigated on caged fluorophores as proof-of-principle.