

Air Pollution from Road Traffic and Systemic Inflammation in Adults: A Cross-Sectional Analysis in the European ESCAPE Project

Timo Lanki,¹ Regina Hampel,² Pekka Tiittanen,¹ Silke Andrich,³ Rob Beelen,⁴ Bert Brunekreef,^{4,5} Julia Dratva,^{6,7} Ulf De Faire,⁸ Kateryna B. Fuks,⁹ Barbara Hoffmann,^{9,10} Medea Imboden,^{6,7} Pekka Jousilahti,¹¹ Wolfgang Koenig,^{12,13} Amir A. Mahabadi,¹⁴ Nino Künzli,^{6,7} Nancy L. Pedersen,¹⁵ Johanna Penell,⁸ Göran Pershagen,⁸ Nicole M. Probst-Hensch,^{6,7} Emmanuel Schaffner,^{6,7} Christian Schindler,^{6,7} Dorothea Sugiri,⁹ Wim J.R. Swart,¹⁶ Ming-Yi Tsai,^{6,7,17} Anu W. Turunen,¹ Gudrun Weinmayr,^{9,18} Kathrin Wolf,² Tarja Yli-Tuomi,¹ and Annette Peters²

¹Department of Health Protection, National Institute for Health and Welfare, Kuopio, Finland; ²Institute of Epidemiology II, German Research Center for Environmental Health, Helmholtz Zentrum München, Neuherberg, Germany; ³Biometry and Epidemiology, Institute for Medical Informatics, University Hospital Essen, University of Duisburg-Essen, Essen, Germany; ⁴Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands; ⁵Julius Center for Health Sciences and Primary Care, Department of Epidemiology, University Medical Center Utrecht, Utrecht, the Netherlands; ⁶Swiss Tropical and Public Health Institute, Basel, Switzerland; ⁷University of Basel, Basel, Switzerland; ⁸Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; ⁹IUF-Leibniz Research Institute for Environmental Medicine, Düsseldorf, Germany; ¹⁰Medical Faculty, Deanery of Medicine, Heinrich Heine University Düsseldorf, Düsseldorf, Germany; ¹¹Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland; ¹²Department of Internal Medicine II-Cardiology, University of Ulm Medical Center, Ulm, Germany; ¹³Deutsches Herzzentrum München, Technische Universität München, Munich, Germany; ¹⁴West-German Heart Center, Department of Cardiology, University of Duisburg-Essen, Essen, Germany; ¹⁵Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden; ¹⁶Centre for Environmental Health, National Institute for Public Health and the Environment, Bilthoven, the Netherlands; ¹⁷Department of Environmental and Occupational Health Sciences, University of Washington, Seattle, USA; ¹⁸Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany

BACKGROUND: Exposure to particulate matter air pollution (PM) has been associated with cardiovascular diseases.

OBJECTIVES: In this study we evaluated whether annual exposure to ambient air pollution is associated with systemic inflammation, which is hypothesized to be an intermediate step to cardiovascular disease.

METHODS: Six cohorts of adults from Central and Northern Europe were used in this cross-sectional study as part of the larger ESCAPE project (European Study of Cohorts for Air Pollution Effects). Data on levels of blood markers for systemic inflammation—high-sensitivity C-reactive protein (CRP) and fibrinogen—were available for 22,561 and 17,428 persons, respectively. Land use regression models were used to estimate cohort participants' long-term exposure to various size fractions of PM, soot, and nitrogen oxides (NO_x). In addition, traffic intensity on the closest street and traffic load within 100 m from home were used as indicators of traffic air pollution exposure.

RESULTS: Particulate air pollution was not associated with systemic inflammation. However, cohort participants living on a busy (> 10,000 vehicles/day) road had elevated CRP values (10.2%; 95% CI: 2.4, 18.8%, compared with persons living on a quiet residential street with < 1,000 vehicles/day). Annual NO_x concentration was also positively associated with levels of CRP (3.2%; 95% CI: 0.3, 6.1 per 20 µg/m³), but the effect estimate was more sensitive to model adjustments. For fibrinogen, no consistent associations were observed.

CONCLUSIONS: Living close to busy traffic was associated with increased CRP concentrations, a known risk factor for cardiovascular diseases. However, it remains unclear which specific air pollutants are responsible for the association.

CITATION: Lanki T, Hampel R, Tiittanen P, Andrich S, Beelen R, Brunekreef B, Dratva J, De Faire U, Fuks KB, Hoffmann B, Imboden M, Jousilahti P, Koenig W, Mahabadi AA, Künzli N, Pedersen NL, Penell J, Pershagen G, Probst-Hensch NM, Schaffner E, Schindler C, Sugiri D, Swart WJ, Tsai MY, Turunen AW, Weinmayr G, Wolf K, Yli-Tuomi T, Peters A. 2015. Air pollution from road traffic and systemic inflammation in adults: a cross-sectional analysis in the European ESCAPE project. *Environ Health Perspect* 123:785–791; <http://dx.doi.org/10.1289/ehp.1408224>

Introduction

Exposure to ambient particulate matter pollution (PM) affects both respiratory and cardiovascular health, and is a major contributor to the global burden of disease (Lim et al. 2012). One of the main sources of particulate air pollution is road traffic. Tailpipe exhaust is a source of fine combustion particles (soot), which are associated with cardiorespiratory health effects and premature mortality (Janssen et al. 2012). However, road traffic also produces road dust consisting mainly of coarse particles. There is increasing evidence on the harmfulness of

coarse particulate matter (Brunekreef and Forsberg 2005).

Ischemic heart disease is the leading contributor to the burden of disease worldwide, and stroke is ranked third (Murray et al. 2012). An American Heart Association Statement has concluded that exposure to fine particulate air pollution is likely to be causally linked to cardiovascular disease (CVD) (Brook et al. 2010). Because of its ubiquitous presence, particulate air pollution is a relevant trigger of myocardial infarction on a population level (Nawrot et al. 2011). There is less, but increasing, evidence on the associations between long-term exposure to

traffic-related air pollutants and the incidence of CVD (Brook et al. 2010).

It has been hypothesized that increased systemic inflammation—a risk factor for, as well as a consequence of, atherosclerosis—is

Address correspondence to T. Lanki, National Institute for Health and Welfare, Department of Health Protection, Neulaniementie 4 (P.O. Box 95), FI-70210 Kuopio, Finland. Telephone: 358 29 524 6326. E-mail: timo.lanki@thl.fi

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1408224>).

The research leading to these results has received funding from the European Community's Seventh Framework Program (FP7/2007–2011) under grant agreement no. 211250. Additional funding for the National FINRISK study came from the Academy of Finland (project no. 129317). The KORA (Cooperative Health Research in the Augsburg Region) research platform and the MONICA (Multinational MONITORING of trends and determinants in Cardiovascular disease) Augsburg studies were initiated and financed by the Helmholtz Zentrum München, German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research and by the State of Bavaria. The HNR (Heinz Nixdorf Recall) study was supported by the Heinz Nixdorf Foundation, the German Ministry of Education and Science, the German Research Foundation (DFG; projects JO-170/8-1, HO 3314/2-1, SI 236/8-1, and SI236/9-1). The Swiss Study on Air Pollution and Lung Disease in Adults (SAPALDIA) was primarily supported by the Swiss National Science Foundation (grants no. 833CSGO-134276/1, 33CSGO-108796, 3247BO-104283, 3247BO-104288, 3247BO-104284, 93247-065896, 3100-059302, 3200-052720, 3200-042532, and 4026-028099) and by the Federal Office for the Environment. The financial support for the combined work with the Stockholm studies was provided by the Swedish Environmental Protection Agency, the Swedish Heart-Lung Foundation, and the Swedish Council for Working Life and Social Research. See Supplemental Material for other funders.

The authors declare they have no actual or potential competing financial interests.

Received: 4 February 2014; Accepted: 23 March 2015; Advance Publication: 27 March 2015; Final Publication: 1 August 2015.

the main underlying mechanism linking long-term PM exposure to incidence of CVD (Brook et al. 2010). A number of studies have associated daily variation in PM with short-term changes in the concentrations of circulating inflammatory markers (Brook et al. 2010). However, it is not known whether recurrent high short-term exposures may also lead to chronic and partly self-sustained systemic inflammation. Associations between long-term within-city variation in PM, typically driven by traffic emissions, and systemic inflammation have been evaluated only in a few population-level studies, and with conflicting results (Chuang et al. 2011; Forbes et al. 2009; Hoffmann et al. 2009; Panasevich et al. 2009; Schwartz 2001). Therefore, we evaluated within the large multi-center ESCAPE project (European Study of Cohorts for Air Pollution Effects) the associations of long-term exposure to traffic and traffic-related air pollution with high-sensitivity C-reactive protein (CRP), a clinically established sensitive marker of systemic inflammation (Emerging Risk Factors Collaboration et al. 2010), and fibrinogen, a hemostatic factor with proinflammatory properties.

Methods

Study population. The ESCAPE project linked data from existing cohorts in different parts of Europe with modeled long-term residential air pollution concentrations. The studies were approved by the local ethical committees and conducted according to the guidelines laid down in the Declaration of Helsinki. All data were collected in accordance with the Declaration of Helsinki, and all participants gave an informed consent before the study. Data on blood levels of high-sensitivity CRP was available for five cohorts in Northern and Central Europe: the National FINRISK study in Finland, KORA (Cooperative Health Research in the Region of Augsburg) and HNR (Heinz Nixdorf Recall) in Germany, SAPALDIA (Swiss Study on Air Pollution and Lung and Heart Disease In Adults) in Switzerland, and TwinGene in Sweden. For the analyses of fibrinogen, four cohorts were available: FINRISK, KORA, HNR, and the “60-year-olds” study in Sweden. Descriptive

statistics of these cohorts are presented in Table 1. See Supplemental Material, “Cohort descriptions” and “Biochemical measurements,” for further information on the cohorts, as well as the methods for estimating CRP and fibrinogen.

Exposure data. Air pollution measurements and modeling followed common study protocols (ESCAPE 2008) and were coordinated by the Institute for Risk Assessment Sciences of Utrecht University. Concentrations of several size fractions of particulate air pollution (PM_{2.5}, aerodynamic diameter < 2.5 μm; PM₁₀, < 10 μm; PM_{coarse}, 2.5 μm ≤ diameter < 10 μm), black carbon (PM_{2.5} absorbance), that is, soot, and nitrogen oxides (NO₂ and NO_x) were measured (Cyrus et al. 2012; Eeftens et al. 2012b). In each study area, pollutants were measured in different locations for 2 weeks during winter, summer, and an intermediate season; that is, 6 weeks altogether in an approximately 1-year period (starting from autumn 2008 for KORA, HNR, and SAPALDIA; from winter 2008 for TwinGene and 60-year-olds study; and from winter 2010 for FINRISK). At the same time, pollutants were measured continuously for 1 year at one regional or urban background site in each study area as described by Eeftens et al. (2012b). Based on these measurements, annual outdoor concentrations of air pollution were modeled at the participants’ home addresses using land use regression (LUR) models (Beelen et al. 2013; Eeftens et al. 2012a). The models were developed for each cohort based on a common manual (ESCAPE 2008). A large number of potential predictors of air pollution concentrations derived from geographic information systems (e.g., traffic intensity, population density, forms of land use) were tested in the models with an aim to maximize explained variability. In SAPALDIA, models for PM were only available for one study area (Lugano, Switzerland). Measurements of air pollution for modeling purposes were conducted later (2008–2011) than the collection of blood samples (1994–2008). Therefore, historical air pollution concentrations were extrapolated for sensitivity analyses using data from routine

monitoring network site(s) as explained by Beelen et al. (2014).

In addition to modeled air pollution concentration, two indices of exposure to traffic were used: traffic intensity (vehicles/day) at the road nearest to the participant’s residence, and traffic load on major roads within 100 m of the residence (measured as the product of traffic intensity and the length of major road fragments within a 100-m buffer). Databases on traffic intensity were constructed for each cohort based on measurements conducted by county- or municipal-level authorities (Eeftens et al. 2012a). The traffic indices were always included in the statistical models together with background NO₂ concentrations. Information on the traffic intensity at the nearest road was not available for the HNR cohort.

Data analyses. One blood sample per person was available for the analyses, taken at the baseline visit except for SAPALDIA (first follow-up visit). High-sensitivity CRP and fibrinogen on continuous scales were used for the current cross-sectional analyses. Based on the distribution of inflammatory blood marker concentrations and model residuals, the (natural) logarithmic value of CRP concentrations was used in the analyses, whereas fibrinogen concentrations were not transformed. Linear regression models in SAS (SAS Institute Inc.) and STATA (StataCorp) software were used for the analyses. In the meta-analyses, cohort-specific effects were treated as having a random component apart from sampling error, and pooled using the empirical Bayes method. Heterogeneity of the cohort-specific estimates was estimated using Cochran’s Q test. Effect estimates are presented as percent difference from the outcome mean together with 95% confidence interval (CI), and are calculated for an increase of 5 μg/m³ in PM_{2.5} and PM_{coarse}, 10 μg/m³ in PM₁₀ and NO₂, 20 μg/m³ in NO_x, and 1 × 10⁻⁵/m in PM_{2.5} absorbance. An association was deemed present when it was statistically significant (*p* < 0.05).

Confounders in the statistical models were selected *a priori* based on literature on the determinants of CRP and fibrinogen concentrations. Four different regression models were

Table 1. Descriptive statistics for the study cohorts.

Study cohort (country)	Time of baseline visit	Age ^a		Sex ^a	BMI ^a	Smoking ^a	C-reactive protein		Fibrinogen	
		Range (years)	Average (years)	Female (%)	Average (kg/m ²)	Current (%)	<i>n</i> of samples	Average ± SD (mg/L)	<i>n</i> of samples	Average ± SD (g/L)
KORA (Germany)	1994–1995, 2000–2001	25–74	50.2	50.1	27.2	23.9	7,137	2.7 ± 4.9	7,151	2.76 ± 0.67
HNR (Germany)	2000–2003	45–76	59.5	49.9	27.9	23.3	4,492	3.1 ± 9.0	4,444	3.3 ± 0.8
SAPALDIA (Switzerland)	2002	18–60	53.8	52.7	25.5	27.7	1,688	2.1 ± 3.2	NA	NA
FINRISK (Finland)	1997, 2002, 2007	25–64	48.9	52.4	26.6	28.3	7,627	2.3 ± 4.5	2,044	3.55 ± 0.81
TwinGene (Sweden)	2004–2008	47–90	64.2	56.2	25.2	19.1	1,617	2.8 ± 4.9	NA	NA
60-year-olds (Sweden)	1997–1999	60	60.4	52	26.8	20.8	NA	NA	3,789	3.02 ± 0.77

Abbreviations: BMI, body mass index; NA, not available.

^aNumbers are based on participants with CRP measurements (except for 60-year-olds cohort). Descriptive statistics were similar when limited to participants with fibrinogen data (data not shown).

used to calculate the effect estimates. All these models included, in addition to exposure variables, an indicator variable for baseline visit for FINRISK and KORA, which had more than one recruitment period, and an area indicator for FINRISK, which consisted of two clearly separate study areas. Additional variables controlled for in the first regression model (the crude model) were age and sex. The second model (the main model) also included education (primary school or less, up to secondary school or equivalent, university degree), body mass index, smoking status (current smoker, ex-smoker, never smoker), physical activity (≤ 3 times/month, once a week, 2–3 times/week, > 3 times/week), and alcohol intake (number of drinks per week; 0, 1–3, 4–6, > 6). Effect estimates were also calculated using an extended model, which included additional covariates not available in all cohorts (alcohol intake excluding wine, wine intake, household income, fish intake, fruit intake, meat intake, CVD, diabetes, arthritis, ulcerative colitis, passive smoking).

The fourth model was a mixed model that included the same covariates as the main model, but also included neighborhood level as a random effect to take into account possible spatial clustering of observations. Depending on data availability for each cohort, neighborhoods either were defined as grids or were based on ZIP code areas. We assumed that the effect of neighborhood on health was mainly attributable to area level socioeconomic status. Therefore, area level median household income or unemployment rate were included in the model as fixed effects.

Effect modification by obesity (cutpoint, 30 kg/m² for body mass index) and sex was evaluated by adding an interaction term in the main model. In addition, a number of sensitivity analyses were conducted using the main model: exclusion of participants with high (> 10 mg/L) CRP values that may indicate a current bacterial infection, exclusion of study participants who moved residence during the 2 years before blood draw (information not available for KORA and for HNR participants from Bochum, Germany; see Supplemental Material, “Cohort descriptions”), use of

back-extrapolated concentrations of NO_x, NO₂, and PM₁₀ (in cohorts where changes over time in concentrations had occurred), inclusion of indicators for medication use (statins and antihypertensives), and finally inclusion of traffic noise for a subset of data in each center [data available only for larger population agglomerates as defined by the European Union (EU), no data available for SAPALDIA]. Traffic noise was modeled according to the guidelines of the EU noise directive (EU 2002). When using meta-analyses to generate combined effect estimates for sensitivity analyses, we substituted effect estimates from the main model for cohorts missing data for the sensitivity analyses.

Results

In the participating cohorts, the average age of the participants ranged from 49 to 64 years, and the proportion of current smokers ranged from 19 to 28% (Table 1). All cohorts consisted of approximately 50% women. There were altogether 22,561 participants with measures of CRP, and 17,428 samples of fibrinogen. Consistently across the cohorts, the standard deviation was much larger for CRP than for fibrinogen.

Modeled air pollution concentrations were the lowest in the Northern cohorts, in Finland and Sweden (Table 2). The highest traffic loads were observed near (within 100 m) the homes of the HNR and SAPALDIA study participants, and the lowest in the KORA cohort.

In the pooled analyses using the main model, CRP serum levels were positively associated with annual NO_x concentration (3.2%; 95% CI: 0.3, 6.1 per 20- $\mu\text{g}/\text{m}^3$ increase in NO_x) and traffic intensity at the nearest road (10.2%; 95% CI: 2.4, 18.6 for the highest category, i.e., $> 10,000$ vehicles/day; reference category $< 1,000$ vehicles/day) without evidence of significant heterogeneity between ESCAPE cohorts (Table 3, Figure 1). Categories of traffic load within 100 m from residence were not consistently associated with CRP. Inclusion of an extended set of individual level confounders (Table 3, Extended model) had only minor effects on

the estimates. When area-level confounding was taken into account, the effect estimate for NO_x decreased and was no longer statistically significant (2.5%; 95% CI: -0.6, 5.6) (Table 3, Area-level model). In contrast, effect estimates for traffic intensity at the nearest road were not affected. Particulate air pollution, including black carbon, and NO₂ were not associated with CRP in the pooled analyses. However, in cohort-specific analyses, high effect estimates were observed for PM_{2.5} in the German cohorts: statistically nonsignificant for KORA (12.9%; 95% CI: -1.4, 29.2%; per 5- $\mu\text{g}/\text{m}^3$ increase in PM_{2.5}), but significant for HNR (16.1%; 95% CI: 0.5, 34.1%) (see Supplemental Material, Figure S1).

In the pooled analyses for fibrinogen, borderline significant ($p < 0.1$) positive associations were observed for NO_x and traffic load within 100 m (in the two highest categories) without evidence of heterogeneity (Table 3, Figure 1). Slightly decreased effect estimates for traffic load were observed when the extended model was used (Table 3). Adding area-level confounders in the models did not affect the results. Other air pollutants were not associated with fibrinogen. However, in cohort-specific analyses, high effect estimates were observed for NO₂ in the German cohorts: statistically significant for KORA (1.6%; 95% CI: 0.5, 2.7% per 10- $\mu\text{g}/\text{m}^3$ increase in NO₂), but nonsignificant for HNR (1.2%; 95% CI: -0.2, 2.7%) (see Supplemental Material, Figure S2). Finally, the middle category of traffic intensity was positively associated with fibrinogen in the pooled analyses, but there was no evidence of association for the other categories.

Estimated effects of air pollution or traffic were not modified by obesity or sex: The confidence intervals were overlapping in the pooled analyses, and no consistent patterns were observed in the cohort-specific analyses (data not shown). Inclusion of environmental noise in the main model for CRP increased effect estimates for traffic intensity in the two highest categories (8.2%; 95% CI: 1.0, 15.9, and 11.9%; 95% CI: 3.4, 21.2, respectively) (see Supplemental Material, Table S1). Inclusion of information on statin use in the model and

Table 2. Descriptive statistics for modeled annual residential air pollution concentrations and indices of traffic intensity.

Study cohort	PM _{2.5} ($\mu\text{g}/\text{m}^3$)			PM ₁₀ ($\mu\text{g}/\text{m}^3$)			PM _{coarse} ($\mu\text{g}/\text{m}^3$)			PM _{2.5} absorbance ($1 \times 10^{-5}/\text{m}$)			NO ₂ ($\mu\text{g}/\text{m}^3$)			NO _x ($\mu\text{g}/\text{m}^3$)			Traffic, nearest road ^a (vehicles/day)		Traffic, within 100 m ^b (vehicles/day \times m)	
	Avg	5th Pctl	95th Pctl	Avg	5th Pctl	95th Pctl	Avg	5th Pctl	95th Pctl	Avg	5th Pctl	95th Pctl	Avg	5th Pctl	95th Pctl	Avg	5th Pctl	95th Pctl	Avg	95th Pctl	Avg	95th Pctl
KORA	13.6	12.5	15.3	20.3	16.5	24.3	6.2	4.9	8.4	1.7	1.5	2	18.7	13.7	25.9	32.6	23.8	47.1	1,600	8,500	434,000	2,790,000
HNR	18.4	16.9	20.4	27.8	25.3	31.6	10	7.3	12.4	1.6	1.2	2.2	30.3	23.4	38.7	50.9	33.6	71.9	NA	NA	1,124,000	4,432,000
SAPALDIA	17.1	14.7	19.9	23.7	19.8	27.4	6.7	4.9	8.6	2	1.3	2.5	27.8	17.1	37.9	46.7	24.0	69.5	3,520	15,500	945,000	4,170,000
FINRISK	7.7	5.6	9.1	14.1	9.5	19.9	6.7	3.9	11	1.3	1.1	1.6	15.4	8.7	24.2	24.4	13.6	41.2	1,700	9,600	638,000	3,750,000
TwinGene	7.2	4.8	9.2	14.8	6.5	20.5	7.2	1.0	11.8	0.6	0.4	0.9	10.8	6.7	19.9	18.7	11.4	39.4	1,500	6,400	570,000	3,300,000
60-year-olds	7.3	4.9	9.2	15.0	6.8	20.6	7.3	1.3	11.8	0.6	0.4	1	10.7	6.5	20.3	18.6	12.0	39.3	1,400	6,300	519,000	3,090,000

Abbreviations: Avg, average; NA, not available; Pctl, percentile.

^aTraffic intensity on the road nearest to the residence. ^bTraffic load on major roads within 100 m from the residence.

exclusion of persons who had recently moved led to decreased estimated effects of NO_x on CRP (2.5%; 95% CI: -0.4, 5.4, and 2.3%; 95% CI: -1.0, 5.8, respectively).

After exclusion of unusually high CRP values (237 observations for KORA, 181 for HNR, 47 for SAPALDIA, 257 for FINRISK, and 62 for TwinGene), effect estimates increased for PM_{coarse} (from 3.0%; 95% CI: -0.7, 6.8 to 6.4%; 95% CI: -0.4, 13.5% for a 5-µg/m³ increase). There were only minor changes in the effect estimates, when back-extrapolated air pollution levels were used in the models for CRP and fibrinogen (data not shown). In general, sensitivity analyses had only minor effects on the effect estimates for fibrinogen (see Supplemental Material, Table S1).

Discussion

In this European multi-center study, modeled annual particulate air pollution concentrations were not associated with systemic

inflammation. However, high traffic intensity on the road nearest to home was associated with increased levels of high-sensitivity CRP, an established marker for systemic inflammation. In the main analyses, long-term exposure to NO_x, an indicator for vehicle exhaust, was also associated with CRP serum levels. However, the effect estimate was more sensitive to further adjustments in the model, such as inclusion of area-level confounders. Fibrinogen was not consistently associated with air pollution, and there was only limited support for a positive association with traffic exposure.

CRP is an acute-phase reactant that has been shown in many cohort studies to be a reliable measure of underlying systemic inflammation and a predictor of future myocardial infarction and stroke (Danesh et al. 2004; Emerging Risk Factors Collaboration et al. 2010). In the present study, NO_x was the only pollutant that was positively associated with CRP in the analyses performed with

the *a priori*-selected main model. However, the association between NO_x and CRP was somewhat sensitive to adjustments in the models, such as inclusion of area-level confounders or information on statin use or moving history. Therefore, further studies are needed to confirm the suggestive association. Furthermore, it is possible that nitrogen oxides are not causally related to systemic inflammation, but that they rather represent markers for traffic-related air pollution mixture, including ultrafine particles, volatile organic compounds, and traffic-related PM_{2.5}.

In the present study, no evidence of an effect was found for soot, measured as PM_{2.5} absorbance, which is considered to be a key component of the near-road traffic-related pollution mixture. One reason for this might be that soot is stable in the atmosphere and is therefore associated not only with fresh but also with aged vehicle exhaust, which may be less toxic due to secondary atmospheric reactions. On the other hand, NO_x is a better

Table 3. Associations of air pollution and indices of traffic intensity with CRP and fibrinogen in the pooled analyses.

Exposures	Crude model ^a		Main model ^b		Extended model ^c		Area-level model ^d	
	% Diff ^e (95% CI)	ρ_{Heter}^f	% Diff ^e (95% CI)	ρ_{Heter}^f	% Diff ^e (95% CI)	ρ_{Heter}^f	% Diff ^e (95% CI)	ρ_{Heter}^f
CRP								
PM _{2.5}	9.4 (-9.7, 32.5)	0.00	2.4 (-7.5, 13.4)	0.05	2.3 (-6.4, 11.7)	0.11	-0.3 (-9.0, 9.2)	0.14
PM ₁₀	2.2 (-7.6, 12.9)	0.05	1.2 (-3.8, 6.4)	0.90	0.7 (-4.3, 5.9)	0.94	0.2 (-4.9, 5.6)	0.76
PM _{coarse}	2.8 (-2.3, 8.2)	0.21	3.0 (-0.7, 6.8)	0.53	2.4 (-1.2, 6.1)	0.66	2.4 (-1.6, 6.5)	0.35
PM _{2.5} absorbance	3.2 (-6.9, 14.4)	0.01	0.4 (-5.2, 6.4)	0.73	0.0 (-5.5, 5.9)	0.86	-2.5 (-8.4, 3.6)	0.78
NO ₂	2.2 (-4.3, 9.2)	0.00	2.1 (-0.8, 5.1)	0.72	2.0 (-0.9, 4.9)	0.82	1.4 (-1.7, 4.6)	0.70
NO _x	3.2 (-2.5, 9.2)	0.00	3.2 (0.3, 6.1)**	0.95	2.9 (0.1, 5.9)**	0.99	2.5 (-0.6, 5.6)	0.94
Traffic intensity at the nearest road (vehicles/day)								
< 1,000	Reference		Reference		Reference		Reference	
1,000–5,000	-0.8 (-5.9, 4.5)	0.81	-1.7 (-6.3, 3.2)	0.69	-2.0 (-6.6, 2.8)	0.81	-1.6 (-6.2, 3.3)	0.66
5,000–10,000	6.3 (-0.9, 14.1)*	0.57	5.5 (-1.2, 12.6)	0.42	4.5 (-2.0, 11.6)	0.55	5.1 (-2.7, 13.6)	0.35
> 10,000	12.8 (4.2, 22.2)**	0.84	10.2 (2.4, 18.6)**	0.77	9.8 (2.1, 18.2)**	0.84	10.2 (2.4, 18.6)**	0.76
Traffic load within 100 m on major roads (vehicles/day × m)								
< 500,000	Reference		Reference		Reference		Reference	
500,000–1,500,000	0.2 (-7.9, 9.0)	0.15	1.4 (-3.0, 6.0)	0.41	1.1 (-3.3, 5.7)	0.59	1.3 (-3.3, 6.1)	0.39
1,500,000–3,000,000	3.6 (-1.6, 9.1)	0.66	1.8 (-3.0, 6.8)	0.51	1.6 (-3.1, 6.5)	0.72	1.5 (-3.3, 6.5)	0.41
> 3,000,000	5.3 (-1.8, 13.0)	0.23	2.6 (-3.4, 9.0)	0.29	2.0 (-3.9, 8.3)	0.30	2.1 (-3.7, 8.3)	0.32
Fibrinogen								
PM _{2.5}	2.8 (0.3, 5.3)**	0.09	0.5 (-1.1, 2.0)	0.66	0.2 (-1.3, 1.8)	0.77	0.4 (-1.3, 2.0)	0.96
PM ₁₀	0.5 (-2.3, 3.2)	0.01	0.1 (-1.4, 1.8)	0.18	0.0 (-1.4, 1.4)	0.27	-0.2 (-1.4, 1.0)	0.39
PM _{coarse}	-0.0 (-1.7, 1.6)	0.03	-0.1 (-1.2, 0.9)	0.24	-0.3 (-1.2, 0.6)	0.37	-0.5 (-1.4, 0.4)	0.80
PM _{2.5} absorbance	1.5 (-0.4, 3.5)	0.25	0.3 (-1.2, 1.9)	0.36	0.1 (-1.3, 1.6)	0.49	-0.5 (-2.0, 1.1)	0.61
NO ₂	0.8 (-1.2, 2.8)	0.00	0.5 (-0.8, 1.8)	0.04	0.5 (-0.7, 1.6)	0.10	0.4 (-0.6, 1.3)	0.30
NO _x	0.9 (-0.9, 2.7)	0.00	0.7 (-0.4, 1.8)	0.08	0.6 (-0.5, 1.6)	0.13	0.5 (-0.4, 1.3)	0.31
Traffic intensity at the nearest road (vehicles/day)								
< 1,000	Reference		Reference		Reference		Reference	
1,000–5,000	0.5 (-0.7, 1.8)	0.43	0.2 (-1.5, 1.8)	0.28	0.0 (-1.2, 1.3)	0.48	0.1 (-1.4, 1.7)	0.29
5,000–10,000	1.8 (0.0, 3.6)*	0.93	1.8 (0.1, 3.5)**	0.99	1.6 (-0.1, 3.3)*	0.93	1.9 (0.1, 3.6)**	0.99
> 10,000	1.3 (-0.7, 3.3)	0.40	0.9 (-1.0, 2.8)	0.84	0.6 (-1.3, 2.5)	0.75	0.9 (-1.1, 2.8)	0.85
Traffic load within 100 m on major roads (vehicles/day × m)								
< 500,000	Reference		Reference		Reference		Reference	
500,000–1,500,000	0.2 (-1.3, 1.7)	0.18	0.3 (-0.8, 1.4)	0.39	0.1 (-0.9, 1.2)	0.37	0.2 (-1.0, 1.4)	0.32
1,500,000–3,000,000	1.4 (0.2, 2.7)**	0.87	1.0 (-0.2, 2.2)*	0.87	0.9 (-0.3, 2.1)	0.86	1.0 (-0.2, 2.2)	0.86
> 3,000,000	1.9 (0.3, 3.5)**	0.35	1.2 (-0.2, 2.6)*	0.59	1.1 (-0.2, 2.5)	0.64	1.2 (-0.2, 2.6)*	0.57

Effect estimates were calculated for a change of 5 µg/m³ in PM_{2.5} and PM_{coarse}, 10 µg/m³ in PM₁₀ and NO₂, 20 µg/m³ in NO_x, and 1 × 10⁻⁵/m in PM_{2.5} absorbance.

^aModel was adjusted for age, sex, baseline visit (KORA and FINRISK), study area (FINRISK). ^bModel was adjusted for age, sex, baseline visit (KORA and FINRISK), study area (FINRISK), education, body mass index, smoking status, physical activity, alcohol intake. ^cModel was adjusted for age, sex, baseline visit (KORA and FINRISK), study area (FINRISK), education, body mass index, smoking status, physical activity in all cohorts, and for wine intake, intake of other types of alcohol, household income, fish intake, fruit intake, meat intake, CVD, diabetes, arthritis, colitis ulcerosa, passive smoking in cohorts where the data available. ^dMixed model (neighborhood as a random effect) was adjusted for age, sex, baseline visit (KORA and FINRISK), study area (FINRISK), education, body mass index, smoking status, physical activity, and area level household income or unemployment rate. ^ePercent difference. ^f ρ -Value for heterogeneity. *Effect estimates with *p*-values < 0.1. **Effect estimates with *p*-values < 0.05.

marker for vehicle exhausts than soot, which also has other (combustion) sources such as residential wood combustion (Healy et al. 2012; Janssen et al. 2012).

Another traffic-related pollutant is the coarse size fraction of particulate matter, which forms the major part of road dust. Road dust is rich in earth minerals (wear of road surface), transition metals (wear of brakes and clutch), and biological material (pollen fragments, bacteria). All these components have been shown in toxicological studies to have considerable proinflammatory potential (Camatini et al. 2012; Jalava et al. 2009; Schwarze et al. 2007). In the present study, effect estimates for coarse particles were consistently positive, but statistical significance was not reached in the main analyses. This could be attributable partly to exposure misclassification: LUR models predicted less of the variance of outdoor concentrations of PM_{coarse} (average R^2 of models = 71%) than of NO_x ($R^2 = 86%$) (Beelen et al. 2013; Eeftens et al. 2012a).

In the pooled analyses, markers of inflammation were not associated with $PM_{2.5}$ or PM_{10} , the spatial variation of which is less strongly associated with road traffic than that of NO_x (Eeftens et al. 2012b). It can be speculated that $PM_{2.5}$ and PM_{10} contain some components that are less toxic than pollution mixture near traffic. Only a few epidemiological studies have previously evaluated associations between long-term exposure to air pollution and CRP. In a cross-sectional study conducted in a representative sample of the English population (Forbes et al. 2009), no association was observed between CRP and PM_{10} or NO_2 . Fibrinogen was not associated with air pollution either. In the study, dispersion models were used to estimate air pollution concentrations for the centroid (1 km²) of each postcode sector; the same exposure estimate was assigned to all study participants within the sector. The spatial resolution of exposure assessment was higher in the present study, which might explain the discrepancy concerning CRP. In a German study that used the HNR cohort and dispersion models, $PM_{2.5}$ was found to be associated with CRP in men but not in women (Hoffmann et al. 2009). In the present study, $PM_{2.5}$ was associated with CRP in the HNR cohort in men and women combined. In an earlier cross-sectional study conducted in Stockholm, Sweden (Panasevich et al. 2009), there was some evidence for an association between dispersion-modeled long-term NO_2 concentration and CRP, but not for fibrinogen, among adults (45–70 years of age). In the same study, NO_2 was also associated with another marker of systemic inflammation, interleukin-6. In a cross-sectional study conducted in a representative sample of adults ≥ 54 years of age in Taiwan

(Chuang et al. 2011), long-term PM_{10} , $PM_{2.5}$, and NO_2 concentrations were all associated with interleukin-6.

Fibrinogen is a hemostatic factor with proinflammatory properties. It has been shown that inflammatory and coagulatory processes are intertwined via many molecular mechanisms, and typically systemic inflammation is associated with increased levels of fibrinogen (Davalos and Akassoglou 2012). One previous cross-sectional study, conducted in a representative sample of the U.S. population (Schwartz 2001), had data on fibrinogen but not on CRP. Both NO_2 and PM_{10} were found to be associated with fibrinogen as well as with platelet counts. In the present study, fibrinogen was not associated with the studied pollutants. Compared with CRP, there were fewer observations (17,428 vs. 22,561) and lower variability in concentrations—both factors that reduce the chance of identifying associations.

The most robust association in the present study was observed between traffic intensity at the road nearest to home and CRP. Cohort participants living on a busy road ($> 10,000$ vehicles/day) had significantly increased CRP compared with participants living on a quiet residential street ($< 1,000$ vehicles/day). The estimated increase was modest (10.2%), but can be seen as clinically significant considering that CVD risk seems to increase linearly

with increasing CRP (Ridker and Cook 2004). It is noteworthy that increased CRP concentration is not necessarily a causal factor in the development of CVD, but might only be an indicator of an increased risk. No statistically significant association was observed between CRP levels and traffic load within 100 m on major roads, another indicator of proximity to dense traffic, perhaps because traffic intensity at the nearest road indicates better exposure to fresh traffic exhaust. An earlier cross-sectional study conducted among adult Puerto Ricans (45–75 years of age) has reported an association between living within 100 m from busy roads ($> 20,000$ vehicles/day) and CRP (Rioux et al. 2010). In the present study, positive but statistically nonsignificant effect estimates were observed for high traffic load.

Inhalation of particulate air pollution is known to cause pulmonary oxidative stress and pulmonary inflammation (Brook et al. 2010), which in turn has been hypothesized to lead further to systemic inflammation. It has been recently shown that oxidative potential of particles sampled near major streets is several times higher than of particles at urban background locations (Boogaard et al. 2012). Systemic inflammation and impaired endothelial function provide a plausible link between exposure to traffic-related pollutants and CVD. Systemic inflammation plays




Figure 1. Cohort-specific and pooled effect estimates (the main model) for the associations of NO_x and traffic with inflammatory markers. veh, vehicles. Estimates calculated for a $20\text{-}\mu\text{g}/\text{m}^3$ increase in NO_x . (A) NO_x , traffic intensity at the nearest road, and CRP; (B) NO_x , traffic load on major roads within 100 m, and fibrinogen.

a major role in the development of atherosclerosis, possibly leading, over time, to clinically detectable CVD (Libby et al. 2002). Acute increases in the inflammatory response and the development of vulnerable arterial plaque may in turn lead to acute exacerbations of CVD. An increasing number of studies link either traffic-related pollutants or traffic intensity with the prevalence, and even incidence, of CVD (Brook et al. 2010).

Most of the evidence on the health effects of PM relates to cardiorespiratory health. However, some recent studies have associated long-term PM exposure with additional outcomes such as rheumatoid arthritis, diabetes, cognitive impairment, and leukemia (Amigou et al. 2011; Hart et al. 2009; Krämer et al. 2010; Ranft et al. 2009). Importantly, an inflammatory component is part of the pathogenesis of many major noncommunicable diseases such as diabetes, Alzheimer's disease, and cancer (Mantovani et al. 2008; Sardi et al. 2011). This adds to the importance of evaluating links between exposure to air pollution from local traffic and systemic inflammation.

As part of the large European ESCAPE project, this study has some major strengths, including the large number of observations, a multi-city study design covering many types of exposure environments and climates, rigorous exposure assessment including extensive quality control measures, and the use of common study manuals for exposure assessment, data management, and data analyses. An obvious limitation is the cross-sectional nature of the study. However, a large number of individual-level and some neighborhood-level covariates were available for inclusion in the statistical models to reduce confounding. It was unfortunate that data on individual-level income, a quantitative indicator of socioeconomic status, was not available. However, we believe that residual confounding by individual level socioeconomic status would have been substantially reduced by adjusting for educational level and lifestyle factors including smoking. The association between traffic intensity at the road nearest to home and CRP was not sensitive to model specifications, which increases the plausibility of our main finding.

Other limitations include the retrospective nature of exposure assessment, varying LUR model performance for different air pollution components, and lack of data on personal exposures. Several recent studies have demonstrated that spatial variation in air pollution exposure can be reliably estimated also retrospectively, even when overall mean levels may change over time (Eeftens et al. 2011; Gulliver et al. 2013). However, the differences in the accuracy of exposure assessment among pollutants should be taken into account when interpreting the present

results. Therefore, even though no statistically significant associations were observed either for PM_{2.5} or coarse particles, the apparent lack of association for the latter could be attributable partly to the difficulty of predicting concentrations. Finally, there is evidence that living close to traffic contributes significantly to personal exposures (Van Roosbroeck et al. 2008). Therefore, indices of traffic intensity near home can be considered as reasonable proxies for exposure to traffic-related air pollution. This is especially true for traffic intensity on the road nearest to home, because emissions occur close to exposed persons, so more complex physicochemical dispersion processes do not affect the concentrations.

Conclusions

Particulate air pollution was not associated with two markers of systemic inflammation in the present analysis. However, compared with levels in other participants, CRP levels were higher among those living next to a busy road. Therefore, the study provides some support for a mechanistic link between proximity to traffic and deteriorating cardiovascular health. Future research should aim at identifying air pollution components responsible for the association between traffic intensity and systemic inflammation.

REFERENCES

- Amigou A, Sermage-Faure C, Orsi L, Leverger G, Baruchel A, Bertrand Y, et al. 2011. Road traffic and childhood leukemia: the ESCALE study (SFCE). *Environ Health Perspect* 119:566–572; doi:10.1289/ehp.1002429.
- Beelen R, Hoek G, Vienneau D, Eeftens M, Dimakopoulou K, Pedeli X, et al. 2013. Development of NO₂ and NO_x land use regression models for estimating air pollution exposure in 36 study areas in Europe—the ESCAPE project. *Atmos Environ* 72:10–23.
- Beelen R, Raaschou-Nielsen O, Stafoggia M, Andersen ZJ, Weinmayr G, Hoffmann B, et al. 2014. Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the multicentre ESCAPE project. *Lancet* 383:785–795.
- Boogaard H, Janssen NA, Fischer PH, Kos GP, Weijsers EP, Cassee FR, et al. 2012. Contrasts in oxidative potential and other particulate matter characteristics collected near major streets and background locations. *Environ Health Perspect* 120:185–191; doi:10.1289/ehp.1103667.
- Brook RD, Rajagopalan S, Pope CA III, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 121:2331–2378.
- Brunekreef B, Forsberg B. 2005. Epidemiological evidence of effects of coarse airborne particles on health. *Eur Respir J* 26:309–318.
- Camatini M, Corvaja V, Pezzolato E, Mantecca P, Gualtieri M. 2012. PM₁₀-biogenic fraction drives the seasonal variation of proinflammatory response in A549 cells. *Environ Toxicol* 27:63–73.
- Chuang KJ, Yan YH, Chiu SY, Cheng TJ. 2011. Long-term air pollution exposure and risk factors for cardiovascular diseases among the elderly in Taiwan. *Occup Environ Med* 68:64–68.
- Cyrys J, Eeftens M, Heinrich J, Ampe C, Armengaud A, Beelen R, et al. 2012. Variation of NO₂ and NO_x concentrations between and within 36 European study areas: results from the ESCAPE study. *Atmos Environ* 62:374–390.
- Danesh J, Wheeler JG, Hirschfield GM, Eda S, Eiriksdottir G, Rumley A, et al. 2004. C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *N Engl J Med* 350:1387–1397.
- Davalos D, Akassoglou K. 2012. Fibrinogen as a key regulator of inflammation in disease. *Semin Immunopathol* 34:43–62.
- Eeftens M, Beelen R, de Hoogh K, Bellander T, Cesaroni G, Cirach M, et al. 2012a. Development of land use regression models for PM_{2.5}, PM_{2.5} absorbance, PM₁₀ and PM_{coarse} in 20 European study areas; results of the ESCAPE project. *Environ Sci Technol* 46:11195–11205.
- Eeftens M, Beelen R, Fischer P, Brunekreef B, Meliefste K, Hoek G. 2011. Stability of measured and modelled spatial contrasts in NO₂ over time. *Occup Environ Med* 68:765–770.
- Eeftens M, Tsai MY, Ampe C, Anwander B, Beelen R, Bellander T, et al. 2012b. Spatial variation of PM_{2.5}, PM₁₀, PM_{2.5} absorbance and PM_{coarse} concentrations between and within 20 European study areas and the relationship with NO₂—results of the ESCAPE project. *Atmos Environ* 62:303–317.
- Emerging Risk Factors Collaboration, Kaptoge S, Di Angelantonio E, Lowe G, Pepys MB, Thompson SG, et al. 2010. C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet* 375:132–140.
- ESCAPE (European Study of Cohorts for Air Pollution Effects). 2008. European Study of Cohorts for Air Pollution Effects (ESCAPE): ESCAPE Study Manual. Available: <http://www.escapeproject.eu/manuals/> [accessed 25 February 2015].
- EU (European Union). 2002. Directive 2002/49/EC of the European Parliament and of the Council of 25 June 2002 relating to the Assessment and Management of Environmental Noise. Available: <http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32002L0049&from=EN> [accessed 26 June 2015].
- Forbes LJ, Patel MD, Rudnicka AR, Cook DG, Bush T, Stedman JR, et al. 2009. Chronic exposure to outdoor air pollution and markers of systemic inflammation. *Epidemiology* 20:245–253.
- Gulliver J, de Hoogh K, Hansell A, Vienneau D. 2013. Development and back-extrapolation of NO₂ land use regression models for historic exposure assessment in Great Britain. *Environ Sci Technol* 47:7804–7811.
- Hart JE, Laden F, Puett RC, Costenbader KH, Karlson EW. 2009. Exposure to traffic pollution and increased risk of rheumatoid arthritis. *Environ Health Perspect* 117:1065–1069; doi:10.1289/ehp.0800503.
- Healy RM, Sciare J, Poulain L, Kamili K, Merkel M, Müller T, et al. 2012. Sources and mixing state of size-resolved elemental carbon particles in a European megacity: Paris. *Atmos Chem Phys* 12:1681–1700.
- Hoffmann B, Moebus S, Dragano N, Stang A, Möhlenkamp S, Schmermund A, et al. 2009. Chronic residential exposure to particulate matter air pollution and systemic inflammatory markers. *Environ Health Perspect* 117:1302–1308; doi:10.1289/ehp.0800362.
- Jalava PI, Hirvonen MR, Sillanpää M, Pennanen AS, Happonen MS, Hillamo R, et al. 2009. Associations of

- urban air particulate composition with inflammatory and cytotoxic responses in RAW 246.7 cell line. *Inhal Toxicol* 21:994–1006.
- Janssen N, Gerlofs-Nijland ME, Lanki T, Salonen RO, Cassee F, Hoek G, et al. 2012. Health Effects of Black Carbon. Bonn:World Health Organization, Regional Office for Europe. Available: http://www.euro.who.int/__data/assets/pdf_file/0004/162535/e96541.pdf [accessed 26 June 2015].
- Krämer U, Herder C, Sugiri D, Strassburger K, Schikowski T, Ranft U, et al. 2010. Traffic-related air pollution and incident type 2 diabetes: results from the SALIA cohort study. *Environ Health Perspect* 118:1273–1279; doi:10.1289/ehp.0901689.
- Libby P, Ridker PM, Maseri A. 2002. Inflammation and atherosclerosis. *Circulation* 105:1135–1143.
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. 2012. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2224–2260.
- Mantovani A, Allavena P, Sica A, Balkwill F. 2008. Cancer-related inflammation. *Nature* 454:436–444.
- Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. 2012. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2197–2223.
- Nawrot TS, Perez L, Künzli N, Munters E, Nemery B. 2011. Public health importance of triggers of myocardial infarction: a comparative risk assessment. *Lancet* 377:732–740.
- Panasevich S, Leander K, Rosenlund M, Ljungman P, Bellander T, de Faire U, et al. 2009. Associations of long- and short-term air pollution exposure with markers of inflammation and coagulation in a population sample. *Occup Environ Med* 66:747–753.
- Ranft U, Schikowski T, Sugiri D, Krutmann J, Krämer U. 2009. Long-term exposure to traffic-related particulate matter impairs cognitive function in the elderly. *Environ Res* 109:1004–1011.
- Ridker PM, Cook N. 2004. Clinical usefulness of very high and very low levels of C-reactive protein across the full range of Framingham Risk Scores. *Circulation* 109:1955–1959.
- Rioux CL, Tucker KL, Mwamburi M, Gute DM, Cohen SA, Brugge D. 2010. Residential traffic exposure, pulse pressure, and C-reactive protein: consistency and contrast among exposure characterization methods. *Environ Health Perspect* 118:803–811.
- Sardi F, Fassina L, Venturini L, Inguscio M, Guerriero F, Rolfo E, et al. 2011. Alzheimer's disease, autoimmunity and inflammation. The good, the bad and the ugly. *Autoimmun Rev* 11:149–153.
- Schwartz J. 2001. Air pollution and blood markers of cardiovascular risk. *Environ Health Perspect* 109(suppl 3):405–409.
- Schwarze PE, Øvrevik J, Hetland RB, Becher R, Cassee FR, Låg M, et al. 2007. Importance of size and composition of particles for effects on cells in vitro. *Inhal Toxicol* 19(suppl 1):17–22.
- Van Roosbroeck S, Hoek G, Meliefste K, Janssen NA, Brunekreef B. 2008. Validity of residential traffic intensity as an estimate of long-term personal exposure to traffic-related air pollution among adults. *Environ Sci Technol* 42:1337–1344.