

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

**Photophysics and Photoredox Catalysis of a Homoleptic Re^{I}
Tris(diisocyanide) Complex**

Journal:	<i>Inorganic Chemistry</i>
Manuscript ID	ic-2017-03258d.R1
Manuscript Type:	Communication
Date Submitted by the Author:	12-Feb-2018
Complete List of Authors:	Larsen, Christopher; University of Basel, Department of Chemistry Wenger, Oliver; University of Basel, Department of Chemistry

SCHOLARONE™
Manuscripts

Photophysics and Photoredox Catalysis of a Homoleptic Re^I Tris(diisocyanide) Complex

Christopher B. Larsen and Oliver S. Wenger*

Department of Chemistry, University of Basel, St. Johannis-Ring 19, Basel CH-4056, Switzerland

Supporting Information Placeholder

ABSTRACT: Herein is communicated a homoleptic Re^I complex bearing three chelating diisocyanide ligands, and its photophysical properties. The complex emits weakly from a high energy ³MLCT excited state with an 8 ns lifetime in de-aerated CH₃CN at 22 °C, and is shown to act as an efficient photoredox catalyst comparable to [Ir(ppy)₃] (ppy = 2-phenylpyridine) in representative test reactions.

Photoredox catalysis is rapidly emerging as a useful method for C-C bond formations.¹⁻² d⁶ precious-metal complexes are the prototypical photoredox catalysts and have played a pivotal role in the development of photoredox catalysis as a field.³⁻⁴ Despite growing interest in organic and earth-abundant metal complex photoredox catalysts,⁵⁻⁸ d⁶ precious-metal complexes remain the first choice for many photoredox transformations – with [Ru(bpy)₃]²⁺ (bpy = 2,2'-bipyridine) and [Ir(ppy)₃] (ppy = 2-phenylpyridine) among the most popular.^{1-3, 9} This largely stems from their ease of synthesis, favourable photophysical properties, tunability, and robustness.

Interestingly, despite significant attention having been paid to the photoluminescent properties of [Re(CO)₃(α-diimine)(X)] complexes (X = Cl⁻, Br⁻, CN⁻, SCN⁻, py)¹⁰⁻¹⁶ and their performance in the catalytic photoreduction of CO₂,^{15, 17-24} Re^I complexes have been essentially unused as catalysts in organic photoredox transformations.

d⁶ metal complexes typically exhibit photoluminescent metal-to-ligand charge-transfer (MLCT) excited states.³ The energies of these excited states are easily tunable through the choice of metal and ligand design. A key feature of such complexes is that in their emissive ³MLCT excited states they are both better oxidants and reductants than in their electronic ground states. The magnitude by which the oxidation potential of a complex is altered between ground and excited state may be estimated using the Rehm-Weller equation:²⁵

$$E_{1/2}(C^+/C^*) \approx E_{1/2}(C^+/C^0) - E_{00}$$

Where $E_{1/2}(C^+/C^0)$ and $E_{1/2}(C^+/C^*)$ are the ground- and excited-state oxidation potentials, respectively, and E_{00} the energy of the electronic origin for the excited state (the difference in energy between the 0th vibrational levels of the

ground and excited states). With the goal of increasing the reducing power of a catalyst, and therefore increasing its substrate scope, the pertinent design principles are to combine an easily oxidized metal (to decrease $E_{1/2}(C^+/C^0)$) with difficult to reduce ligands (to increase E_{00} for the MLCT excited state). Using [Ru(bpy)₃]²⁺ and [Ir(ppy)₃] as examples, the metal-centred (Ru^{II} and Ir^{III}, respectively) ground-state oxidation potentials of these complexes are 1.29 and 0.77 V vs SCE, respectively, and the ligand-based (bpy vs ppy, respectively) reductions -1.33 and -2.19 V vs SCE, yielding E_{00} values of 2.10 and 2.50 eV, respectively.⁶ Inserting these values into the Rehm-Weller equation, it is unsurprising that [Ir(ppy)₃] is the more powerful photoreductant with $E_{1/2}(C^+/C^*)$ of -1.73 V vs SCE, compared to -0.81 V vs SCE for [Ru(bpy)₃]²⁺.

Our recent investigations into photoluminescent earth-abundant zero-valent d⁶ metal complexes have identified chelating diisocyanides as promising ligands for the development of strongly photoreducing complexes.²⁶⁻²⁸ These ligands are more difficult to reduce than bpy or ppy, and their strong σ-donor character makes metal-based oxidations more facile. In combination with their strong π-accepting character, these attributes also increase the energy of deactivating ligand-field states relative to desirable MLCT states. Furthermore, the strong-field character of isocyanides is akin to that of carbonyls,²⁹⁻³⁰ stabilizing the low-valent Re^I.³¹⁻³² Prior studies of Re^I complexes with monodentate isocyanide ligands demonstrated that this compound class has very promising photophysical properties,³³⁻³⁵ but analogues with chelating ligands seem to be unknown so far.


We have therefore prepared a homoleptic Re^I tris(diisocyanide) complex, [Re(L^I)₃]PF₆ (Scheme 1), with the goal of developing a strong photoreductant, anticipating that the chelating nature of the ligand would increase robustness compared to Re^I complexes with monodentate arylisocyanide ligands.²⁷ The relevant photophysical properties of the complex are presented, in combination with the photoredox catalytic performance for two representative test reactions.

The Re^I complexation of L^I was achieved through adaptation of the method of Wang *et al.*³⁶ for the synthesis of [Re(^{Me}bpy)₃]PF₆ from [ReCl₄(THF)₂] to afford [Re(L^I)₃]PF₆ in 76% yield (Scheme 1). L^I and [ReCl₄(THF)₂] were prepared according to literature procedures.^{26, 37}

Electronic absorption and emission spectra of [Re(L^I)₃]PF₆ in CH₃CN are presented in Figure 1, and in THF, CH₂Cl₂ and

CH₃CN in Figure S1. Both absorption and emission profiles exhibit only minor differences across this solvent range.

Scheme 1. Synthesis of [Re(L¹)₃]PF₆



The electronic absorption spectrum of [Re(L¹)₃]PF₆ in CH₃CN (blue trace) exhibits a maximum at 318 nm ($\epsilon = 34,000 \text{ M}^{-1} \text{ cm}^{-1}$) not present in the ligand spectrum (dotted black trace), which tails past 400 nm and is assigned as MLCT in nature. Upon photoexcitation into this MLCT absorption feature ($\lambda_{\text{ex}} = 350 \text{ nm}$) in de-aerated CH₃CN at 22 °C, photoluminescence is observed at ~480 nm (red trace in Fig. 1) with a quantum yield of 0.6% and lifetime of 8 ns (inset in Fig. 1). Photoluminescence is quenched in the presence of oxygen, and is therefore characterized as ³MLCT in nature. From the intersection of the absorption and photoluminescence spectra, a ³MLCT excited state energy of 2.96 eV is estimated.

The low quantum yield and short excited-state lifetime may be explained by thermal population of ligand-field excited states. These states are typically not readily accessible in [Re(CO)₃(α -diimine)(X)] complexes;^{10-16, 38} however, our complex has a higher energy ³MLCT state than [Re(CO)₃(α -diimine)(X)] complexes,^{16, 39} and the isocyanide ligands destabilize ligand-field states to a lesser degree than carbonyls, resulting in thermally accessible ligand-field states. A previously studied homoleptic Re^I hexakis(arylisocyanide) complex with monodentate ligands was non-emissive at room temperature due to thermal occupation of ligand-field states.⁴⁰ Thus, the chelating ligand leads to markedly improved photophysical properties.

Electrochemical studies performed on [ReCl(CO)₃(N-N)] complexes typically exhibit an irreversible Re^{II}/Re^I oxidation near the solvent limit,⁴¹ and for the [Re(2,6-dimethylphenylisocyanide)₆]⁺ complex a potential of 1.08 V vs. SSCE was reported.⁴⁰ [Re(L¹)₃]PF₆ exhibits such an oxidation at 1.36 V vs SCE in CH₃CN (Figure S3). In combination with the excited-state energy (see above), an estimated $E_{1/2}(\text{C}^+/\text{C}^*)$ of -1.60 V vs SCE can be obtained. [Re(L¹)₃]PF₆ is photostable in de-aerated CH₃CN, with its electronic absorption spectrum in deaerated CH₃CN remaining unchanged after irradiation at 405 nm (1 W) for 24 h (Figure S2). This is an important finding in view of the known photolability of the [Re(2,6-dimethylphenylisocyanide)₆]⁺ complex, which undergoes efficient ligand exchange in presence of halide anions.⁴²

Many organic photoredox catalysts have excited-state lifetimes that are similarly short as that of [Re(L¹)₃]PF₆,^{5, 43} and it therefore seemed reasonable to perform photoredox catalysis

experiments with this complex, particularly in view of its comparatively high reducing power and photostability.





Figure 1. Electronic absorption (blue trace) and emission (red trace, $\lambda_{\text{ex}} = 350 \text{ nm}$) spectra of [Re(L¹)₃]PF₆, recorded in de-aerated CH₃CN at 22 °C. Inset: Photoluminescence decay of [Re(L¹)₃]PF₆ recorded in de-aerated CH₃CN ($\lambda_{\text{ex}} = 405 \text{ nm}$). *Stray excitation light. Black dotted trace represents ligand UV-Vis spectrum in CH₃CN.

Our initial approach was to try and match the excited-state oxidation potential of the complex with the reduction potential of a known substrate of interest. To this end, we chose the dimerization of benzyl bromide ($E_{\text{red}}^{\circ} = -1.3 \text{ V}$ vs SCE)¹ as a test reaction (Table 1).⁴⁴ Reactions were performed in flame-sealed NMR tubes containing the substrate (0.1 M), catalyst (1 mol %) and *N,N*-diisopropylethylamine (DIPEA, 4 mol eq.) as a sacrificial electron donor in de-aerated CD₂Cl₂. Over the course of 20 h under 405 nm (1 W) irradiation at room temperature, [Re(L¹)₃]PF₆ completely converted the substrate, with 81% dimerizing to afford dibenzyl and 19% presumably abstracting an H-atom from trace water present in the solvent to afford toluene. Catalyst-based photoluminescence was still observable after consumption of the substrate with no obvious loss of intensity, indicating that the catalyst is stable over long reaction times. As a control experiment, [Ru(bpy)₃](PF₆)₂ ($E_{1/2}(\text{C}^+/\text{C}^*) = -0.72 \text{ V}$ vs SCE) converted only 49% of the substrate under 455 nm irradiation for the same duration at room temperature, yielding dibenzyl and toluene in 45% and 4% yields, respectively. The different excitation source was used to better coincide with the absorption maximum of the [Ru(bpy)₃](PF₆)₂ MLCT absorption band. No reaction occurred in the absence of either catalyst or light.


Encouraged by the catalytic performance of [Re(L¹)₃]PF₆ in this test reaction, we chose to explore the scope of the complex for more challenging catalytic photoreductive transformations. Given an $E_{1/2}(\text{C}^+/\text{C}^*)$ of -1.60 V vs SCE for [Re(L¹)₃]PF₆, the reduction of iodobenzene ($E_{\text{red}}^{\circ} \approx -1.6 \text{ V}$ vs SCE in DMSO)⁴⁵ to benzene was chosen as a demanding transformation (Table 2). Reactions were once again performed in flame-sealed NMR tubes with de-aerated solvent, using conditions modified from Nguyen *et al.* for [Ir(ppy)₃] (Table 2).⁴⁶

Table 1. Benzyl bromide dimerization.


Catalyst	Irradiation Wavelength / nm	Benzyl Bromide Consumption / %	Dibenzyl Formation / %	Toluene Formation / %
[Re(L') ₃]PF ₆	405	100	81	19
[Ru(bpy) ₃](PF ₆) ₂	455	49	45	4

Conditions: 20 h continuous irradiation in vacuum flame-sealed NMR tubes. Conversions obtained from ¹H NMR integrals. See the Supporting Information for details.

Over the course of 20 h under 405 nm irradiation at room temperature, both [Re(L')₃]PF₆ and [Ir(ppy)₃] catalyzed the reaction to completion. No reaction occurred in the absence of either catalyst or light. Next we attempted the even more demanding reduction of bromobenzene ($E^{\circ}_{\text{red}} \approx -2.1$ V vs SCE in DMSO)⁴⁵ to benzene. Under exactly the same conditions as for the reduction of iodobenzene to benzene, [Re(L')₃]PF₆ catalyzed the reaction to 22%, and [Ir(ppy)₃] to 18%. These values are within experimental error, indicating that despite the short excited-state lifetime, [Re(L')₃]PF₆ operates at a similar catalytic efficiency to [Ir(ppy)₃].

Table 2. Reduction of halobenzenes.


Catalyst	X	Conversion / %
[Re(L') ₃]PF ₆	I	100
[Ir(ppy) ₃]	I	100 ^a
[Re(L') ₃]PF ₆	Br	22
[Ir(ppy) ₃]	Br	18

Conditions: 20 h continuous irradiation in vacuum flame-sealed NMR tubes. Conversions obtained from ¹H NMR integrals. ^a Obtained from Nguyen *et al.*⁴⁶ See the Supporting Information for details.

In conclusion, we have herein communicated a new homoleptic Re^I tris(diisocyanide) complex with a high energy ³MLCT excited state that can be exploited for photoredox catalysis. The low photoluminescence quantum yield and the comparatively short excited state lifetime presumably arise from thermal population of ligand field excited states.⁴⁰ Previously investigated Re^I complexes with monodentate arylisocyanide ligands were not emissive in solution at room temperature and seemed to undergo more rapid photodecomposition,^{40, 42} hence chelating isocyanide ligands lead to much improved photophysical properties and greater

chemical robustness, in analogy to what we recently found for Cr⁰ and Mo⁰ complexes.²⁶⁻²⁸ The Re^I tris(diisocyanide) complex catalyses representative photoredox transformations with similar efficiencies to the popular [Ir(ppy)₃] catalyst.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, variable solvent UV-Vis and emission spectra, UV-Vis photostability experiment, cyclic voltammetry and photoredox catalysis NMR experiments. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

Email: oliver.wenger@unibas.ch

Funding Sources

Financial support by the Swiss National Science Foundation through grant number 200021_156063/1 is gratefully acknowledged.

ACKNOWLEDGMENT

C. B. L. acknowledges a Swiss Government Excellence Postdoctoral Scholarship for Foreign Scholars. The authors thank Dr. L. A. Büldt and Dr. X. Guo for insightful discussion.

REFERENCES

- (1) Ravelli, D.; Protti, S.; Fagnoni, M., Carbon–Carbon Bond Forming Reactions via Photogenerated Intermediates. *Chem. Rev.* **2016**, *116*, 9850–9913.
- (2) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C., Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81*, 6898–6926.
- (3) Arias-Rotondo, D. M.; McCusker, J. K., The Photophysics of Photoredox Catalysis: A Roadmap for Catalyst Design. *Chem. Soc. Rev.* **2016**, *45*, 5803–5820.
- (4) Yoon, T. P.; Ischay, M. A.; Du, J., Visible Light Photocatalysis as a Greener Approach to Photochemical Synthesis. *Nat. Chem.* **2010**, *2*, 527.
- (5) Romero, N. A.; Nicewicz, D. A., Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116*, 10075–10166.
- (6) Larsen, C. B.; Wenger, O. S., Photoredox Catalysis with Metal Complexes Made from Earth-Abundant Elements. *Chem. Eur. J.* **2018**, *24*, 2039–2058.
- (7) Hernandez-Perez, A. C.; Collins, S. K., Heteroleptic Cu-Based Sensitizers in Photoredox Catalysis. *Acc. Chem. Res.* **2016**, *49*, 1557–1565.
- (8) Reiser, O., Shining Light on Copper: Unique Opportunities for Visible-Light-Catalyzed Atom Transfer Radical Addition Reactions and Related Processes. *Acc. Chem. Res.* **2016**, *49*, 1990–1996.
- (9) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C., Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322–5363.
- (10) Yam, V. W.-W.; Lau, V. C.-Y.; Cheung, K.-K., Luminescent Rhenium(I) Carbon Wires: Synthesis, Photophysics, and Electrochemistry. X-ray Crystal Structure of [Re(tBu₃bpy)(CO)₃(C≡CC≡C)Re(tBu₃bpy)(CO)₃]. *Organometallics* **1996**, *15*, 1740–1744.
- (11) Wong, K. M.-C.; Lam, S. C.-F.; Ko, C.-C.; Zhu, N.; Roué, S.; Lapinte, C.; Fathallah, S.; Costuas, K.; Kahlal, S.; Halet, J.-F., Electroswitchable Photoluminescence Activity: Synthesis, Spectroscopy, Electrochemistry, Photophysics, and X-ray Crystal and Electronic Structures of

[Re(bpy)(CO)₃(C≡CC₆H₄C≡C)Fe(C₅Me₅)(dppe)][PF₆]_n (n = 0, 1). *Inorg. Chem.* **2003**, *42*, 7086-7097.

(12) Schoonover, J. R.; Strouse, G. F.; Dyer, R. B.; Bates, W. D.; Chen, P.; Meyer, T. J., Application of Time-Resolved, Step-Scan Fourier Transform Infrared Spectroscopy to Excited-State Electronic Structure in Polypyridyl Complexes of Rhenium(I). *Inorg. Chem.* **1996**, *35*, 273-274.

(13) Lundin, N. J.; Blackman, A. G.; Gordon, K. C.; Officer, D. L., Synthesis and Characterization of a Multicomponent Rhenium(I) Complex for Application as an OLED Dopant. *Angew. Chem. Int. Ed.* **2006**, *45*, 2582-2584.

(14) van der Salm, H.; Larsen, C. B.; McLay, J. R. W.; Fraser, M. G.; Lucas, N. T.; Gordon, K. C., Stretching the Phenazine MO in Dppz: The Effect of Phenyl and Phenyl-Ethynyl Groups on the Photophysics of Re(I) Dppz Complexes. *Dalton Trans.* **2014**, *43*, 17775-17785.

(15) Coleman, A.; Brennan, C.; Vos, J. G.; Pryce, M. T., Photophysical Properties and Applications of Re(I) and Re(I)-Ru(II) Carbonyl Polypyridyl Complexes. *Coord. Chem. Rev.* **2008**, *252*, 2585-2595.

(16) Sacksteder, L.; Zipp, A. P.; Brown, E. A.; Streich, J.; Demas, J. N.; DeGraff, B. A., Luminescence Studies of Pyridine α -Diimine Rhenium(I) Tricarbonyl Complexes. *Inorg. Chem.* **1990**, *29*, 4335-4340.

(17) Kou, Y.; Nabetani, Y.; Masui, D.; Shimada, T.; Takagi, S.; Tachibana, H.; Inoue, H., Direct Detection of Key Reaction Intermediates in Photochemical CO₂ Reduction Sensitized by a Rhenium Bipyridine Complex. *J. Am. Chem. Soc.* **2014**, *136*, 6021-6030.

(18) Morimoto, T.; Nakajima, T.; Sawa, S.; Nakanishi, R.; Imori, D.; Ishitani, O., CO₂ Capture by a Rhenium(I) Complex with the Aid of Triethanolamine. *J. Am. Chem. Soc.* **2013**, *135*, 16825-16828.

(19) Rohacova, J.; Ishitani, O., Rhenium(I) Trinuclear Rings as Highly Efficient Redox Photosensitizers for Photocatalytic CO₂ Reduction. *Chem. Sci.* **2016**, *7*, 6728-6739.

(20) Hawecker, J.; Lehn, J.-M.; Ziessel, R., Photochemical and Electrochemical Reduction of Carbon Dioxide to Carbon Monoxide Mediated by (2,2'-Bipyridine)tricarbonylchlororhenium(I) and Related Complexes as Homogeneous Catalysts. *Helv. Chim. Acta* **1986**, *69*, 1990-2012.

(21) Hayashi, Y.; Kita, S.; Brunschwig, B. S.; Fujita, E., Involvement of a Binuclear Species with the Re-C(O)O-Re Moiety in CO₂ Reduction Catalyzed by Tricarbonyl Rhenium(I) Complexes with Diimine Ligands: Strikingly Slow Formation of the Re-Re and Re-C(O)O-Re Species from Re(dmb)(CO)₃S (dmb = 4,4'-Dimethyl-2,2'-bipyridine, S = Solvent). *J. Am. Chem. Soc.* **2003**, *125*, 11976-11987.

(22) Kurz, P.; Probst, B.; Spingler, B.; Alberto, R., Ligand Variations in [ReX(diimine)(CO)₃] Complexes: Effects on Photocatalytic CO₂ Reduction. *Eur. J. Inorg. Chem.* **2006**, *2006*, 2966-2974.

(23) Windle, C. D.; George, M. W.; Perutz, R. N.; Summers, P. A.; Sun, X. Z.; Whitwood, A. C., Comparison of Rhenium-Porphyrin Dyads for CO₂ Photoreduction: Photocatalytic Studies and Charge Separation Dynamics Studied by Time-Resolved IR Spectroscopy. *Chem. Sci.* **2015**, *6*, 6847-6864.

(24) Matlachowski, C.; Braun, B.; Tschierlei, S.; Schwalbe, M., Photochemical CO₂ Reduction Catalyzed by Phenanthroline Extended Tetramesityl Porphyrin Complexes Linked with a Rhenium(I) Tricarbonyl Unit. *Inorg. Chem.* **2015**, *54*, 10351-10360.

(25) Rehm, D.; Weller, A., Kinetics of Fluorescence Quenching by Electron and H-Atom Transfer. *Isr. J. Chem.* **1970**, *8*, 259-271.

(26) Büldt, L. A.; Guo, X.; Prescimone, A.; Wenger, O. S., A Molybdenum(o) Isocyanide Analogue of Ru(2,2'-Bipyridine)₃²⁺: A Strong Reductant for Photoredox Catalysis. *Angew. Chem. Int. Ed.* **2016**, *55*, 11247-11250.

(27) Büldt, L. A.; Wenger, O. S., Chromium(o), Molybdenum(o), and Tungsten(o) Isocyanide Complexes as Luminophores and Photosensitizers with Long-Lived Excited States. *Angew. Chem. Int. Ed.* **2017**, *56*, 5676-5682.

(28) Büldt, L. A.; Guo, X.; Vogel, R.; Prescimone, A.; Wenger, O. S., A Tris(diisocyanide)chromium(o) Complex Is a Luminescent Analog of Fe(2,2'-Bipyridine)₃²⁺. *J. Am. Chem. Soc.* **2017**, *139*, 985-992.

(29) Barnett, B. R.; Figueroa, J. S., Zero-Valent Isocyanides of Nickel, Palladium and Platinum as Transition Metal σ -Type Lewis Bases. *Chem. Commun.* **2016**, *52*, 13829-13839.

(30) Margulieux, G. W.; Weidemann, N.; Lacy, D. C.; Moore, C. E.; Rheingold, A. L.; Figueroa, J. S., Isocyanide Analogues of [Co(CO)₄]_n: A Tetra(isocyanide) of Cobalt Isolated in Three States of Charge. *J. Am. Chem. Soc.* **2010**, *132*, 5033-5035.

(31) Cameron, C. J.; Walton, R. A.; Edwards, D. A., The Vibrational Spectra of the Homoleptic Isocyanide Complexes [Re(CNR)₆]PF₆, (R = Me, Ph, and 4-MeC₆H₄). *J. Organomet. Chem.* **1984**, *262*, 335-346.

(32) Hahn, F. E., The Coordination Chemistry of Multidentate Isocyanide Ligands. *Angew. Chem. Int. Ed.* **1993**, *32*, 650-665.

(33) Ko, C.-C.; Cheung, A. W.-Y.; Lo, L. T.-L.; Siu, J. W.-K.; Ng, C.-O.; Yiu, S.-M., Syntheses and Photophysical Studies of New Classes of Luminescent Isocyanide Rhenium(I) Diimine Complexes. *Coord. Chem. Rev.* **2012**, *256*, 1546-1555.

(34) Ko, C.-C.; Siu, J. W.-K.; Cheung, A. W.-Y.; Yiu, S.-M., Synthesis, Characterization, and Photophysical and Emission Solvatochromic Study of Rhenium(I) Tetra(isocyanide) Diimine Complexes. *Organometallics* **2011**, *30*, 2701-2711.

(35) Ko, C.-C.; Lo, L. T.-L.; Ng, C.-O.; Yiu, S.-M., Photochemical Synthesis of Intensely Luminescent Isocyanide Rhenium(I) Complexes with Readily Tunable Structural Features. *Chem. Eur. J.* **2010**, *16*, 13773-13782.

(36) Wang, M.; Weyhermüller, T.; Bill, E.; Ye, S.; Wieghardt, K., Structural and Spectroscopic Characterization of Rhenium Complexes Containing Neutral, Monoanionic, and Dianionic Ligands of 2,2'-Bipyridines and 2,2':6,2''-Terpyridines: An Experimental and Density Functional Theory (DFT)-Computational Study. *Inorg. Chem.* **2016**, *55*, 5019-5036.

(37) Allen, E. A.; Johnson, N. P.; Rosevear, D. T.; Wilkinson, W., Complexes of Rhenium-(IV) and -(V) with Ligands Containing Group VIB Atoms. *J. Chem. Soc. A* **1969**, 788-791.

(38) Kuimova, M. K.; Alsindi, W. Z.; Blake, A. J.; Davies, E. S.; Lampus, D. J.; Matousek, P.; McMaster, J.; Parker, A. W.; Towrie, M.; Sun, X.-Z.; Wilson, C.; George, M. W., Probing the Solvent Dependent Photophysics of fac-[Re(CO)₃(dppz-X₂)Cl] (dppz-X₂ = 11,12-X₂-dipyrido[3,2-a:2',3'-c]phenazine); X = CH₃, H, F, Cl, CF₃). *Inorg. Chem.* **2008**, *47*, 9857-9869.

(39) Connick, W. B.; Di Bilio, A. J.; Hill, M. G.; Winkler, J. R.; Gray, H. B., Tricarbonyl(1,10-phenanthroline) (imidazole) Rhenium(I): A Powerful Photooxidant for Investigations of Electron Tunneling in Proteins. *Inorg. Chim. Acta* **1995**, *240*, 169-173.

(40) Villegas, J. M.; Stoyanov, S. R.; Reibenspies, J. H.; Rillema, D. P., Spectroscopic and Computational Investigations of the Temperature-Dependent Emission Behavior of [Re(CN)₅Cl] and [Re(CN)₅]⁺ Complexes. *Organometallics* **2005**, *24*, 395-404.

(41) Kim, T. Y.; Elliott, A. B. S.; Shaffer, K. J.; John McAdam, C.; Gordon, K. C.; Crowley, J. D., Rhenium(I) Complexes of Readily Functionalized Bidentate Pyridyl-1,2,3-triazole "Click" Ligands: A Systematic Synthetic, Spectroscopic and Computational Study. *Polyhedron* **2013**, *52*, 1391-1398.

(42) Stacy, N. E.; Conner, K. A.; McMillin, D. R.; Walton, R. A., Studies of the Photoinduced Ligand Substitution Reactions of Hexakis(2,6-dimethylphenyl isocyanide)rhenium(I) in the Presence of Halide Ions. *Inorg. Chem.* **1986**, *25*, 3649-3652.

(43) Hari, D. P.; König, B., Synthetic Applications of Eosin Y in Photoredox Catalysis. *Chem. Commun.* **2014**, *50*, 6688-6699.

(44) Zhang, Y.; Petersen, J. L.; Milsman, C., A Luminescent Zirconium(IV) Complex as a Molecular Photosensitizer for Visible Light Photoredox Catalysis. *J. Am. Chem. Soc.* **2016**, *138*, 13115-13118.

(45) Kokes, F. C.; Hine, J., Kinetics of the Hydrolysis of Fluoromethyl Methyl Ether in Neutral to Alkaline Solution. *J. Org. Chem.* **1976**, *41*, 1976-1979.

(46) Nguyen, J. D.; D'Amato, E. M.; Narayanam, J. M. R.; Stephenson, C. R. J., Engaging Unactivated Alkyl, Alkenyl and Aryl Iodides in Visible-Light-Mediated Free Radical Reactions. *Nat. Chem.* **2012**, *4*, 854-859.

A homoleptic Re^{I} complex bearing chelating diisocyanide ligands and exhibiting a weakly photoluminescent high-energy $^3\text{MLCT}$ excited state is studied and exploited for photoredox catalysis.

