

RESEARCH ARTICLE

The prevalence of brucellosis and bovine tuberculosis in ruminants in Sidi Kacem Province, Morocco

Hind Yahyaoui Azami^{1,2,3*}, Marie J. Ducrot⁴, Mohammed Bouslikhane¹, Jan Hattendorf^{2,3}, Mike Thrusfield⁵, Raquel Conde-Álvarez⁶, Ignacio Moriyón⁶, Amaia Zúñiga-Ripa⁶, Pilar M. Muñoz Álvaro⁷, Virginie Mick⁸, Ward Bryssinckx⁹, Sue C. Welburn⁴, Jakob Zinsstag^{2,3}

1 Department of Veterinary Pathology and Public Health, Agronomic and Veterinary Institute Hassan II, Rabat, Maroc, **2** Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Basel, Switzerland, **3** University of Basel, Basel, Switzerland, **4** Division of Infection and Pathway Medicine, Centre for Infectious Diseases, School of Biomedical Sciences, College of Medicine and Veterinary Medicine, The University of Edinburgh, Edinburgh, United Kingdom, **5** Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Easter Bush Veterinary Centre, Roslin, Midlothian, United Kingdom, **6** IDISNA - Instituto de Salud Tropical y Depto. Microbiología y Parasitología, Universidad de Navarra, Edificio de Investigación, Pamplona, Spain, **7** Unidad de Producción y Sanidad Animal, Instituto Agroalimentario de Aragón- IA2 - (CITA-Universidad de Zaragoza), Zaragoza, Spain, **8** Paris-Est University/Anses, EU/OIE/FAO & National Reference Laboratory for brucellosis, Animal Health Laboratory, Maisons-Alfort, France, **9** Avia-GIS, Zoersel, Belgium

* yahyaouiiazamihind@gmail.com



OPEN ACCESS

Citation: Yahyaoui Azami H, Ducrot MJ, Bouslikhane M, Hattendorf J, Thrusfield M, Conde-Álvarez R, et al. (2018) The prevalence of brucellosis and bovine tuberculosis in ruminants in Sidi Kacem Province, Morocco. PLoS ONE 13(9): e0203360. <https://doi.org/10.1371/journal.pone.0203360>

Editor: Roy Martin Roop, II, East Carolina University Brody School of Medicine, UNITED STATES

Received: December 11, 2017

Accepted: August 20, 2018

Published: September 18, 2018

Copyright: © 2018 Yahyaoui Azami et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The present study is a component of a large European Union entitled the Integrated Control of Neglected Zoonoses (ICONZ), grant agreement n° 221948. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. One of

Abstract

Bovine tuberculosis (BTB) and brucellosis are major endemic zoonoses in ruminants in Morocco that impact on both animal and human health. This study presents an assessment of the epidemiological and socioeconomic burden of bacterial zoonoses in Sidi Kacem Province in Northern Morocco from a cross-sectional survey of 125 cattle and/or small ruminant-owning households. In total, 1082 sheep and goats were examined from 81 households. The single intradermal comparative cervical test to screen for bovine tuberculosis was undertaken on 1194 cattle from 123 households and all cattle were blood sampled. Cattle and small ruminant sera were tested for brucellosis using the standard Rose Bengal Test (sRBT) and the modified Rose Bengal Test (mRBT). Bacteriology was performed on 21 milk samples obtained from cattle that were seropositive for brucellosis for isolation and phenotyping of circulating *Brucella* strains. Individual and herd prevalence for BTB in cattle of 20.4% (95% CI 18%-23%) and 57.7% (95% CI 48%-66%), respectively, were observed in this study. The prevalence of brucellosis in cattle at individual and herd level was 1.9% (95% CI 1.2%-2.8%) and 9% (95% CI 4.5%-1.5%), respectively. *Brucella* pathogens were isolated from three cattle milk samples and were identified as *B. abortus* using Brucladder[®] multiplex PCR and *B. abortus* biovar 1 by classical phenotyping. All small ruminants were seronegative to sRBT, two were positive to mRBT. A higher risk of BTB and brucellosis was observed in cattle in intensive livestock systems, in imported and crossed breeds and in animals from larger herds (>15). The three risk factors were usually present in the same herds, leading to higher transmission risk and persistence of both zoonoses. These results highlight the importance of implementing control strategies for both BTB and brucellosis to

the authors was employed by a commercial company: Avia-GIS. The funder provided support in the form of salaries for author WB, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

Competing interests: The former commercial affiliation of WB with Avia-GIS does not alter our adherence to PLOS ONE policies on sharing data and materials.

reduce productivity losses and the risk of transmission to humans. Prioritising control for BTB and brucellosis in intensive livestock production systems is essential for human and animal health.

Introduction

Bovine tuberculosis (BTB) and brucellosis are bacterial zoonoses endemic in cattle, and brucellosis is endemic in small ruminants, in Morocco. These diseases are prioritized in Moroccan veterinary legislation [1,2] but remain poorly controlled. Infectious diseases impose a heavy financial burden on the livestock sector [3] and zoonoses have a dual impact through human disease burden and productivity losses of livestock, on which rural families depend for their livelihoods [4]. While brucellosis and BTB have been controlled and/or eliminated in many developed countries [5,6], in developing nations, these diseases are neglected [7] with the World Health Organization considering control of zoonotic tuberculosis to be a major priority [8]. Rapid growth and intensification of livestock systems is expected to result in an increase in prevalence of both brucellosis and BTB [9].

The infectious agent of bovine BTB is *Mycobacterium bovis*, a member of the *Mycobacterium tuberculosis* complex. Despite a host preference for cattle [10], *M. bovis* can infect a wide range of domestic and wild animals [11,12]. Cattle to cattle transmission occurs via direct contact (aerosols) and depends on a number of factors including the number of bacilli excreted and herd density [13]. Transmission of BTB to humans occurs mainly through consumption of infected raw milk, although direct transmission can occur [14]. Bovine tuberculosis is still common in Morocco. A national tuberculosis survey in cattle in 2004, using the single intradermal tuberculin test, showed individual and herd prevalence of 18% (n = 13021) and 33% (n = 2263), respectively [15]. Bovine tuberculosis is responsible for meat losses due to carcass condemnation, and causes a decrease in herd productivity and milk yields [16]. In Morocco, Government BTB control initiatives include tuberculin testing and the slaughtering of positive animals (reactors). At national level, the mandatory BTB control strategy is for test and slaughter but legislation is poorly enforced and no programme of systematic BTB screening of cattle is in place [2,17].

Brucellosis is caused by gram-negative bacteria of the genus *Brucella*. *Brucella melitensis* and *B. abortus* cause disease mostly in small ruminants and cattle, respectively [18]. Although it displays a preferential host-range, *B. melitensis* can infect cattle in mixed breeding areas where it coexists with small ruminants [19]. Also, in West Africa *B. abortus* infection in small ruminants is noted to occur in areas where the animals are in contact with cattle and where *B. melitensis* is absent [20] [19]. Brucellosis is spread through contact with abortion products and vaginal fluids and by milk feeding, through semen or congenitally. Sheep can also be infected by *B. ovis*, a non-zoonotic species. *Brucella melitensis* and *B. abortus*, together with *B. suis* and *B. canis*, cause human brucellosis and contact with infected animals and consumption of raw dairy products is the most common source of transmission [21]. Brucellosis causes economic losses in livestock due to abortions and infertility.

National epidemiological surveys for brucellosis in Morocco in 1996 and 2010 showed that bovine brucellosis is more prevalent in the north-west coastal and central zones where the cattle density is the highest, with a mean individual and herd prevalence of 2.1% (n = 8991) and 4.9% (n = 1168), respectively (Government survey, 2010). Herd seroprevalences have remained at a similar level to those reported in 1977 (4.6%) and in 1988 (4.9%) [19]. Initiatives

to control brucellosis in Morocco have had varied success. A national vaccination campaign using S19 from 1989 to 1994 showed little impact on herd prevalence [22]. By contrast, a public-private initiative (2007) that included RB51 and/or S19 vaccination and test and slaughter on farms that are members of professional associations or cooperatives reduced brucellosis herd seroprevalence from 40% to 0.4% in member farms [23]. In total 81230 cattle were serologically tested for brucellosis, 55869 were vaccinated and 2901 were culled at a cost of \$US 2.6 million. A large bacteriological study comprising 500 samples from 357 cattle isolated *B. abortus* biovar 1 and 3 in the 1980s [24,25].

Brucellosis in small ruminants is a recognized problem across the northern Mediterranean zone of Morocco and in inland mountainous areas where sheep and goat populations dominate. In the mid-1990s, small ruminant brucellosis emerged in in the Oriental region where Morocco shares a border with Algeria and prompted a programme of mass vaccination of small ruminants with the live attenuated vaccine *B. melitensis* Rev1 via the conjunctival route in the zone between 1997 and 2003. Premature ending of the vaccination campaign resulted in disease re-emergence and between 2009 and 2013 48 individual cases of small ruminant brucellosis were reported in the Northeastern region. A National survey in 2006 testing 11609 small ruminants yielded eight sero-positives, five of which were from the Oriental region (19). Bacteriological evidence of brucellosis in small ruminants in Morocco is limited to only three studies and they reported isolation of *B. melitensis* biovar 3 [26–28].

This study aimed to estimate the current prevalence of bovine tuberculosis in cattle, and the seroprevalence of the zoonotic *Brucella* species in cattle, sheep and goats, and to assess BTB and brucellosis risk factors in Sidi Kacem province in Morocco.

Material and methods

Study area

Sidi Kacem province comprises 29 communes and is sub-divided into two agro-hydrologic zones: “rainfed” (bour) in the northeast and “irrigated” in the southwest. The irrigated zone is characterized by low-lying plains, contrasting with the rainfed zone, which is mountainous (altitude from 150 to 500m). The number of cattle in the rainfed zone is 40% of the total cattle population of the province. The irrigated zone is dominated by an intensive mode of livestock rearing characterized by larger herd sizes (15 cattle/herd) [29] and European dairy breeds (Holstein-Friesian, Montbeliarde, Tarentaise) or cross-bred cattle. The livestock production system in the rainfed zone is more extensive, with dominance of local cattle breeds and smaller herd sizes (5 cattle/herd).

Sampling

Sample size was determined using standard formulae for cluster surveys [30], we considered the village as a cluster, assuming the following parameters, i) a BTB prevalence of 18% [national survey 2004 [15]], ii) a mean number of cattle per household of 9, iii) an average number of 2 households selected per cluster and iv) an intra-class correlation coefficient of 0.2 [31,32]. An average of 62 villages (douars) and 124 households are required to estimate the prevalence with a precision (defined as one half-length of the 95% CI) of 5%-points. Clusters were randomly selected based on the official village lists available for Sidi Kacem province. Two households per village were randomly selected based on livestock-owning households provided by the chief of the commune upon arrival in each village; all cattle in cattle owning households were sampled. As small ruminant flock numbers were high, not all sheep and goats were sampled in small-ruminant owning households, a maximum of 20 animals were sampled per household.

Herd and animal level data

At herd level, GPS coordinates (Fig 1.), livestock production system, grazing system and herd size were recorded.

Most sampled households owned cattle, but not small ruminants. Table 1 shows the number of households and the species present within the household. For every animal, age, gender, breed and body condition score (BCS) were recorded.

Diagnosis of bovine tuberculosis

The single intradermal comparative cervical skin test (SICCT) was used following OIE Terrestrial Manual standards for BTB screening [33]. Briefly, injection sites were clipped and cleaned; a fold of skin within each clipped area was measured with callipers and 0.2 ml each of bovine and avian purified protein derivative (PPD) (25000 IU/ml) was injected in the left neck region. Injections were performed using a separate intradermal gun for each PPD in two spots separated by approximately 15 cm. The skin-fold thickness at each injection site was re-measured 72h after injection by the same individual. Any exudate, oedema and pain observed in the injection sites were recorded. The SICCT was interpreted using the OIE recommended cut off and an animal was considered positive if the increase in skin thickness of the bovine PPD injection site was greater than the skin thickness of the Avian PPD injection site by 4 mm or more. The reaction was considered inconclusive if the difference was between 1 and 4 mm. The reaction was interpreted as negative if the increase in the skin thickness was equal for the bovine and avian PPD injection sites [33].

Brucellosis sero-diagnosis and bacteriology

Blood from 1194 cattle and 1082 small ruminants across 125 households was collected (Fig 1) and all sera were stored in cool boxes before laboratory processing. After 24h storage at 4°C to allow serum separation, the tubes were centrifuged, and sera extracted, aliquoted and tested for antibodies to smooth *Brucella* using the Rose Bengal Test (RBT). There are two OIE accepted [34] variants of the test: the standard (sRBT) and the modified (mRBT) RBT that differ in the proportion of serum:antigen used: equal volumes (25–30 µl) in the sRBT and a higher volume of serum (75 µl serum and 25 µl antigen) in the mRBT. The mRBT increases the sensitivity when testing sera from small ruminants [35,36] but this modification is not recommended when testing cattle sera because it reduces the specificity in areas where S19 vaccination is implemented. Accordingly, the following testing strategy was used. Immediately after collection, one aliquot was used for screening of both cattle and small ruminant sera using the mRBT. This initial screening aimed to collect milk samples from seropositive lactating females for bacteriological investigations; using the mRBT increased the diagnostic sensitivity thereby maximising the number of milk samples collected. A second aliquot was stored at -18°C and sent to the “Instituto de Salud Tropical y Depto. Microbiología y Parasitología, Universidad de Navarra (UNAV)” for screening. Cattle samples were screened using the sRBT protocol and small ruminant samples were screened using sRBT and mRBT in parallel. The antigen used was a suspension of fully smooth *B. abortus* 1119 standardized according to international guidelines [34] and controlled for quality using a panel of brucellosis positive and negative serum samples [37].

Bacteriology was undertaken on milk samples obtained from 21 seropositive cattle (three CITA and three Farrell's selective media plates for each sample) [38]. After incubation for 4–7 days at 37°C (5–10% CO₂ atmosphere), suspicious colonies were re-plated on the same media for preliminary identification. Isolates were considered presumptive for *Brucella* by agglutination with anti-*Brucella* serum coated staphylococci and oxidase and urease tests [39]. Colonies




Fig 1. Sidi Kacem rainfed and irrigated regions and geo-localisation of the household screened.

<https://doi.org/10.1371/journal.pone.0203360.g001>

found to be suspicious were stored at -20°C in vials of Tryptic Soy Broth (TSB) with 5% DMSO for further typing at the WHO and OIE collaborating centres for the diagnosis of animal and human brucellosis at the “Universidad de Navarra (UNAV)” and “Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA)” in Spain. Isolates were confirmed as *Brucella*

Table 1. Number of sampled households in terms of present species.

Species owned by household	Number of households
Cattle only	44
Sheep only	2
Goats only	0
Cattle sheep and goats	4
Cattle and sheep	75
Cattle and goats	0
Sheep and goats	0
TOTAL	125

<https://doi.org/10.1371/journal.pone.0203360.t001>

species and typed at species level using Bruceladder[®] multiplex PCR [40] and, at biovar level, using classical phenotyping methods [39], i.e. oxidase and urease tests, CO₂ requirement, agglutination with monospecific anti-A/anti-M sera, lysis with phages (Tb, Wb, Iz1 and R/C), sensitivity to dyes (fucine, thionin and saphranine) and Crystal Violet exclusion test (to assess absence of dissociation) [39].

Statistical analyses

Data were entered in Access 2012 and analysed using Stata 12.1. Generalized linear mixed models (GLMM) for binary outcomes were used to estimate prevalence and to identify risk factors. The GLMMs included a logit link function to estimate the odds ratios (OR) to quantify the relationship between different risk factors and the diseases [30]. Initial inspection of the variance components using multilevel models showed that most of the variance in the response is at household level rather than at village level and therefore we included households as random effect to account for clustering. Prevalence and corresponding confidence intervals were estimated fitting a constant only model. The predictor variables evaluated were livestock production system (categorized as intensive, semi-intensive, extensive), rainfed or irrigated grazing, age (up to 1 year, up to 3 years and older), sex, breed and body condition score (BSC, 1 to 2, 2.5 to 3, 3.5+). Categories with very few observations, i.e. extensive production system and BSC above 3.5, were collapsed with their adjacent categories for the analysis. The multivariable model was pre-specified and included all predictor variables. Because the individual level brucellosis prevalence was low we had to exclude predictor variables without positive animals in one of their categories to avoid (quasi) complete separation. All inconclusive diagnostic test results were omitted prior analysis.

Considering that SICTT has a high specificity and an average sensitivity, we calculated both apparent and true individual prevalence. To calculate true prevalence, we used the Rogan Gladen-estimator [41] assuming a diagnostic sensitivity of 51.1% and a specificity of 98.9% as previously reported by Müller et al. from Chadian cattle using SICTT [42]. The true individual of BTB prevalence was estimated as: $\{\text{True prevalence} = (\text{Apparent prevalence} + \text{sp} - 1) / (\text{sp} + \text{se} - 1)\}$. For brucellosis, given the high sensitivity and specificity of sRBT we can assume that apparent prevalence is close to true prevalence [42]. Sensitivity analysis used villages as clusters.

Results

Cattle characteristics

In total, 1201 bovines from 125 farms in 62 villages were examined of which 78.4% (n = 938) were female and 85% were crossbreeds (n = 1016), and almost all (96%, n = 1150) had a BCS equal to or less than 3. Most cattle (78%) (n = 936) were over 12 months. Cattle were almost equally distributed between the rain fed and the irrigated area, most from semi intensive herds (82.8%, n = 994) and only a few (n = 20) from extensive farming systems. Most herd sizes were below 15 (Table 2).

Prevalence for bovine tuberculosis and brucellosis

Tuberculin skin test results were obtained for 1194 cattle. Of these, 107 animals were found inconclusive for BTB using the OIE interpretation criteria of the SICCT and were not included in the analysis. An individual apparent BTB prevalence of 20.4% (95% CI 18%-23%) and a herd prevalence of 57.7% (95% CI 48%-67%) were found for the remaining animals. Assuming a sensitivity of 51% and specificity of 99% as reported by Müller et al [43], a true individual BTB prevalence of 38.2% was estimated.

Table 2. Basic characteristics of 1201 cattle sampled in Sidi Kacem, Morocco.

<i>Characteristics</i>	<i>Classes</i>	<i>N (%)</i>
Herd level		
Livestock production system	Intensive	15 (12.2)
	Semi-intensive	105 (85.4)
	Extensive	3 (2.4)
Grazing system	Rainfed	58 (46.4)
	Irrigated	67 (53.6)
Herd size	1–15	92 (74.8)
	>15	31 (25.2)
Individual level		
Livestock production system	Intensive	187 (15.6)
	Semi-intensive	994 (82.8)
	Extensive	20 (1.6)
Grazing system	Rainfed	596 (49.6)
	Irrigated	605 (50.4)
Age (months)	0–12	265 (22.1)
	13–36	397 (33.1)
	>36	539 (44.8)
Sex	Female	938 (78.4)
	Male	258 (21.6)
Breed	Crossed	1016 (84.9)
	Imported	100 (8.4)
	Local	81 (6.7)
Body condition score	1–2	366 (30.5)
	2.5–3	784 (65.3)
	3.5–4	15 (1.3)

<https://doi.org/10.1371/journal.pone.0203360.t002>

Brucellosis results using sRBT were available for 1179 cattle. Prevalence of bovine brucellosis was 1.9% (95% CI 1%-3%) at individual level and 9% (95% CI 5%-15%) at herd level. None of the 1044 sheep or 51 goats screened were sRBT positive and only 2/1044 sheep were mRBT positive.

Risk factors analysis

a. Bovine tuberculosis. The uni- and multivariable analysis of BTB risk factors are shown in Table 3. Age higher than 36 months was significantly associated with a higher risk of BTB compared with age below 12 months (28.2% vs 13.7% OR: 2.6, 95% CI 1.4–4.8). Imported breeds were observed to have a lower risk of BTB compared to crossbreeds (16.7% vs 21.7%, OR: 0.4, 95% CI 0.1–0.8). We observed the lowest prevalence in local breeds (9.6%) but the confidence interval was broad and included unity (OR: 0.5 95% CI 0.2–1.4). Male animals showed a lower prevalence (12.0%) compared to female animals (22.7%) but this relationship was not statistically significant in the multivariable model (OR: 0.8 95% CI 0.4–1.4), likely due to the fact that only very few males were older than 36 months.

At herd level, irrigated grazing systems showed a significantly higher risk of BTB compared with rainfed systems (30.1% vs 11%, OR: 3.1 95% CI 1.6–6.2). Large herd size showed a higher risk of BTB than small to medium herd size (26.2% vs 15.7%, OR: 2.1, 95% CI 1.0–4.3).

Table 3. Individual and herd risk factors of BTB in 1087 cattle.

Risk factor	Classes	N screened	% (N positive)	OR (95% CI)	mOR (95% CI)
Herd level					
Livestock production system	Intensive	170	34.7 (59)	Ref	—
	Semi-intensive & extensive	922	17.8 (164)	0.3 (0.1–0.9)	0.5 (0.2–1.4)
Grazing system	Rainfed	550	11.0 (60)	Ref	—
	Irrigated	544	30.1 (164)	3.5 (1.8–6.5)	3.1 (1.6–6.2)
Herd size (animals)	1–15	600	15.7 (94)	Ref	—
	>15	492	26.2 (129)	2.0 (0.9–4.1)	2.1 (1.0–4.3)
Individual level					
Breed	Crossed	918	21.7 (199)	Ref	—
	Imported	96	16.7 (16)	0.3 (0.1–0.7)	0.4 (0.1–0.8)
	Local	73	9.6 (7)	0.5 (0.2–1.3)	0.5 (0.2–1.4)
Age (months)	0–12	241	13.7 (33)	Ref	—
	13–36	367	14.4 (53)	1.4 (0.8–2.5)	1.2 (0.7–2.3)
	>36	475	28.2 (134)	3.9 (2.3–6.6)	2.6 (1.4–4.8)
Sex	Female	845	22.7 (192)	Ref	—
	Male	241	12.0 (29)	0.3 (0.2–0.6)	0.8 (0.4–1.4)
Body condition score	1–2	325	29.5 (96)	Ref	—
	2.5–4	732	16.7 (122)	0.6 (0.4–0.8)	0.8 (0.5–1.3)

mOR: multivariable analysis OR

Ref: Reference category

<https://doi.org/10.1371/journal.pone.0203360.t003>

b. Brucellosis. Uni- and multivariable analysis of risk factors for brucellosis are shown in [Table 4](#). Due to the low number of positive brucellosis cases, the study was not sufficiently powered to detect statistically significant differences.

We observed a lower prevalence in animals with a BCS lower than 2 (0.7% vs 4.7%), and a higher prevalence in irrigated grazing systems (3.2% vs 0.5%) and in large herds (3.2% vs 0.8%).

Characterization of *Brucella* isolates

Three *Brucella* strains isolated from female cattle were identified as *B. abortus* biovar 1 through classical typing. Isolates were from cows belonging to the same herd located in the irrigated zone. The cows were over 72 months old; two had a history of abortions and BCS of less than 2.

Discussion

The overall BTB individual apparent prevalence in this study was 20.4%, similar to that reported in 2004 (18%) during national tuberculin skin testing. However, given the reported low sensitivity of BTB testing these figures are likely to be underestimated. The BTB 57.7% herd prevalence in this study is higher than that previously reported at 33% in 2004 at national level [15]. Despite that the 2004 survey applied the single intradermal tuberculin skin test (SITT), which is less specific than the SICTT used in the present study. While the single tuberculin skin test (SITT) may result in a high number of false positives, the comparative tuberculin skin test (SICTT) was shown to reduce false positives and cross reactions (i.e. the single test has a higher diagnostic sensitivity and the comparative test is more specific) [44]. The BTB prevalence in the present study is also far higher than those reported in Uganda (6%) [45],

Table 4. Individual and herd risk factors of bovine brucellosis in 1177 cattle.

Risk factor	Classes	N screened	% (N positive)	OR (95% CI)	mOR (95% CI)
Herd level					
Livestock production system	Intensive	187	6.9 (13)	Ref	--
	Semi-intensive & extensive	994	0.9 (9)	0.3 (0.05–2.02)	0.6 (0.1–4.3)
Grazing system	Rainfed	595	0.5 (3)	Ref	--
	Irrigated	586	3.2 (19)	4.7 (0.8–26.7)	3.4 (0.6–21.0)
Herd size (animals)	1–15	643	0.77 (5)	Ref	--
	>15	538	3.16 (17)	2.8 (0.5–15.1)	2.3 (0.5–11.9)
Individual level					
Breed	Crossed	997	2.2 (22)	Ref	--
	Imported	100	0.0 (0)	Nd	--
	Local	80	0.0 (0)	Nd	--
Age (months)	0–12	257	0.4 (1)	Ref	--
	13–36	388	0.3 (1)	0.5 (0.03–8.5)	0.4 (0.02–7.41)
	>36	527	3.8 (20)	6.9 (0.8–60.4)	5.1 (0.6–45.6)
Sex	Female	928	2.4 (22)	Ref	--
	Male	248	0.0 (0)	Nd	--
Body condition score	1–2	363	4.7 (17)	Ref	--
	2.5–4	783	0.6 (5)	0.2 (0.05–0.7)	0.5 (0.1–1.8)

mOR: multivariable analysis OR

Ref: Reference category

Nd: Not determined because of perfect prediction

<https://doi.org/10.1371/journal.pone.0203360.t004>

Niger (3.6%) [46], rural Ethiopia (5.5%) [47], and Tanzania (2.4%) [48]. The high prevalence in Morocco may be explained by more intensive dairy cattle farming practices (see below), similar to Central Ethiopia, where a herd prevalence of 50% was registered in 2012 using SICTT [49]. On the other hand, BTB prevalence in Morocco (20.4%) was lower than that reported in Zambia (49.8%) and Mozambique (39.6%) [50,51].

The gold standard for brucellosis diagnosis is isolation and identification of *Brucella* spp. However, bacteriological culture is cumbersome, expensive and requires skill and facilities rarely available in resource-poor countries and consequently, indirect testing of anti-*Brucella* antibodies in serum is commonly applied for brucellosis screening. In this work, brucellosis apparent seroprevalence was examined using the sRBT for cattle and the mRBT for small ruminants. A rigorous meta-analysis of diagnostic tests for bovine brucellosis [52] based on solid data gathered by a panel of experts [53] has shown that the sRBT performs with diagnostic sensitivity and specificity values of 98.1% (95% CI [96.8%–99.1%]) and 99.8% (95% CI [99.7%–99.8%]), respectively, where vaccination is not practiced. Accordingly, and taking into account that vaccination was not undertaken in the area, no further serological testing by the so-called complementary tests was necessary. This is worth commenting upon because there are a number of works assuming that RBT lacks specificity in the absence of vaccination and needs to be confirmed by a second test. The above-cited meta-analysis and a recent rigorous analysis of publications meeting strict scientific criteria [42] prove that this is a misconception, possibly resulting from a misinterpretation of the use of tests in programs where eradication through vaccination in parallel to test and slaughter is applied (for a discussion see [42,54]). This is also true for small ruminants testing where optimal sensitivity is obtained using the mRBT [35]. Indeed, mRBT has been shown to be useful for reducing the cost and the time of

brucellosis screening in *B. melitensis* eradication campaigns [55]. The standard serum agglutination test is not recommended by OIE due to its low sensitivity, and the complement fixation test is technically cumbersome and does not outperform RBT in terms of specificity and sensitivity in the absence of vaccination [35,42]. Indirect ELISA shows similar sensitivity and specificity to RBT in the absence of vaccination, but is more expensive [52] and need to be validated in the target population since its diagnostic performance depends on the manufacturer and the selection of an adequate cut-off to discriminate positive and negative samples [56] [42]. Despite original reports [57], recent studies show that competitive ELISAs do not have optimal sensitivity [58]. Thus, as Morocco is a country where no official brucellosis vaccination is currently practiced, sRBT and mRBT can be recommended for serological surveys in cattle and small ruminants respectively.

Using these methods, individual apparent brucellosis prevalence found in this work in cattle (1.9%) is near the 2.1% reported for the national survey of 2010–2011. National statistics from 1982 to 1992 registered a mean herd prevalence ranging from 2.1% to 4.9%, which is lower than brucellosis herd prevalence (9%) found in the present study [26]. A recent study using mRBT to investigate bovine brucellosis in 25 farms (221 cattle) reported a prevalence of 33.48%, which is higher than that reported by our study [59]. However, the high prevalence could have been due to the irrigated study area and the targeted animals (females older than 18 months). Individual brucellosis prevalence found in Sidi Kacem was also lower than that found in Egypt (23.8%) [60], Uganda (5%) [61], Nigeria in 2011 (4.04%) [62] and Ethiopia in 2006 (2.9%) [63]. The *Brucella* strains isolated from the three Friesian/Holstein cows and belonging to the same herd in the irrigated zone were found to be *B. abortus* biovar 1, and phylogenomic studies (not described here) show that these strains have some homogeneity with Spanish strains. This is in line with a study conducted in the mid-1980s yielding 8 *B. abortus* biovar 1 and 28 *B. abortus* biovar 3 strains from 500 samples [24,25]. A subsample of these Moroccan strains was examined as part of a study characterising *B. abortus* strains of African origin and found the 12 Moroccan strains to be identical to *B. abortus* biotypes isolated in Europe.

In this study only two sheep from two different herds in the irrigated area showed a positive mRBT result. The mRBT positive sheep were from a herd that had several cases of cattle brucellosis. Livestock from this village grazed on shared pasture and the sheep may have developed seropositivity due to contact with *B. abortus* infected cattle. These sheep could have developed antibodies but cleared the infection and hence be of negative infection status or alternatively be infected and a source of contagion to other ruminants or humans. A study performed in Northern Morocco and the Middle Atlas in 2013, investigated brucellosis among 23 sheep and goat herds using mRBT. The results of the study showed an individual and herd prevalence of 13.3% and 43% respectively. The high prevalence registered could be explained by the condition of animals sampled, which were exclusively aborting females, so abortion could be considered as a confounding factor for that study [64]. Very little is known about the course of infection by *B. abortus* in sheep, which seems to be a very rare event even in areas free of *B. melitensis* and where cattle are infected by *B. abortus* [65]. Regrettably, slaughter for collection of necropsy samples for bacteriological confirmation were beyond the scope of this study. Further investigations are necessary to clarify the epidemiology of small ruminant brucellosis in the areas targeted by the study.

Consistent with a recent review [9], we observed a higher BTB and brucellosis prevalence in intensive livestock production systems, although the relationships were not statistically significant. Intensive dairy farming conditions within a confined environment with less access to sun, air flow and high humidity conditions may enhance transmission of *Brucella* and *Mycobacteria* between animals. The higher stocking density, larger herd sizes, and propensity to buy

in animals, together with higher calving frequency in intensive systems may also increase the risk of disease transmission. On the other hand, extensive systems, where animals are grazed in lower density on communal pasture, may have a lower risk of transmission because of the effect on the sun and heat on *Brucella* and *Mycobacteria* environmental contaminants [19]. The shift to a more intensive livestock production observed in rapidly urbanizing developing countries such as Morocco, could lead to an increase of zoonotic diseases transmission in the absence of animal recording systems, movement controls and well managed veterinary services [19,66].

As has been previously reported, Indigenous animals showed a lower risk of BTB, [51] compared to imported and cross breed animals. This may be explained by their non-adaptation to local conditions. Local indigenous breeds may not be a protective factor, however, because in Morocco extensive herds are dominated by local and cross-bred cattle, while intensive herds are dominated by imported breeds. Consequently, the breed could be a confounding factor for the production system [18]. Older animals showed a higher risk of BTB and bovine brucellosis as described in other studies [67]. All positive brucellosis cases were found to be females, but it must be interpreted in light of the fact that the sample was composed predominantly of female cattle. In Nigeria, females were described to be more susceptible to *B. abortus* infection [68]. In the present study, a low BCS showed a higher risk of BTB. BCS has previously been linked with BTB infection [69,70], although recent studies in Ireland and Tanzania showed no evidence of the association of low BCS and high BTB prevalence [71,72]. Low BCS could be linked to clinically advanced BTB [73], but in this study the initial status of the sampled animals was unknown.

Based on the findings of this study we conclude that bovine tuberculosis and brucellosis are prevalent in Sidi Kacem. Considering the economic losses caused by BTB and brucellosis, in addition to their public health impact, additional efforts should be deployed to design an integrated control strategy. Test and slaughter has been shown to be the most efficient elimination strategy for BTB in several countries. Many factors contribute to the success of a control and elimination campaign. Trust between all stakeholders especially the farming industry, the government and the farmers are a very important component which contributed to the success of brucellosis and BTB control program in other contexts. Correct application of livestock biosecurity measures, early diagnosis of the disease, and application of movement restrictions also affect the success of a control campaign [74]. However, in Morocco, relationships between livestock keepers, local authorities and veterinary services are characterized by mistrust. Solid and sustainable control cannot be achieved without the conviction and participation of all stakeholders. Sensitisation and education campaigns of all stakeholders are required to improve adherence to and acceptance of control programs by local populations and decision makers. A rigorous application of the decided strategy and involvement of the animal owners in the decision process are pivotal for the success of control and elimination programs [75].

The best strategy for controlling brucellosis in Morocco would be conjunctival mass vaccination every two years just before the natural breeding season or immediately after calving/lambing/kidding. This is the best option as prevalence is high and the veterinary services are not able to apply individual tagging allowing vaccination of young replacements only [76]. However, vaccination needs to be sustained over time to be effective. Premature removal of S19 or Rev1, or replacement with a less effective vaccine e.g. RB51, has led to failure of previous attempts for control in Morocco (19).

A BTB transmission model for Morocco indicated that BTB could be controlled within 25 years, if 50% of cattle were tested annually, and infected animals were slaughtered at an estimated cost of 1.55 billion Euros [77]. Taking into consideration the current infectious diseases prioritized by Morocco (e.g. foot and mouth disease, sheep pox virus and Peste des Petits

Ruminants [PPR]), such a control strategy is currently deemed unaffordable for a middle-income country. Wildlife reservoirs can also complicate control operations. The presence of a wildlife reservoir (e.g. badgers in Ireland) has caused reemergence of BTB in developed countries [78]. Wildlife reservoirs have been shown to exist in Africa (feral baboons in Kenya [79] and warthog and buffalo in Uganda [80]). The role of a wildlife reservoir in BTB transmission in Morocco is unknown, and other barriers to control are considered more critical including: lack of efficient organization of veterinary services; prioritization of other highly infectious viral and parasitic diseases; limited technical capacity and financial constraints [81].

As for other neglected zoonosis, evidence and advocacy is necessary to convince policy-makers and communities of the benefits of disease control [82,83]. The evaluation of the cost of BTB should take into account the cost of the human disease, and for this purpose an investigation of the prevalence of *M. bovis* in humans is required, as well as a calculation of the direct and indirect cost of a human TB case. Cross-sectoral socio-economic analysis of the cost of both diseases is needed. In addition, support from private industry (e.g. the milk industry) could sustain BTB and brucellosis control campaigns, as they will also benefit from the control measures. Novel methods for innovative financing should be examined to mobilize investment for interventions that contribute towards elimination of neglected zoonosis [84,85] and benefit human and animal health.

The present study confirms that there is added value in investigating multiple zoonoses simultaneously, especially for zoonoses with a reservoir overlap. Undertaking brucellosis and BTB screening in parallel and in multiple hosts is logistically and technically feasible. The added value of an integrated approach to epidemiological investigations on zoonoses has been demonstrated in Chad [86]. Morocco could consider a parallel elimination campaign for BTB and brucellosis that optimizes use of human and economic resources.

Ethical clearance

The methodology of the study including the household questionnaire was reviewed and validated by international and national research expert partners within the ICONZ project. In addition, the Moroccan veterinary services (ONSSA) approved and granted authorization for the study. The purpose of the study was explained to the local authorities and to the farmers. The tests used for the screening of BTB and brucellosis are routinely used by the veterinary services of Morocco.

Supporting information

S1 Dataset. HouseholdSurvey_ICONZ_2012.

(XLSX)

S2 Dataset. ICONZ_Sidi Kacem_Database_Prevalence.

(XLSX)

S3 Dataset. Serology_Brucellosis_MoroccoICONZ.

(XLSX)

Acknowledgments

The present study is a component of a large European Union entitled the Integrated Control of Neglected Zoonoses (ICONZ), grant agreement n° 221948. The authors are grateful to the veterinary services (ONSSA) at National and Regional level for their contribution to the study.

Author Contributions

Conceptualization: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Mike Thrusfield, Raquel Conde- Álvarez, Ignacio Moriyón, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

Data curation: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Mike Thrusfield, Raquel Conde- Álvarez, Ignacio Moriyón, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

Formal analysis: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Raquel Conde- Álvarez, Ignacio Moriyón, Amaia Zúñiga-Ripa, Pilar M. Muñoz Álvaro, Virginie Mick, Sue C. Welburn, Jakob Zinsstag.

Funding acquisition: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Mike Thrusfield, Raquel Conde- Álvarez, Ignacio Moriyón, Amaia Zúñiga-Ripa, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

Investigation: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Raquel Conde- Álvarez, Ignacio Moriyón, Amaia Zúñiga-Ripa, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

Methodology: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Mike Thrusfield, Raquel Conde- Álvarez, Ignacio Moriyón, Amaia Zúñiga-Ripa, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

Project administration: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Raquel Conde- Álvarez, Ignacio Moriyón, Sue C. Welburn, Jakob Zinsstag.

Resources: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Raquel Conde- Álvarez, Ignacio Moriyón, Sue C. Welburn, Jakob Zinsstag.

Software: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Raquel Conde- Álvarez, Ignacio Moriyón, Ward Bryssinckx, Sue C. Welburn, Jakob Zinsstag.

Supervision: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Raquel Conde- Álvarez, Ignacio Moriyón, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

Validation: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Raquel Conde- Álvarez, Ignacio Moriyón, Pilar M. Muñoz Álvaro, Virginie Mick, Sue C. Welburn, Jakob Zinsstag.

Visualization: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Raquel Conde- Álvarez, Ignacio Moriyón, Amaia Zúñiga-Ripa, Pilar M. Muñoz Álvaro, Virginie Mick, Sue C. Welburn, Jakob Zinsstag.

Writing – original draft: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Raquel Conde- Álvarez, Ignacio Moriyón, Amaia Zúñiga-Ripa, Pilar M. Muñoz Álvaro, Virginie Mick, Ward Bryssinckx, Sue C. Welburn, Jakob Zinsstag.

Writing – review & editing: Hind Yahyaoui Azami, Marie J. Ducrotoy, Mohammed Bouslikhane, Jan Hattendorf, Mike Thrusfield, Raquel Conde- Álvarez, Ignacio Moriyón, Pilar M. Muñoz Álvaro, Sue C. Welburn, Jakob Zinsstag.

References

1. LES BRUCELLOSES ANIMALES [Internet]. [cited 1 Mar 2016]. http://www.onssa.gov.ma/fr/index.php?option=com_content&view=article&id=176&Itemid=117
2. Tuberculose bovine [Internet]. 2016 [cited 1 Mar 2016]. http://www.onssa.gov.ma/fr/index.php?option=com_content&view=article&id=181&Itemid=122
3. Narrod C, Zinsstag J, Tiongco M. A One Health Framework for Estimating the Economic Costs of Zoonotic Diseases on Society. *Ecohealth*. 2012; 9: 150–162. <https://doi.org/10.1007/s10393-012-0747-9> PMID: 22395956
4. Zinsstag J, Schelling E, Roth F, Bonfoh B, de Savigny D, Tanner M. Human Benefits of Animal Interventions for Zoonosis Control. *Emerg Infect Dis*. 2007; 13: 527–531. <https://doi.org/10.3201/eid1304.060381> PMID: 17553265
5. Morris RS. Diseases, dilemmas, decisions—Converting epidemiological dilemmas into successful disease control decisions. *Prev Vet Med*. 2015; 122: 242–252. <https://doi.org/10.1016/j.prevetmed.2015.05.003> PMID: 26072199
6. Cousins DV, Roberts JL. Australia's campaign to eradicate bovine tuberculosis: the battle for freedom and beyond. *Tuberculosis*. 2001; 81: 5–15. <https://doi.org/10.1054/tube.2000.0261> PMID: 11463220
7. Okello AL, Gibbs EPJ, Vandersmissen A, Welburn SC. One Health and the neglected zoonoses: turning rhetoric into reality. *Vet Rec*. 2011; 169: 281–285. <https://doi.org/10.1136/vr.d5378> PMID: 21908565
8. WHO. Tuberculosis Report. 2016.
9. Jones BA, Grace D, Kock R, Alonso S, Rushton J, Said MY, et al. Zoonosis emergence linked to agricultural intensification and environmental change. *Proc Natl Acad Sci U S A*. 2013; 110: 8399–8404. <https://doi.org/10.1073/pnas.1208059110> PMID: 23671097
10. Amanfu W. The situation of tuberculosis and tuberculosis control in animals of economic interest. *Tuberculosis*. 2006; 86: 330–335. <https://doi.org/10.1016/j.tube.2006.01.007> PMID: 16644282
11. Palmer MV. *Mycobacterium bovis*: characteristics of wildlife reservoir hosts. *Transbound Emerg Dis*. 2013; 60 Suppl 1: 1–13. <https://doi.org/10.1111/tbed.12115> PMID: 24171844
12. Pesciaroli M, Alvarez J, Boniotti MB, Cagiola M, Di Marco V, Marianelli C, et al. Tuberculosis in domestic animal species. *Res Vet Sci*. 2014; 97 Suppl: S78–85. <https://doi.org/10.1016/j.rvsc.2014.05.015> PMID: 25151859
13. Pollock JM, Rodgers JD, Welsh MD, McNair J. Pathogenesis of bovine tuberculosis: the role of experimental models of infection. *Vet Microbiol*. 2006; 112: 141–150. <https://doi.org/10.1016/j.vetmic.2005.11.032> PMID: 16384665
14. Cosivi O, Grange JM, Daborn CJ, Raviglione MC, Fujikura T, Cousins D, et al. Zoonotic tuberculosis due to *Mycobacterium bovis* in developing countries. *Emerg Infect Dis*. 1998; 4: 59–70. <https://doi.org/10.3201/eid0401.980108> PMID: 9452399
15. FAO. Principales réalisations depuis l'ouverture de la Représentation de la FAO à Rabat en 1982. 2011 Juillet. Report No.: ISBN 978-92-5-206938-6.
16. Bernués A, Manrique E, Maza MT. Economic evaluation of bovine brucellosis and tuberculosis eradication programmes in a mountain area of Spain. *Prev Vet Med*. 1997; 30: 137–149. [https://doi.org/10.1016/S0167-5877\(96\)01103-8](https://doi.org/10.1016/S0167-5877(96)01103-8) PMID: 9234417
17. Ministère Marocain de l'agriculture, du développement et des eaux et forêts. Arrêté du ministre de l'agriculture, du développement rural et des eaux et forêts n° 2017–01 du 19 chaabane 1422 (5 novembre 2001) relatif aux mesures complémentaires et spéciales pour lutter contre la tuberculose bovine. 2001 pp. 42–46.
18. Ducrot M, Ammary K, Ait Lbacha H, Zouagui Z, Mick V, Prevost L, et al. Narrative overview of animal and human brucellosis in Morocco: intensification of livestock production as a driver for emergence? *Infect Dis Poverty*. 2015; 4: 57. <https://doi.org/10.1186/s40249-015-0086-5> PMID: 26690090
19. Ducrot M, Bertu WJ, Matope G, Cadmus S, Conde-Álvarez R, Gusi AM, et al. Brucellosis in Sub-Saharan Africa: Current challenges for management, diagnosis and control. *Acta Trop*. 2017; 165: 179–193. <https://doi.org/10.1016/j.actatropica.2015.10.023> PMID: 26551794
20. Ocholi RA, Kwaga JKP, Ajogi I, Bale JOO. Abortion due to *Brucella abortus* in sheep in Nigeria. *Rev Sci Tech Int Off Epizoot*. 2005; 24: 973–979.
21. Corbel MJ, Alton G.G., Banai M., Díaz R., Dranovskaia B.A., Elberg S.S., et al. Brucellosis in Humans and Animals. World Health Organization; 2006.
22. MARA. Plan national de la lutte control la brucellose bovine (1989–1994). Morocco: Ministère de l'Agriculture et de la Reforme de l'Elevage, Direction de l'Elevage, Royaume du Maroc; 1989 pp. 15–35.

23. Abrak A, Boubia Y. Brucellose des ruminants domestiques: evaluation des plans de lutte actuels et perspectives d'eradication. Seminaire sur la vaccination contre la brucellose chez les ruminants; 2013 Février; Rabat.
24. Taoudi A, Fassi-Fehri MM, Johnson DW, Fagouri S. [Brucella abortus biotypes found in cattle in Morocco: a preliminary study]. *Dev Biol Stand.* 1983; 56: 123–128.
25. Jm Verger, Grayon M. Caractéristiques de 273 souches de *Brucella abortus* d'origine africaine. *Dev Biol Stand.* 1984; 56: 63–71. PMID: [6436118](#)
26. Benhabyles N, Benkirane A, Boudilmi B, Benchouk S, Bouayoun H. Epidemiologie de la brucellose humaine et animale au Maghreb. *CIHEAM Publ Neth No 1.* 1992;
27. MAMVA. Brucellose des petits ruminants: enquête épidémiologique dans l'oriental. Projet FAO: Ministère de l'Agriculture et de la Mise en Valeur agricole. Direction de l'élevage. 1996.
28. El Moudni Y. Actions du laboratoire régional d'analyses et de recherches vétérinaires d'Oujda dans l'épidémiologie de la brucellose ovine et caprine: Direction de l'élevage, Laboratoire Régional d'Analyses et de Recherches Vétérinaires d'Oujda. 1997;
29. MAPM. Province de Sidi Kacem. Division des Statistiques (DS), Direction de la Stratégie et des Statistiques (DSS), Ministère de l'Agriculture et de la Pêche Maritime. 2010.
30. Thrusfield M. *Veterinary Epidemiology.* Third edition. Blackwell Science Limited; 2007.
31. Otte M, Gumm I. Intra-cluster correlation coefficients of 20 infections calculated from the results of cluster-sample surveys. *Prev Vet Med.* 1997;
32. Thrusfield M. *Veterinary Epidemiology.* (fourth ed.) Blackwell Science Limited, USA; 2018.
33. OIE. *OIE Terrestrial Manual 2009, Bovine Tuberculosis.* 2009.
34. OIE. *Brucellosis in: Manual of Diagnostic Tests and Vaccines for Terrestrial Animals.* 2016.
35. Blasco JM, Garin-Bastuji B, Marin CM, Gerbier G, Fanlo J, Jiménez de Bagués MP, et al. Efficacy of different Rose Bengal and complement fixation antigens for the diagnosis of *Brucella melitensis* infection in sheep and goats. *Vet Rec.* 1994; 134: 415–420. PMID: [8036772](#)
36. Díaz-Aparicio E, Marín C, Alonso-Urmeneta B, Aragón V, Pérez-Ortiz S, Pardo M, et al. Evaluation of serological tests for diagnosis of *Brucella melitensis* infection of goats. *J Clin Microbiol.* 1994; 32: 1159–1165. PMID: [8051240](#)
37. Blasco JM, Marín C, Jiménez de Bagués M, Barberán M, Hernández A, Molina L, et al. Evaluation of allergic and serological tests for diagnosing *Brucella melitensis* infection in sheep. *J Clin Microbiol.* 1994; 32: 1835–1840. PMID: [7989528](#)
38. De Miguel MJ, Marín CM, Muñoz PM, Dieste L, Grilló MJ, Blasco JM. Development of a selective culture medium for primary isolation of the main *Brucella* species. *J Clin Microbiol.* 2011; 49: 1458–1463. <https://doi.org/10.1128/JCM.02301-10> PMID: [21270216](#)
39. Alton GG. *Techniques for the brucellosis laboratory.* Paris : Institut National de la Recherche Agronomique; 1988.
40. García-Yoldi D, Marín CM, de Miguel MJ, Muñoz PM, Vizmanos JL, López-Goñi I. Multiplex PCR assay for the identification and differentiation of all *Brucella* species and the vaccine strains *Brucella abortus* S19 and RB51 and *Brucella melitensis* Rev1. *Clin Chem.* 2006; 52: 779–781. <https://doi.org/10.1373/clinchem.2005.062596> PMID: [16595839](#)
41. Rogan WJ, Gladen B. Estimating prevalence from the results of a screening test. *Am J Epidemiol.* 1978; 107: 71–76. PMID: [623091](#)
42. Ducrotoy MJ, Muñoz PM, Conde-Álvarez R, Blasco JM, Moriyón I. A systematic review of current immunological tests for the diagnosis of cattle brucellosis. *Prev Vet Med.* 2018; 151: 57–72. <https://doi.org/10.1016/j.prevetmed.2018.01.005> PMID: [29496108](#)
43. Müller B, Vounatsou P, Ngandolo BNR, Diguimbaye-Djaïbe C, Schiller I, Marg-Haufe B, et al. Bayesian receiver operating characteristic estimation of multiple tests for diagnosis of bovine tuberculosis in Chadian cattle. *PLoS One.* 2009; 4: e8215. <https://doi.org/10.1371/journal.pone.0008215> PMID: [20011046](#)
44. OIE. *Bovine tuberculosis in: Manual of the Diagnostic Tests and Vaccines for Terrestrial Animals.* 2009.
45. Bernard F, Vincent C, Matthieu L, David R, James D. Tuberculosis and brucellosis prevalence survey on dairy cattle in Mbarara milk basin (Uganda). *Prev Vet Med.* 2005; 67: 267–281. <https://doi.org/10.1016/j.prevetmed.2004.11.002> PMID: [15748756](#)
46. Boukary AR, Thys E, Abatih E, Gamatié D, Ango I, Yenikoye A, et al. Bovine tuberculosis prevalence survey on cattle in the rural livestock system of Torodi (Niger). *PloS One.* 2011; 6: e24629. <https://doi.org/10.1371/journal.pone.0024629> PMID: [21961039](#)

47. Dejene SW, Heitkönig IMA, Prins HHT, Lemma FA, Mekonnen DA, Alemu ZE, et al. Risk Factors for Bovine Tuberculosis (bTB) in Cattle in Ethiopia. *PLoS One*. 2016; 11: e0159083. <https://doi.org/10.1371/journal.pone.0159083> PMID: 27404387
48. Katala BZ, Mbugi EV, Karimuribo ED, Keyyu JD, Kendall S, Kibiki GS, et al. Prevalence and risk factors for infection of bovine tuberculosis in indigenous cattle in the Serengeti ecosystem, Tanzania. *BMC Vet Res*. 2013; 9: 267. <https://doi.org/10.1186/1746-6148-9-267> PMID: 24377705
49. Firdessa R, Tschopp R, Wubete A, Sombo M, Hailu E, Erenso G, et al. High Prevalence of Bovine Tuberculosis in Dairy Cattle in Central Ethiopia: Implications for the Dairy Industry and Public Health. *PLOS ONE*. 2012; 7: e52851. <https://doi.org/10.1371/journal.pone.0052851> PMID: 23285202
50. Munyeme M, Muma JB, Skjerve E, Nambota AM, Phiri IGK, Samui KL, et al. Risk factors associated with bovine tuberculosis in traditional cattle of the livestock/wildlife interface areas in the Kafue basin of Zambia. *Prev Vet Med*. 2008; 85: 317–328. <https://doi.org/10.1016/j.prevetmed.2008.03.006> PMID: 18455816
51. Moiane I, Machado A, Santos N, Nhambir A, Inlamea O, Hattendorf J, et al. Prevalence of Bovine Tuberculosis and Risk Factor Assessment in Cattle in Rural Livestock Areas of Govuro District in the Southeast of Mozambique. *PLOS ONE*. 2014; 9: e91527. <https://doi.org/10.1371/journal.pone.0091527> PMID: 24632593
52. Greiner M, Verloo D, de Massis F. Meta-analytical equivalence studies on diagnostic tests for bovine brucellosis allowing assessment of a test against a group of comparative tests. *Prev Vet Med*. 2009; 92: 373–381. <https://doi.org/10.1016/j.prevetmed.2009.07.014> PMID: 19766334
53. EFSA. Opinion of the Scientific Panel on Animal Health and Welfare (AHAW) on a request from the Commission concerning Brucellosis Diagnostic Methods for Bovines, Sheep, and Goats: Opinion of the Scientific Panel on Animal Health and Welfare (AHAW) on a request from the Commission concerning Brucellosis Diagnostic Methods for Bov. *EFSA J*. 2007; 5: 432. <https://doi.org/10.2903/j.efsa.2007.432>
54. Ducrotoy MJ, Conde-Álvarez R, Blasco JM, Moriyón I. A review of the basis of the immunological diagnosis of ruminant brucellosis. *Vet Immunol Immunopathol*. 2016; 171: 81–102. <https://doi.org/10.1016/j.vetimm.2016.02.002> PMID: 26964721
55. Ferreira AC, Cardoso R, Dias IT, Mariano I, Belo A, Preto IR, et al. Evaluation of a modified Rose Bengal test and an indirect Enzyme-Linked Immunosorbent Assay for the diagnosis of *Brucella melitensis* infection in sheep. *Vet Res*. 2003; 34: 297–305. <https://doi.org/10.1051/vetres:2003005> PMID: 12791239
56. Caprine and ovine brucellosis (excluding *Brucella ovis*), in: *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals (OIE)*. 2009.
57. Nielsen K. Diagnosis of brucellosis by serology. *Vet Microbiol*. 2002; 90: 447–459. [https://doi.org/10.1016/S0378-1135\(02\)00229-8](https://doi.org/10.1016/S0378-1135(02)00229-8) PMID: 12414164
58. Praud A, Durán-Ferrer M, Fretin D, Jaÿ M, O'Connor M, Stournara A, et al. Evaluation of three competitive ELISAs and a fluorescence polarisation assay for the diagnosis of bovine brucellosis. *Vet J Lond Engl 1997*. 2016; 216: 38–44. <https://doi.org/10.1016/j.tvjl.2016.06.014> PMID: 27687924
59. Lucchese L, Benkirane A, Hakimi I, El Idrissi A, Natale A. Seroprevalence study of the main causes of abortion in dairy cattle in Morocco. *Vet Ital*. 2016; 52: 13–19. <https://doi.org/10.12834/VetIt.388.1813.1> PMID: 27033527
60. El-Diasty MM, Ahmed HA, Sayour AE, El Hofy FI, Tahoun ABMB, Shafik SM. Seroprevalence of *Brucella* spp. in Cattle, Molecular Characterization in Milk, and the Analysis of Associated Risk Factors with Seroprevalence in Humans, Egypt. *Vector Borne Zoonotic Dis Larchmt N*. 2016; 16: 758–764. <https://doi.org/10.1089/vbz.2016.1985> PMID: 27754795
61. Makita K, Fèvre EM, Waiswa C, Eisler MC, Thrusfield M, Welburn SC. Herd prevalence of bovine brucellosis and analysis of risk factors in cattle in urban and peri-urban areas of the Kampala economic zone, Uganda. *BMC Vet Res*. 2011; 7: 60. <https://doi.org/10.1186/1746-6148-7-60> PMID: 22004574
62. Mohammed FU, Ibrahim S, Ajogi I, Olaniyi BJO. Prevalence of Bovine Brucellosis and Risk Factors Assessment in Cattle Herds in Jigawa State. *Int Sch Res Not*. 2011; 2011: e132897. <https://doi.org/10.5402/2011/132897> PMID: 23738094
63. Jergefa T, Kelay B, Bekana M, Teshale S, Gustafson H, Kindahl H. Epidemiological study of bovine brucellosis in three agro-ecological areas of central Oromiya, Ethiopia. *Rev Sci Tech Int Off Epizoot*. 2009; 28: 933–943.
64. Benkirane Abdelali, Essamkaoui Soukaina, El Idrissi Ahmed, Lucchese Laura, Natale Alda. A sero-survey of major infectious causes of abortion in small ruminants in Morocco. *Vet Ital*. 2015; 25–30. <https://doi.org/10.12834/VetIt.389.1814.1> PMID: 25842210
65. Bertu WJ, Ducrotoy MJ, Muñoz PM, Mick V, Zúñiga-Ripa A, Bryssinckx W, et al. Phenotypic and genotypic characterization of *Brucella* strains isolated from autochthonous livestock reveals the dominance

- of *B. abortus* biovar 3a in Nigeria. *Vet Microbiol.* 2015; 180: 103–108. <https://doi.org/10.1016/j.vetmic.2015.08.014> PMID: 26315770
66. Ducrotoy MJ, Bertu WJ, Ocholi RA, Gusi AM, Bryssinckx W, Welburn S, et al. Brucellosis as an Emerging Threat in Developing Economies: Lessons from Nigeria. *PLoS Negl Trop Dis.* 2014; 8: e3008. <https://doi.org/10.1371/journal.pntd.0003008> PMID: 25058178
 67. Brooks-Pollock E, Conlan AJ, Mitchell AP, Blackwell R, McKinley TJ, Wood JL. Age-dependent patterns of bovine tuberculosis in cattle. *Vet Res.* 2013; 44: 97. <https://doi.org/10.1186/1297-9716-44-97> PMID: 24131703
 68. Joseph OA, Oluwatoyin AV, Comfort AM, Judy S, Babalola CSI. Risk factors associated with brucellosis among slaughtered cattle: Epidemiological insight from two metropolitan abattoirs in Southwestern Nigeria. *Asian Pac J Trop Dis.* 2015; 5: 747–753. [https://doi.org/10.1016/S2222-1808\(15\)60925-2](https://doi.org/10.1016/S2222-1808(15)60925-2)
 69. Cook AJC, Tuchiili LM, Buve A, Foster SD, Godfrey-Faussett P, Pandey GS, et al. Human and bovine tuberculosis in the Monze District of Zambia—a cross-sectional study. *Br Vet J.* 1996; 152: 37–46. PMID: 8634864
 70. Griffin JM, Haheisy T, Lynch K, Salman MD, McCarthy J, Hurley T. The association of cattle husbandry practices, environmental factors and farmer characteristics with the occurrence of chronic bovine tuberculosis in dairy herds in the Republic of Ireland. *Prev Vet Med.* 1993; 17: 145–160.
 71. Costello E, Doherty ML, Monaghan ML, Quigley FC, O'Reilly PF. A study of cattle-to-cattle transmission of *Mycobacterium bovis* infection. *Vet J.* 1998; 155: 245–250. PMID: 9638070
 72. Kazwala RR, Kambarage DM, Daborn CJ, Nyange J, Jiwa SFH, Sharp JM. Risk factors associated with the occurrence of bovine tuberculosis in cattle in the Southern Highlands of Tanzania. *Vet Res Commun.* 2001; 25: 609–614. PMID: 11767006
 73. Humblet M-F, Boschioli ML, Saegerman C. Classification of worldwide bovine tuberculosis risk factors in cattle: a stratified approach. *Vet Res.* 2009; 40: 1–24.
 74. Collins JD. Tuberculosis in cattle: strategic planning for the future. *Vet Microbiol.* 2006; 112: 369–381. <https://doi.org/10.1016/j.vetmic.2005.11.041> PMID: 16330164
 75. Radunz B. Surveillance and risk management during the latter stages of eradication: Experiences from Australia. *Vet Microbiol.* 2006; 112: 283–290. <https://doi.org/10.1016/j.vetmic.2005.11.017> PMID: 16321479
 76. Ducrotoy M, Bertu WJ, Matope G, Cadmus S, Conde-Álvarez R, Gusi AM, et al. Brucellosis in Sub-Saharan Africa: Current challenges for management, diagnosis and control. *Acta Trop.* 2015; <https://doi.org/10.1016/j.actatropica.2015.10.023> PMID: 26551794
 77. Abakar F, Yahyaoui Azami H, Justus Bless P, Crump L, Lohmann P, Laager M, et al. Transmission Dynamics and Elimination Potential of Zoonotic Tuberculosis in Morocco. *PLoS Negl Trop Dis.* 2016;
 78. Collins JD. Tuberculosis in cattle: new perspectives. *Tuberculosis.* 2001; 81: 17–21. <https://doi.org/10.1054/tube.2000.0262> PMID: 11463221
 79. Sapolsky RM, Share LJ. A Pacific Culture among Wild Baboons: Its Emergence and Transmission. *PLOS Biol.* 2004; 2: e106. <https://doi.org/10.1371/journal.pbio.0020106> PMID: 15094808
 80. Woodford MH. Tuberculosis in wildlife in the Ruwenzori National Park Uganda (part I). *Trop Anim Health Prod.* 1982; 14: 81–88. PMID: 7201688
 81. Ayele WY, Neill SD, Zinsstag J, Weiss MG, Pavlik I. Bovine tuberculosis: an old disease but a new threat to Africa. *Int J Tuberc Lung Dis Off J Int Union Tuberc Lung Dis.* 2004; 8: 924–937.
 82. Maudlin I, Eisler MC, Welburn SC. Neglected and endemic zoonoses. *Philos Trans R Soc Lond B Biol Sci.* 2009; 364: 2777–2787. <https://doi.org/10.1098/rstb.2009.0067> PMID: 19687045
 83. Welburn SC, Beange I, Ducrotoy MJ, Okello AL. The neglected zoonoses—the case for integrated control and advocacy. *Clin Microbiol Infect.* 2015; 21: 433–443. <https://doi.org/10.1016/j.cmi.2015.04.011> PMID: 25911990
 84. Welburn SC, Bardosh KL, Coleman PG. Novel Financing Model for Neglected Tropical Diseases: Development Impact Bonds Applied to Sleeping Sickness and Rabies Control. *PLoS Negl Trop Dis.* 2016; 10: e0005000. <https://doi.org/10.1371/journal.pntd.0005000> PMID: 27855156
 85. Welburn SC, Coleman PG, Zinsstag J. Rabies Control: Could Innovative Financing Break the Deadlock? *Front Vet Sci.* 2017; 4. <https://doi.org/10.3389/fvets.2017.00032> PMID: 28337440
 86. Schelling E, Diguimbaye C, Daoud S, Nicolet J, Boerlin P, Tanner M, et al. Brucellosis and Q-fever seroprevalences of nomadic pastoralists and their livestock in Chad. *Prev Vet Med.* 2003; 61: 279–293. PMID: 14623412