

Asteroid trace fossils from Lower Cretaceous shallow- to marginal-marine deposits in Patagonia

Diana E. Fernández^{1,2*}, Marcos Comerio³, Luciana Giachetti¹, Pablo J. Pazos^{1,2}, Andreas Wetzel⁴

¹ Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales. Departamento de Ciencias Geológicas. Ciudad Universitaria, Buenos Aires, Argentina.

elizabeth@gl.fcen.uba.ar; giachetti.luciana@gmail.com; pazos@gl.fcen.uba.ar

² CONICET - Universidad de Buenos Aires. Instituto de Estudios Andinos Don Pablo Groeber (IDEAN). Ciudad Universitaria, Buenos Aires, Argentina

³ Centro de Tecnología de Recursos Minerales y Cerámica (CETMIC). CONICET. Manuel B. Gonnet, Argentina. marcoscomerio@gmail.com

⁴ Geologisches Institut der Universität Basel, Bernoullistrasse 32, CH-4056 Basel, Switzerland. andreas.wetzel@unibas.ch

* Corresponding author

Keywords: *Asteriacites*, Asteroids, Palaeoecology, Predator-prey interaction, Ichnotaxonomy Morphometry

Published in: Cretaceous Research, 93: 120–128

<https://doi.org/10.1016/j.cretres.2018.09.010>

Abstract

Most stellate trace fossils of the ichnogenus *Asteriacites* are attributed to asterozoan producers in general and the majority is the result of the work of ophiuroids. The fossil record of asterozoans is scarce in South America, particularly for the Mesozoic. *Asteriacites* specimens found in shallow- to marginal-marine Lower Cretaceous (upper Hauterivian-lower Barremian) deposits in the Neuquén Basin (Patagonia, Argentina) exhibit sculpture and morphometry typical of asteroid producers. This is the second record of asteroids from the Lower Cretaceous of South America. The close association between these *Asteriacites* possibly produced by astropectinids and traces assignable to Siphonichnidae are suggestive of a predator-prey interaction, adding palaeo-ecological information for community-structure reconstruction of these deposits. For ichno-taxonomic evaluation, morphometric parameters of *Asteriacites* were elaborated using simple photogrammetric procedures applied on negative epirelief specimens and undertraces to define edges of the stellate trace fossils.

1. Introduction

The ichnogenus *Asteriacites* von Schlotheim, 1820 includes stellate trace fossils (Seilacher, 1953; Häntzschel, 1970). Most records of *Asteriacites* are attributed to asterozoans as producers in general and the majority is the result of the work of ophiuroids (Mángano et al., 1999, and references therein). Such trace fossils provide a valuable record of asterozoan activities (see recent review by Knaust and Neumann, 2016), because the producing organisms tend to disarticulate after death very rapidly (e.g., Villier et al., 2004). Especially, for South America, the asterozoan record is scarce for both body and trace fossils (e.g., Martín-Medrano and García-Barrera, 2013; Fernández et al., 2014; Martínez and del Río, 2015). So far, asterozoan trace fossils are known from two units of the Neuquén Basin, the Jurassic Las Lajas Formation, where the presence of *Asteriacites* has only been mentioned (McIlroy et al., 2005), and the Cretaceous Mulichinco Formation where *Asteriacites lumbricalis* has been found, described and attributed to ophiuroids (Rodríguez et al., 2007; Fernández, 2013).

The new finds of *Asteriacites* in the shallow- to marginal-marine, upper Hauterivian-lower Barremian Agua de la Mula Member (Agrido Formation, Neuquén Basin, Patagonia), thus, provides new data to the asterozoan record in general and in particular to the community structure of these deposits. In addition, excellently preserved primary sedimentary structures help to decipher the environmental setting of the Agua de la Mula Member in more detail and the habitat of the *Asteriacites* producers in general. It is the purpose of this paper (i) to describe these specimens in detail, (ii) to evaluate their ichnotaxonomy, (iii) to outline their palaeoecological implications, and (iv) to refine the palaeobathymetric interpretation of the studied deposits.

2. Geological and palaeontological setting

The Neuquén Basin is located in west-central Argentina between 34° and 41°S (Fig. 1A). It represents an epicontinental basin in back-arc position to the Palaeo-PacificAndean subduction zone (e.g., Howell et al., 2005; Zapata and Folguera, 2005). It contains more than 7000 m thick marine and continental deposits being Late Triassic to Palaeogene in age (Vergani et al., 1995; Legarreta and Uliana, 1999). Most of the Jurassic and Early Cretaceous deposits formed in marine settings while temporarily enhanced subsidence and sea-level fluctuations accentuated the diverse, fossiliferous facies (Howell et al., 2005).

The Agua de la Mula Member (Leanza et al., 2001; Fig. 1B) of the Agrido Formation (Weaver, 1931) was studied in detail in the Neuquén Province. It is late Hauterivian to early Barremian in age (Aguirre-Urreta et al., 2007, 2008, 2015; Aguirre-Urreta and Rawson, 2012). The depositional environment of the Agua de la Mula Member has mainly been interpreted as an open marine ramp (Spalletti et al., 2001a; Lazo et al., 2005; Ballent et al., 2006). However, marginal-marine intervals

are recurrently present, particularly in the upper part of the unit (Pazos and Fernández, 2010; Fernández and Pazos, 2012, 2013; Pazos et al., 2012).

The Agua de la Mula Member was studied in detail at Bajada del Agrio, the type locality of the Agrio Formation, located between 38°25'S/70°00'W and 38°26'S/70°01'W (Figs. 1 and 2). There, the entire unit is 470 m thick. At Bajada del Agrio, the upper part of the Agrio Formation is interpreted to have formed in shallow subtidal to proximal outer ramp sub-environments influenced by fair-weather and storm waves (e.g., Spalletti et al., 2001b). The base of the logged section is located at GPS coordinates 38°25'23"S/70°00'42"W. The studied interval coincides with the uppermost part of the stratigraphic sequence Sq3 and the lowermost part of Sq4 of Guler et al. (2013). Within an inner ramp setting, it occupies a transitional position between basically shoreface deposits with intervals of upper shoreface sediments (Fernández et al., 2018) and shallow to marginal-marine sediments comprising tidal-flat deposits affected by enhanced and/or fluctuating salinity (Fernández and Pazos, 2012). Tides influenced the depositional area considerably as observed in other localities (Pazos et al., 2012) rather than storms as traditionally proposed (e.g., Spalletti et al., 2001b). The marginal-marine, tidally influenced deposits constitute the uppermost 63 m of the unit.




Fig. 1. Neuquén Basin and stratigraphic column of the Agua de la Mula Member (Agrio Formation). (A) Regional map of the Neuquén Basin in west-central Argentina showing outcrops of the Agrio Formation (modified from Aguirre-Urreta et al., 2015). (B) Generalized stratigraphic column of the Agua de la Mula Member. Note the studied interval, approximately 340 m above the contact with the underlying Avilé Member (Agrio Formation). Modified from Aguirre-Urreta et al. (2007).




Figure 2.

Logged section beginning 340 m above the base of the Agua de la Mula Member at Bajada del Agrio and ending at an oolitic limestone level about 60 m below the base of the Huitrin Fm. (see Fernández and Pazos, 2012). Depositional sequences taken from Guler et al. (2013).

The Agua de la Mula Member contains varied and abundant macrofossils including bivalves, corals, ammonoids, gastropods, serpulids, sponges, echinoids, decapods, etc. (e.g., Cichowolski, 2003; Lazo et al., 2005, 2009; Rodríguez, 2007; Taylor et al., 2009; Aguirre-Urreta et al., 2011; Cataldo, 2013; Luci et al., 2013). Some groups are only known through their trace fossils (e.g. xiphosurids; Fernández and Pazos, 2013). So far, there is no record of asteroids in the unit.

3. Material and methods

The logged section begins 340 m above the transgressive base of the Agua de la Mula Member and ends in an oolitic-skeletal bar (Figs. 1B and 2). The latter represents the first level of the section analyzed in detail by Fernández and Pazos (2012). To provide a sedimentological and environmental framework, in total 73 m of over- and underlying strata were logged (Figs. 2 and 3). While the Agua de la Mula Member is considered as homoclinal siliciclastic carbonate ramp (e.g. Lazo et al., 2005; Guler et al., 2013), the depositional settings were accordingly classified based on the scheme of Burchette and Wright (1992).

Two specimens of stellate trace fossils were studied in detail (see field photographs on Figs. 3D and 4). Specimens A and B were found at 17.8 m and 18.6 m of the logged section, respectively (Fig. 2). Because of their occurrence in the field, the specimens were not recoverable. For enhancement of certain morphologic characteristics of the trace fossils, the images were processed with the photogrammetric software Agisoft PhotoScan Professional 1.0.4 build 1847 (64 bit), using standard methodology (e.g. Mallison and Wings, 2014). 3D meshes with texture overprint were used to enhance the imprints within each arm (Fig. 4E). 3D surfaces with shaded relief and contour lines as well as different profiles perpendicular to the arm axes were created using the software Golden Surfer (Fig. 4D). Given the scarcity of specimens the photogrammetric analyses were only accessory in the description as visual aid to better define the edge of the trace fossils (Fig. 4E). Morphometric analysis and ichnotaxonomic evaluation is based on Knaust and Neumann (2016).

4. Depositional setting

4.1. Observations

The studied section is characterized by coarsening- and shallowing-upward packages (Figs. 2 and 3). They range from 4 to 10 m in thickness (Figs. 2 and 3A) and are considered as parasequences in a sequence stratigraphic framework (e.g., Cichowolski et al., 2012). These packages have thin bioclastic conglomerates at the base, indicating episodes of reduced siliciclastic input. Above follow massive or laminated mudstones, thin lenticular and wavy bedding and finally fine-grained sandstones either massive or with wave-ripple lamination. Within the sandstone interval, occasionally beds with current-ripple lamination, swaley cross-stratification, and in some cases syndepositional deformation structures are present. At the top of a package, sandstone bed surfaces may exhibit wrinkle structures (Fig. 3B), interference ripples and parting lineation. Some ripples are flat-topped (Fig. 3E and F).

The *Asteriacites* specimens were found in two levels with dominant wave-ripple lamination (Fig. 3C and D) in the package beginning at 14.5 m above the base of the logged section (Fig. 2). Besides *Asteriacites*, other trace fossils occur in the logged section, such as cf. *Arenicolites* Salter, 1857, *Bolonia lata* Meunier, 1886, *Chondrites ?intricatus* (Brongniart, 1823), *Gyrochorte comosa* Heer, 1865, *Rhizocorallium* Zenker, 1836, *Siphonichnus* Stanistreet et al., 1980, tetrapod swimming traces, arthropod trackways and invertebrate trails. *Chondrites*, *Gyrochorte* and *Rhizocorallium* are also present in the overlying interval on top of the Agua de la Mula Member (Fernández and Pazos, 2012).




Fig. 3. Examples of sedimentary features of the logged section. (A) Example of coarsening- and shallowing-upward packages. Graphic scale = 1 m. (B) Wrinkle structures. Graphic scale = 1 cm. (C) Wave ripples. Graphic scale = 6 cm. (D) *Asteriacites*, specimen A, and associated ripples. Graphic scale = 2 cm. (E) and (F) Flat-topped ripples. Rock pick = 33 cm long. Coin = 2.1 cm in diameter.

4.2. Interpretation

The logged section is interpreted to represent marginal-marine deposits accumulated in a proximal mid-ramp to inner ramp setting (cf. Burchette and Wright, 1992). Wave ripples indicate a bathymetric range above the fair-weather wave base, while swaley cross-stratification is a consequence of storm events in rather shallow water (e.g., Dumas and Arnott, 2006). The massive sandstone beds are either the result of intense indistinct bioturbation or rapid deposition from decelerating flows or wave pumping in association or not to soft-sediment deformation (e.g., Gingras et al., 2014; Liu et al., 2017). Intense bioturbation is the most likely process in intervals without syndepositional deformation. The heterolithic deposits are interpreted to have accumulated in a shallowing-upward setting characterized by mixed traction and suspension deposition forming first lenticular and then wavy bedding with oscillatory ripples that become replaced by current ripples (e.g., Reineck and Singh, 1980). The association of wrinkle structures with interference and flat-topped ripples suggests a marginal-marine setting. Interference ripples are typical of a very shallow depositional setting above the fair-weather wave base, while the flat-topped ripples reflect temporary subaerial exposure (e.g., Reineck and Singh, 1980). Although wrinkle structures occur within a wide palaeobathymetric range (e.g., Mata and Bottjer, 2009), they are most common in marginal-marine environments with shallow water experiencing intermittent exposure, such as tidal flats (e.g., Allen, 1984; Hagadorn and Bottjer, 1997). In addition, wrinkle structures have also been reported from a marine-marginal setting at the top of the Agua de la Mula Member exposed in another locality. There, clear evidence of tidal influence and temporary subaerial exposure have been observed (Fernández and Pazos, 2013). Indicators of subaerial exposure are present in three levels within the logged section, but not in the *Asteriacites*-bearing levels. Consequently, these intervals were formed under shallow subtidal conditions above fair-weather wave base.

5. Ichnology

5.1. Ichnogenus *Asteriacites* von Schlotheim, 1820

Type ichnospecies. *Asteriacites lumbricalis* von Schlotheim, 1820; by subsequent designation of Seilacher (1953).

Diagnosis. Star-shaped, commonly pentamerous imprints and shallowly excavated pits with arms grading from a central area and continuously tapering distally (modified by Knaust and Neumann, 2016).

Remarks. The latest revision of *Asteriacites* applied hierarchically ordered ichnotaxobases: The overall appearance defines the ichnofamily Asteriacitidae, while the basic morphology differentiates *Asteriacites* from related ichnogenera (Knaust and Neumann, 2016). Sculpture was commonly applied as an ichnospecific ichnotaxobase (e.g., Seilacher, 1953; Crimes and Crossley, 1991), but due to its possible taphonomic bias it is now considered to be of subordinate value compared with other morphometric data such as the length/width ratio of the arms (Knaust and Neumann, 2016).

Description. Two specimens of pentamerous imprints preserved as negative epirelief (specimens A and B; Fig. 4). The sand laminae wherein the imprints are preserved are approximately 2–3 mm thick, while the whole bed in both cases is 3–4 cm thick. The arms are mostly straight except a slight curvature of one of them (Fig. 4D). The length from the central point to the tip of each individual arm is between 54.1 and 64.5 mm in specimen A (Fig. 4A), and 51.2 and 60.7 mm in specimen B (Fig. 4C). The largest width of the arms measured at their base varies slightly in each arm between 28.8 and 30 mm in specimen A and between 18.1 and 27.5 mm in specimen B. Some of the arms appear diffuse at their tips. The cross-sectional geometry of the arms is roughly V-shaped (Fig. 4D). Within each specimen, two areas are defined. The inner area close to the median line of an arm is 2–4.5 mm deep, sculptured, and rather regular in outline, while the outer area is

less deep (1–2 mm), unornamented or finely striated, and less regular in outline (Fig. 4B). The imprints in both areas are oriented perpendicular to oblique to the arm axes.

5.2. Associated trace fossils

In the levels where the *Asteriacites* specimens occur, *Bolonia lata* and *Gyrochorte comosa* are common (Fig. 2). While *Gyrochorte comosa* is quite similar to those found further up in the section, as described by Fernández and Pazos (2012), *Bolonia lata* is reported for the first time from this unit. However, a detailed description of these trace fossils is beyond the scope of this study. Vertical burrows with a circular to oval cross-section, composed of a central core and surrounded by a concentric mantle, are assigned to the ichnogenus *Siphonichnus* (e.g., Knaust, 2015). These burrows are found closely associated with the *Asteriacites* traces (Fig. 4A, D, E). *Asteriacites* and the associated burrows do not cross-cut each other.




Fig. 4. Field photographs of *Asteriacites* specimens and examples of image analyses used as visual aid for the description. Graphic scales = 1 cm in A, C and D, 2 mm in B and E. (A) *Asteriacites*, specimen A. Si = trace fossil assignable to *Siphonichnus*. Si? = trace fossil assigned to *Siphonichnus* with doubts. (B) Detail of the rectangle in A showing part of the arm imprint; line drawing outlining striation features. WS = wide striations, RS = medium-sized striations, FS = fine striations. The first two types of striae are restricted to the inner area of the arm imprint (as defined in text), while fine striations are found in the outer area. (C) Specimen B. (D) Shaded grey 3D photogrammetric model of specimen A and example of a relative morphometric profile (location marked with white line) perpendicular to the arm axis. The Xs mark the areas of the profile that helped defined the trace boundary. (E) Detail of the white rectangle marked in D showing example of delimitation of the trace boundary.

5.3. Ichnotaxonomic and preservational discussion

Three ichnospecies of *Asteriacites* are regarded as valid (Knaust and Neumann, 2016): *A. lumbricalis* von Schlotheim, 1820, *A. stelliformis* (Miller and Dyer, 1878), and *A. quinquefolius* (Quenstedt, 1876). *A. lumbricalis* is characterised by slender arms distinct from a central disc-shaped area (Knaust and Neumann, 2016), a feature not observed in the studied specimens. *A. stelliformis* presents arms having a length:width ratio >2 , while in *A. quinquefolius* this ratio is <2 (Knaust and Neumann, 2016). The morphometric analysis of the studied specimens shows that for the arms the length:width ratio varies between 1.8 and 2.9. For 7 of the 10 arms the length:width ratio matches the value typical of *A. stelliformis* (>2). In specimen B the length:width ratio is >2 for every arm, but for specimen A three arms exhibit a length:width ratio <2 being characteristic for *A. quinquefolius*. In addition, the arm sculpture with respect to morphology and distribution of imprints resembles more that of the holotype of *A. quinquefolius* (Seilacher, 1953, pl. 10, fig. 2), while the striae are not continuous as typical of *A. lumbricalis* (Seilacher, 1953, 2007; Mángano et al., 1999; Knaust and Neumann, 2016; Ishida et al., 2017). Based on the morphometric parameters of the current ichnospecies diagnoses, specimen A is assigned to *A. stelliformis*, while specimen B could be assigned to *A. stelliformis* or *A. quinquefolius*. However, because of the sculpture and while the majority of the arms shows a length:width ratio <2 , an assignment to *A. quinquefolius* is suggested. It has been found in other *Asteriacites* that the width:length ratio differs between the arms of the same specimen, suggesting two different ichnospecific assignments (Singh et al., 2017).

The *Asteriacites* traces are preserved as negative epirelief in two separate sandstone beds showing dominance of wave-ripple lamination (Fig. 2). They are covered by thin mudstone layers. In neoichnological experiments with burrowing asteroids (astropectinids), rather indistinct depressions are formed on the sand surface, while the undertraces exhibit the structures more clearly (Seilacher, 1953). The outer arm area observed in the studied specimens (Fig. 4B) resembles that of the surface impression shown by Seilacher (1953, fig. 1a), while a rather sharp sculpture of the inner area resembles that of the undertrace. Furthermore, in aquarium well-preserved surface impressions show small mounds of sediment on both sides of the arms (e.g., Ishida et al., 2017). However, such structures are not present in the studied traces. They are, therefore, considered to represent an intermediate case between the two preservational types described by Seilacher (1953). Because the depth of the epirelief (3–5 mm) exceeds the thickness of individual sand laminae (1–2 mm) that formed during wave action, it is possible that the upper part (outer area) of the trace was affected by sand deposition during trace production and represents the original surface imprint, while the lower part (inner area) remained nearly unaffected or even could represent an undertrace. In the parts where the outer areas are unornamented, redeposition of sand by wave-pumping effects (Liu et al., 2017) and/or microavalanches due to originally unstable parts of the flanks cannot be ruled out as processes that “smoothed” the original profile and/or sculpture. In other cases, where specimens assignable to *A. stelliformis* are attributed to asteroids (e.g., Ishida et al., 2013, fig. 3, specimen with arm length:width ratio >2), preservation as shallow undertrace would explain nicely the high length:width ratio of arm imprints, which would be lower for surface imprints of the same specimen.

6. Palaeoecological implications

Asteriacites is attributed to both asteroids and ophiuroids as producers. Usually considered a resting trace, it documents shallow burrowing. The movements of arms and tube feet might result in arm impressions wider than the producing body parts, and distinctive sculpture inside the trace, respectively (Seilacher, 1953, 2007). Asteroids usually have too broad arms to move the sediment from one side of the arm to the other by a single tube foot. Seilacher (1953) described the imprints of the tube feet of asteroids as discontinuous and irregular while ophiuroids produce transverse and

continuous striae. Recent neoichnological experiments focused on tube-feet movement and the resultant striations ornamenting the traces of asteroids and ophiuroids (Ishida et al., 2017).

For example, in the case of asteroids, in particular astropectinids, after forming a shallow burrow and starting to move again in a particular direction, the striations differ within each arm depression. The striations are wider and more regular in the half of the arm imprint opposite to the direction of starfish movement, while they are fine and irregular in the other half. In the case of ophiuroids, the striations are fine, regular and continuous from one side of the arm to the other. In both cases, the striations are mostly perpendicular to the arm axis (Ishida et al., 2017).

The sculpture of the studied specimens resembles that resultant from the work of tube feet of asteroids. Along each arm, areas with different ornamentation are found (see above). While the striations are mostly perpendicular to the arm axis, they are not continuous (Fig. 4B). In neoichnological experiments, tube-feet movement was observed on the surface of the substratum as well as on the ventral and lateral sides of the glass beaker, and striations form at different depths (Ishida et al., 2017). Thus, two areas having different striations are likely to be present in shallow undertraces.

A. stelliformis can be produced by both ophiuroids and asteroids, but *A. quinquefolius* is interpreted to be produced only by asteroids (Knaust and Neumann, 2016). In summary, morphometric data and sculpture of the *Asteriacites* in particular the arm-width:arm-length ratio point to asteroids as producers of the studied specimens. They were produced in very shallow marine, subtidal environments as recorded in the *Asteriacites*-bearing levels.

So far there is only one record of asteroids from the Neuquén Basin from the Mulichinco Formation and was assigned to *Tethyaster antares* Fernández et al., 2014, belonging to the family Astropectinidae. Comparison of morphometric data between asteroid body fossils and trace fossils has been used as tool for identification of possible producers (Ishida et al., 2013). The armlength:arm-width ratio of *T. antares* is between 2 and 2.26. When compared to the ratio given for asterozoan trace fossils by Knaust and Neumann (2016) it mostly coincides with that typical of *A. stelliformis* (>2), but it is also within the range of *A. quinquefolius* (<2). Therefore, the only asteroid species known from the Lower Cretaceous, in particular early Valanginian of the Neuquén Basin (Fernández et al., 2014), has a similar arm-length:arm-width ratio as the studied asterozoan trace fossils found in the Agrio Formation (late Valanginian-early Barremian).

Extant astropectinids and paxillosidans in general are semiinfaunal, capable of self-burial in soft sandy or muddy substrates and are common in marine settings from intertidal areas to the deep sea (Heddle, 1967; Blake and Aronson, 1998; Jagt, 2000; Byrne and O'Hara, 2017). These starfish are voracious predators that usually feed on other invertebrates like bivalves and worms. They search for prey and shallowly dig into the substrate to get it while extruding the stomach (Ortega et al., 2011 and references therein). Self-burial of asteroids evolved in relation to the presence of fascioles that allow unobstructed water flow near the animals's body (Blake and Aronson, 1998; Blake and Reid, 1998). These structures were already present in Valanginian times (Fernández et al., 2014). Therefore, the producers of the studied *Asteriacites* were capable to burrow. Consequently, the studied *Asteriacites* could be associated with different behaviour types: (i) not-sustained resting (cubichnia) while shallowly burrowing; (ii) temporally prolonged shallow burial, typical of a shallow semi-infaunal life style; (iii) active searching for prey within the uppermost centimeters of the substrate; or (iv) combination of (i) to (iii).

Trace fossils of the ichnofamily Siphonichnidae are common in marginal-marine and shallow-marine deposits. They are produced by infaunal bivalves adjusting their position within the sediment (Knaust, 2015 and references therein). The presence of these trace fossils in close association with *Asteriacites* are suggestive of a behaviour of the *Asteriacites* tracemaker to burrow shallowly to locate prey. This adds information about the epifaunal-infaunal structure of the studied deposits and possible ecological interactions. In the modern environments, some starfish do not dig

randomly, but accurately locate and capture infaunal bivalves; they produce a pit often marked by furrows, which mirror the position of the arms (e.g., Smith, 1961; Doering, 1982). Although a possible predator-prey interaction is highly likely for trace fossils of asteroids and infaunal bivalves, it has rarely been reported (Quenstedt, 1858; Seilacher, 1953; Mángano et al., 1999). *Asteriacites* produced by asteroids have been usually linked to concealment and hiding activities (e.g., Häntzschel, 1970; Seilacher, 2007). Deeper burrows attributed to asterozoan activities are usually considered typical of dwelling, escaping or hunting purposes (Knaust and Neumann, 2016). Nevertheless, surface trails and shallow burrows produced by modern sea stars have been attributed to prey detection (Martin, 2013).

The latest neoichnological experiments provided additional insights into the morphology of shallow asteroid burrows and their sculpture (e.g., Ishida et al., 2017). Future neoichnological studies with asteroids and prey such as bivalves might help clarify the resultant morphology of the shallow burrows while the tracemaker is involved in prey detection and/or hunting, short-term vs. long-term burial, etc.

7. Fossil record of asteroids in South America

Fossil asteroids are scarce in South America. Body fossils are known from the Lower Devonian of Argentina and Brazil (Melo, 1988; Haude, 1995), the Upper Cretaceous of Brazil (Maury, 1930; de Castro Manso, 2006), and from Lower Cretaceous deposits in Argentina (Fernández et al., 2014). In Chile, Lower Cretaceous *Asteriacites* have been assigned to *A. lumbricalis* and ascribed to asteroids (Bell, 2004). So far it is the only South American record of trace fossils with asteroids as putative producers. Based on the figured material and the latest revision of the ichnogenus by Knaust and Neumann (2016), the specimens described by Bell (2004) should be classified as *A. stelliformis*. The arm length:width ratio ranges between 4.4 and 5.5 and hence, it is significantly larger than the established threshold value of 2. Furthermore, there is “no evidence of a central disc” (Bell, 2004, p. 56). Regarding the assignment of producers, when present the striae appear to be continuous from side to side, as typical of ophiuroid trace fossils. Therefore, this material needs to be revised, but it is currently unavailable (pers. comm. Servicio Nacional de Geología y Minería, Chile).

8. Conclusions

Stellate trace fossils are described in detail for the first time from the Lower Cretaceous Agrio Formation (Neuquén Basin, Patagonia). One specimen is assigned to *Asteriacites stelliformis*, and the other tentatively to *A. stelliformis* or *A. quinquefolius*, because three arms match the morphometric characteristics of the former ichnospecies and two arms those of the latter. However, the sculpture resembles that of *A. quinquefolius*. The sculpture and morphometric data imply asteroids as producers of these *Asteriacites*.

The clear definition of the boundary in asterozoan trace fossils, necessary to apply the morphometric parameters used in their ichnotaxonomic assignment can be improved in negative epirelief specimens through simple photogrammetric procedures.

The *Asteriacites* represent the second record of asteroids in the Neuquén Basin, the first evidence of this group in the Agrio Formation, and one of the very few finds in South America at all. Until now, Astropectinidae is the only family of asteroids known from the Neuquén Basin and individuals belonging to it are considered as possible producers. The only astropectinid species described in the basin and the studied *Asteriacites* have a similar arm length:width ratio supporting the attribution to astropectinids. These animals are active predators capable of shallow burrowing. The close association of *Asteriacites* with bivalve traces assigned to Siphonichnidae suggests a predator-prey interaction between asteroids and bivalves in these deposits, adding information about the community structure.

The studied interval represents shallow-marine to marginal-marine deposits in a proximal mid-ramp and inner ramp setting. Interference and flat-topped ripples and wrinkle structures point to very shallow water depth even implying temporary subaerial exposure. Therefore, the depositional environment for this part of the Agrio Formation is shallower than previously assumed.

Acknowledgements

This study was partially funded by PICT 189 and UBACyT grants to M.B. Aguirre-Urreta and UBACyT grant (2015–2017) to Pablo J. Pazos. Esteban Beckwith provided support and encouragement during fieldwork. The Bunge and Born Foundation gave a grant to Diana Elizabeth Fernández. M.B. Aguirre-Urreta, D. Pérez, L. Luci and Y. Ishida shared literature. Andreas Wetzel was supported by Swiss National Foundation (grant 200021_169042/1). All these contributions are gratefully acknowledged. This work has been greatly improved by constructive comments and suggestions from Dirk Knaust, one anonymous reviewer and Marcin Machalski (Associate Editor). Eduardo Koutsoukos is thanked for his work as Editor-in-Chief. This is contribution R-259 of the Instituto de Estudios Andinos don Pablo Groeber.

References

- Aguirre-Urreta, M.B., Rawson, P.F., 2012. Lower Cretaceous ammonites from the Neuquén Basin, Argentina: a new heteromorph fauna from the uppermost Agrio Formation. *Cretaceous Research* 35, 208–216. <https://doi.org/10.1016/j.cretres.2012.01.001>.
- Aguirre-Urreta, M.B., Mourgues, F.A., Rawson, P.F., Bulot, L.G., Jaillard, E., 2007. The Lower Cretaceous Chãnarillo and Neuquén Andean basins: ammonoid biostratigraphy and correlations. *Geological Journal* 42, 143–173. <https://doi.org/10.1144/GSL.SP.2005.252.01.04>.
- Aguirre-Urreta, M.B., Pazos, P.J., Lazo, D.G., Fanning, C.M., Litvak, V.D., 2008. First U–Pb SHRIMP age of the Hauterivian stage, Neuquén Basin, Argentina. *Journal of South American Earth Sciences* 26, 91–99. <https://doi.org/10.1016/j.jsames.2008.01.001>.
- Aguirre-Urreta, M.B., Lazo, D.G., Griffin, M., Vennari, V., Parras, A.M., Cataldo, C., Garberoglio, R., Luci, L., 2011. Megainvertebrados del Cretácico y su importancia bioestratigráfica. In: *Relatorio del XVIII Congreso Geológico Argentino*, pp. 465–488. Neuquén, Argentina.
- Aguirre-Urreta, M.B., Lescano, M., Schmitz, M.D., Tunik, M., Concheyro, A., Rawson, P.F., Ramos, V.A., 2015. Filling the gap: new precise Early Cretaceous radioisotopic ages from the Andes. *Geology Magazine* 152, 1–8. <https://doi.org/10.1017/S001675681400082X>.
- Allen, J.R.L., 1984. Wrinkle marks: an intertidal sedimentary structure due to aseismic soft-sediment loading. *Sedimentary Geology* 41, 75–95. [https://doi.org/10.1016/0037-0738\(84\)90003-4](https://doi.org/10.1016/0037-0738(84)90003-4).
- Ballent, S., Concheyro, A., Sagasti, G., 2006. Bioestratigrafía y paleoambiente de la Formación Agrio (Cretácico Inferior), en la Provincia de Mendoza, Cuenca Neuquina, Argentina. *Revista Geologica de Chile* 33, 47–79. <https://doi.org/10.4067/S0716-02082006000100003>.
- Bell, C.M., 2004. Asteroid and ophiuroids trace fossil from the lower cretaceous of Chile. *Paleontology* 47, 51–66. <https://doi.org/10.1111/j.0031-0239.2004.00347.x>.
- Blake, D.B., Aronson, R.B., 1998. Stellerioidea of the Late Eocene La Meseta Formation, Seymour Island, Antarctic Peninsula. *Journal of Paleontology* 72, 339–353. <https://doi.org/10.1017/S0022336000036325>.

- Blake, D.B., Reid, R., 1998. Some Albian (Cretaceous) asteroids (Echinodermata) from Texas and their paleobiological Implications. *Journal of Paleontology* 72, 512–532. <https://doi.org/10.1017/S002233600002429X>.
- Brongniart, A.T., 1823. Observations sur les Fucoïdes. *Mémoire de la Société d'Historie Naturelle de Paris* 1, 301–320.
- Burchette, T.P., Wright, V.P., 1992. Carbonate ramp depositional systems. *Sedimentary Geology* 79, 3–57.
- Byrne, M., O'Hara, T.D., 2017. *Australian Echinoderms: Biology, Ecology and Evolution*. CSIRO Publishing, Melbourne.
- de Castro Manso, C.L., 2006. Primeiro registro de Goniasteridae (Echinodermata, Asteroidea) no Cretáceo do Brasil: Turoniano da Bacia Potiguar. *Geociencias* 25, 255–260.
- Cataldo, C.S., 2013. A new Early Cretaceous nerineoid gastropod from Argentina and its palaeobiogeographic and palaeoecological implications. *Cretaceous Research* 40, 51–60.
- Cichowolski, M., 2003. The nautiloid genus *Cymatoceras* from the Cretaceous of the Neuquén and Austral basins, Argentina. *Cretaceous Research* 24, 375–390. [https://doi.org/10.1016/S0195-6671\(03\)00050-8](https://doi.org/10.1016/S0195-6671(03)00050-8).
- Cichowolski, M., Pazos, P.J., Tunik, M.A., Aguirre-Urreta, M.B., 2012. An exceptional storm accumulation of nautilids in the Lower Cretaceous of the Neuquén Basin, Argentina. *Lethaia* 45, 121–138. <https://doi.org/10.1111/j.1502-3931.2011.00271.x>.
- Crimes, T.P., Crossley, J.D., 1991. A diverse ichnofauna from Silurian flysch of the Aberystwyth Grits Formation, Wales. *Geological Journal* 26, 27–64. <https://doi.org/10.1002/gj.3350260104>.
- Doering, P.H., 1982. Reduction of sea star predation by the burrowing response of the hard clam *Mercenaria mercenaria* (Mollusca: Bivalvia). *Estuaries Research Federation* 5, 310–315. <https://doi.org/10.2307/1351754>.
- Dumas, S., Arnott, R.W.C., 2006. Origin of hummocky and swaley cross stratification: the controlling influence of unidirectional current strength and aggradation rate. *Geology* 34, 1073–1076. <https://doi.org/10.1130/G22930A.1>.
- Fernández, D.E., 2013. *Ichнологía de facies marino-marginales en el Cretácico Inferior de la cuenca Neuquina*. PhD Thesis. Universidad de Buenos Aires, 440 p.
- Fernández, D.E., Pazos, P.J., 2012. Ichnology of marginal marine facies of the Agrio Formation (Lower Cretaceous, Neuquén Basin, Argentina) at its type locality. *Ameghiniana* 49, 505–524.
- Fernández, D.E., Pazos, P.J., 2013. Xiphosurid trackways in a Lower Cretaceous tidal flat in Patagonia: palaeoecological implications and the involvement of microbial mats in trace-fossil preservation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 375, 16–29. <https://doi.org/10.1016/j.palaeo.2013.02.008>.
- Fernández, D.E., Pérez, D.E., Luci, L., Carrizo, M.A., 2014. An Early Cretaceous astropectinid (Echinodermata, Asteroidea) from Patagonia (Argentina): a new species and the oldest record of the family for the Southern Hemisphere. *Andean Geology* 41, 210–223. <https://doi.org/10.5027/andgeoV41n1-a08>.
- Fernández, D.E., Comerio, M., Pazos, P.J., 2018. *Nereites* in Lower Cretaceous marginal-marine facies from Patagonia: ichnotaxonomic and ethological implications. *Cretaceous Research* 81, 51–63. <https://doi.org/10.1016/j.cretres.2017.09.002>.
- Gingras, M.K., Pemberton, S.G., Smith, M., 2014. Bioturbation: reworking sediments for better and worse. *Oilfield Review Winter 2014/2015* 26 (4), 46–58.

- Guler, M.V., Lazo, D.G., Pazos, P.J., Borel, C.M., Ottone, E.G., Tyson, R.V., Cesaretti, N., Aguirre-Urreta, M.B., 2013. Palynofacies analysis and palynology of the Agua de la Mula Member (Agrio Formation) in a sequence stratigraphy framework, Lower Cretaceous, Neuquén Basin, Argentina. *Cretaceous Research* 41, 65–81.
- Hagadorn, J.W., Bottjer, D.J., 1997. Wrinkle structures: microbially mediated sedimentary structures common in subtidal siliciclastic settings at the Proterozoic-Phanerozoic transition. *Geology* 25, 1047–1050. [https://doi.org/10.1130/0091-613\(1997\)025<1047:WSMMS>2.3.CO;2](https://doi.org/10.1130/0091-613(1997)025<1047:WSMMS>2.3.CO;2)
- Häntzschel, W., 1970. Star-like trace fossils. In: Crimes, T.P., Harper, J.C. (Eds.), *Trace Fossils*. Geological Journal, Special Issue 3, 201–214.
- Haude, R., 1995. Echinodermen aus dem Unter-Devon der argentinischen Praekordillere. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 197, 37–86.
- Heddle, D., 1967. Versatility of movement and the origin of the asteroids. In: Millott, N. (Ed.), *Echinoderm Biology, Symposium of the Zoological Society of London* 20, 125–141.
- Heer, O., 1865. *Die Urwelt der Schweiz*. Friedrich Schultze, Zürich, 662 pp.
- Howell, J.A., Schwarz, E., Spalletti, L.A., Veiga, G.D., 2005. The Neuquén Basin: An overview. In: Veiga, G.D., Spalletti, L.A., Howell, J.A., Schwarz, E. (Eds.), *The Neuquén Basin, Argentina: A case study in sequence stratigraphy and basin dynamics*. Geological Society of London, Special Publications 252, 1–14.
- Ishida, Y., Röper, M., Fujita, T., 2013. Comparison of asteroids and ophiuroids trace fossils in the Jurassic of Germany with resting behavior of extant asteroids and ophiuroids. In: Johnson, C. (Ed.), *13th International Echinoderms Conferences: Echinoderms in a Changing World*, pp. 87–94.
- Ishida, Y., Fujita, T., Kiyomoto, M., Röper, M., Komatsu, T., Kato, K., Kamada, K., Shigeta, Y., Kumagae, T., 2017. How striations of ophiuroid and asteroid trace fossils were produced – Observations of tube-feet movement in living ophiuroids and asteroids. *Palaentological Society of Japan, Paleontological Research* 21, 27–36. <https://doi.org/10.2517/2016PR003>.
- Jagt, J.W.M., 2000. Late Cretaceous-Early Palaeogene echinoderms and the K/T boundary in the southeast Netherlands and northeast Belgium – Part 5: Asteroids. *Scripta Geologica* 121, 377–503.
- Knaust, D., 2015. Siphonichnidae (new ichnofamily) attributed to the burrowing activity of bivalves: ichnotaxonomy, behaviour and palaeoenvironmental implications. *Earth Science Reviews* 150, 497–519. <https://doi.org/10.1016/j.earscirev.2015.07.014>.
- Knaust, D., Neumann, C., 2016. *Asteriacites* von Schlotheim, 1820 – the oldest valid ichnogenus name – and other asterozoan produced trace fossils. *Earth-Science Reviews* 157, 111–120. <https://doi.org/10.1016/j.earscirev.2016.04.003>.
- Lazo, D.G., Cichowolski, M., Rodríguez, D.L., Aguirre-Urreta, M.B., 2005. Lithofacies, palaeoecology and palaeoenvironments of the Agrio Formation, Lower Cretaceous of the Neuquén Basin, Argentina. In: Veiga, G.D., Spalletti, L.A., Howell, J.A., Schwarz, E. (Eds.), *The Neuquén Basin, Argentina: A Case Study in Sequence Stratigraphy and Basin Dynamics*. Geological Society of London, Special Publications 252, 295–315.
- Lazo, D.G., Concheyro, G.A., Ottone, E.G., Guler, M.V., Aguirre-Urreta, M.B., 2009. Bioestratigrafía integrada de la Formación Agrio en su localidad tipo, Cretácico temprano de cuenca. *Neuquina. Revista de la Asociación Geológica Argentina* 65, 322–341.

- Leanza, H.A., Hugo, C.A., Repol, D., 2001. Hoja geológica 3969-I, Zapala, provincia del Neuquén, vol. 275. Instituto de Geología y Recursos Minerales. Servicio Geológico Minero Argentino, Boletín, Buenos Aires, 128 p.
- Legarreta, L., Uliana, M.A., 1999. El Jurásico y Cretácico de la Cordillera Principal y la Cuenca Neuquina. I, Facies sedimentarias. In: Caminos, R. (Ed.), Geología Argentina. SEGEMAR, Buenos Aires, pp. 399–416.
- Liu, X., Jia, Y., Zheng, J., Wen, M., Shan, H., 2017. An experimental investigation of wave-induced sediment responses in a natural silty seabed: new insights into seabed stratification. *Sedimentology* 64, 508–529. <https://doi.org/10.1111/sed.12312>.
- Luci, L., Garberoglio, R.M., Lazo, D.G., 2013. Serpulids and other calcareous tube-dwelling encrusting polychaetes from the Early Cretaceous Agrio Formation (Neuquén Basin, Argentina). *Geobios* 46, 213–224. <https://doi.org/10.1016/j.geobios.2012.06.003>.
- Mallison, H., Wings, O., 2014. Photogrammetry in paleontology e a practical guide. *Journal of Paleontological Techniques* 12, 1–31.
- Mángano, M.G., Buatois, L.A., West, R.R., Maples, C.G., 1999. The origin and paleoecologic significance of the trace fossil *Asteriacites* in the Pennsylvanian of Kansas and Missouri. *Lethaia* 32, 17–30.
- Martin, A.J., 2013. *Life Traces of the Georgia Coast: Revealing the Unseen Lives of Plants and Animals*. Indiana University Press, Bloomington and Indianapolis, Indiana. 670 p.
- Martin-Medrano, L., García-Barrera, P., 2013. Análisis del registro fósil de ofiuroideos (Echinodermata) en el Continente Americano. *Boletín de la Sociedad Geológica Mexicana* 65, 497–509.
- Martínez, S., del Río, C., 2015. Fossil Asterozoa (Echinodermata) of Argentina. *Revista de Biología Tropical* 63, 1–6. <https://doi.org/10.15517/rbt.v63i2.23123>.
- Mata, S.A., Bottjer, D.J., 2009. The paleoenvironmental distribution of Phanerozoic wrinkle structures. *Earth Science Reviews* 96, 181–195. <https://doi.org/10.1016/j.earscirev.2009.06.001>.
- Maury, C.J., 1930. O Cretaceo da Parahyba do Norte. Serviço Geologico e Mineralogico do Brasil Monographia 8, 305 pp.
- McIlroy, D., Flint, S., Howell, J.A., Timms, N., 2005. Sedimentology of the tide-dominated Jurassic Lajas Formation, Neuquén Basin, Argentina. In: Veiga, G.D., Spalletti, L.A., Howell, J.A., Schwarz, E. (Eds.), *The Neuquén Basin, Argentina: A case Study in Sequence Stratigraphy and Basin Dynamics*. Geological Society of London, Special Publications 252, 83–107.
- Melo, J.H.G., 1988. The Malvinokaffric realm in the Devonian of Brazil. In: McMillan, N.J., Embry, A.F., Glass, D.J. (Eds.), *Devonian of the World*, vol. 1. Canadian Society of Petroleum Geologists, pp. 669–703.
- Meunier, S., 1886. Sur quelques empreintes problématiques des couches boloniennes du Pas-de-Calais. *Bulletin de la Société Géologique de France, Série 3*, 14, 564–568.
- Miller, S.A., Dyer, C.B., 1878. Contributions to paleontology, descriptions of Cincinnati and Niagaran fossils. *Cincinnati Society of Natural History Journal* 1, 24–39.
- Ortega, I., Martín, A., Díaz, Y., 2011. Distribución, parámetros poblacionales y dieta de *Astropecten marginatus* (Asteroidea: Astropectinidae) en el Atlántico venezolano. *Revista de Biología Tropical* 59, 53–69.

- Pazos, P.J., Fernández, D.E., 2010. Three-dimensionally integrated trace-fossils from shallow-marine deposits in the Lower Cretaceous of the Neuquén Basin (Argentina): *Hillichnus agrioensis* isp. nov. *Acta Geologica Polonica* 60, 105–118.
- Pazos, P.J., Lazo, D.G., Tunik, M.A., Marsicano, C.A., Fernández, D.E., Aguirre-Urreta, M.B., 2012. Paleoenvironmental framework of dinosaur track-sites and other ichnofossils in Early Cretaceous mixed siliciclastic-carbonate deposits in the Neuquén Basin, northern Patagonia (Argentina). *Gondwana Research* 22, 1125–1140.
- Quenstedt, F.A., 1858. *Der Jura*. Verlag der H. Laupp'schen Buchhandlung, Tübingen, 842 pp., 100 pl.
- Quenstedt, F.A., 1876. *Petrefaktenkunde Deutschlands*. 4. Die Asteriden und Encriniden nebst Cysti und Blastoideen. Fues, Leipzig, 742 pp.
- Reineck, H.E., Singh, I.B., 1980. *Depositional sedimentary environments*. Springer, Berlin. 549 pp.
- Rodríguez, D.L., 2007. Equinoideos mesozoicos de las cuencas andinas del centrooeste de Argentina. PhD Thesis. Universidad de Buenos Aires, 398 pp.
- Rodríguez, D.L., Pazos, P.J., Aguirre-Urreta, M.B., 2007. Cretaceous ophiuroid trace fossils from the Neuquén Basin, Argentina. In: Bromley, R.G., Buatois, L.A., Mángano, G., Genise, J.F., Melchor, R.N. (Eds.), *Sediment-Organism Interactions: A Multifaceted Ichnology*. Society for Sedimentary Geology, Special Publication 88, 97–105.
- Salter, J.W., 1857. On annelide-burrows and surface-markings from the Cambrian rocks of the Longmynd, No. 2. *Quarterly Journal of the Geological Society of London* 13, 19–206.
- Schlotheim, E.F. von, 1820. *Die Petrefaktenkunde auf ihrem jetzigen Standpunkte durch die Beschreibung seiner Sammlung versteinertes und fossiler Überreste des Thier- und Pflanzenreichs der Vorwelt erläutert*. Becker, Gotha, 437 p.
- Seilacher, A., 1953. *Studien zur Palichnologie II. Die fossilen Ruhespuren (Cubichnia)*. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 98, 87–124.
- Seilacher, A., 2007. *Trace Fossil Analysis*. Springer-Verlag, Berlin Heidelberg, p. 226.
- Singh, B.P., Bhargava, O.N., Mikuláš, R., Prasad, S.K., Singla, G., Kaur, R., 2017. *Asteriacites* and other trace fossils from the Po Formation (Visean–Serpukhovian), Ganmachidam Hill, Spiti Valley (Himalaya) and its paleoenvironmental significance. *Geologica Carpathica* 68, 464–478. <https://doi.org/10.1515/geoca-2017-0030>.
- Smith, L.S., 1961. Clam-digging behavior in the starfish *Pisaster brevispinus* (Stimpson 1857). *Behaviour* 18 (1/2), 148–153.
- Spalletti, L.A., Poiré, D.G., Schwarz, E., Veiga, G.D., 2001a. Sedimentologic and sequence stratigraphic model of a Neocomian marine carbonate-siliciclastic ramp: Neuquén basin, Argentina. *Journal of South American Earth Sciences* 14, 609–624.
- Spalletti, L.A., Poire, D., Pirrie, D., Matheos, S., Doyle, P., 2001b. Respuesta sedimentológica a cambios en el nivel de base en una secuencia mixta clástica carbonática del Cretácico de la cuenca Neuquina, Argentina. *Revista de la Sociedad Geologica de Espana* 14, 57–74.
- Stanistreet, I.G., le Blanc Smith, G., Cadle, A.B., 1980. Trace fossils as sedimentological and palaeoenvironmental indices in the Ecca Group (Lower Permian) of the Transvaal. *Transactions of the Geological Society of South Africa* 83, 333–344.
- Taylor, P.D., Lazo, D.G., Aguirre-Urreta, M.B., 2009. Lower Cretaceous bryozoans from Argentina: a ‘by-catch’ fauna from the Agrio Formation (Neuquén Basin). *Cretaceous Research* 30, 193–203. <https://doi.org/10.1016/j.cretres.2008.07.003>.

- Vergani, G.D., Tankard, A.J., Belotti, J., Welsink, H.J., 1995. Tectonic evolution and paleogeography of the Neuquén Basin, Argentina. In: Tankard, A.J., Suárez, R., Welsink, H.J. (Eds.), *Petroleum Basins of South America*. American Association of Petroleum Geologists, Memoir 62, 383–402.
- Villier, L., Kutscher, M., Mah, C.L., 2004. Systematics and palaeoecology of middle Toarcian Asteroidea (Echinodermata) from the ‘Seuil du Poitou’, Western France. *Geobios* 37, 807–825. <https://doi.org/10.1016/j.geobios.2003.05.010>.
- Weaver, C.E., 1931. Paleontology of the Jurassic and Cretaceous of West Central Argentina, vol. 1. Memoir of the University of Washington, pp. 1–469.
- Zapata, T., Folguera, A., 2005. Tectonic evolution of the Andean Fold and thrust belt of the southern Neuquén Basin, Argentina. In: Veiga, G.D., Spalletti, L.A., Howell, J.A., Schwarz, E. (Eds.), *The Neuquén Basin, Argentina: A Case Study in Sequence Stratigraphy and Basin Dynamics*. Geological Society of London, Special Publications 252, 37–56. <https://doi.org/10.1144/GSL.SP.2005.252.01.03>.
- Zenker, J.C., 1836. *Historisch-topographisches Taschenbuch von Jena und seiner Umgebung besonders in naturwissenschaftlicher und medicinischer Beziehung*. Frommann, Jena, 338 pp.