

**Remarks on the degree
growth of
birational transformations**

C. Urech

Departement Mathematik und Informatik
Fachbereich Mathematik
Universität Basel
CH-4051 Basel

Preprint No. 2016-19
September 2016

www.math.unibas.ch

REMARKS ON THE DEGREE GROWTH OF BIRATIONAL TRANSFORMATIONS

CHRISTIAN URECH

ABSTRACT. We look at sequences of positive integers that can be realized as degree sequences of iterates of rational dominant maps of smooth projective varieties over arbitrary fields. New constraints on the degree growth of endomorphisms of the affine space and examples of degree sequences are displayed. We also show that the set of all degree sequences of rational maps is countable; this generalizes a result of Bonifant and Fornaess.

CONTENTS

1. Introduction and results	1
2. Preliminaries	5
3. Proofs	9
4. Remarks	11
References	11

1. INTRODUCTION AND RESULTS

1.1. Groups of birational transformations and degree sequences. Let X_k be a projective variety defined over a field k ; denote by $\text{Bir}(X_k)$ the group of birational transformations of X_k . A group Γ is called a *group of birational transformations* if there exists a field k and a projective variety X_k over k such that $\Gamma \subset \text{Bir}(X_k)$. More generally, one can consider $\text{Rat}(X_k)$, the monoid of dominant rational maps of X_k . Accordingly, we call a monoid Δ a *monoid of rational dominant transformations*, if there exists a field k and a projective variety X_k over k such that $\Delta \subset \text{Bir}(X_k)$.

If X_k is a smooth projective variety, an interesting tool to study the structure of monoids of rational dominant transformations are *degree functions*. Fix a polarization of X_k , i.e. an ample divisor class H of X_k . Then one can associate to every element $f \in \text{Rat}(X_k)$ its degree $\deg_H(f) \in \mathbb{Z}^+$ with respect to H , which is defined by

$$\deg_H(f) = f^* H \cdot H^{d-1},$$

Date: July 2016.

2010 *Mathematics Subject Classification.* 14E07; 37P05; 32H50.

The author gratefully acknowledges support by the Swiss National Science Foundation Grant "Birational Geometry" PP00P2 128422 /1 as well as by the Geldner-Stiftung, the FAG Basel, the Janggen Pöhn-Stiftung and the State Secretariat for Education, Research and Innovation of Switzerland.

where d is the dimension of X_k and f^*H is the total transform of H under f . For a smooth projective variety X_k over a field of characteristic zero k , one has for $f, g \in \text{Bir}(X_k)$

$$\deg_H(f \circ g) \leq C(X_k, H) \deg_H(f) \deg_H(g),$$

where $C(X_k, H)$ is a constant only depending on X_k and the choice of polarization H (see [DS05]). For a generalization of this result to fields of positive characteristic, see [Tru15].

Let f be a rational self map of \mathbb{P}_k^d . With respect to homogeneous coordinates $[x_0 : \dots : x_d]$ of \mathbb{P}_k^d , f is given by $[x_0 : x_1 : \dots : x_d] \mapsto [f_0 : f_1 : \dots : f_d]$, where $f_0, \dots, f_d \in k[x_0, \dots, x_d]$ are homogeneous polynomials of the same degree and without a common factor. We define $\deg(f) = \deg(f_i)$. Note that if f is dominant, then $\deg(f) = \deg_H(f)$ for $H = \mathcal{O}(1)$. So in case $X_k = \mathbb{P}_k^d$ we can extend the notion of degree to all rational self maps. Note that if f is an endomorphism of \mathbb{A}_k^d defined by $(x_1, \dots, x_d) \mapsto (f_1, \dots, f_d)$ with respect to affine coordinates (x_1, \dots, x_d) of \mathbb{A}_k^d , where $f_1, \dots, f_d \in k[x_1, \dots, x_d]$, then $\deg(f)$ is the maximal degree of the f_i .

Let $\Delta \subset \text{Bir}(X_k)$ be a finitely generated monoid of rational dominant transformations of a smooth projective variety X_k with a finite set of generators S . We define

$$D_{S,H}: \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$$

by

$$D_{S,H}(n) := \max_{\delta \in B_S(n)} \{\deg_H(\gamma)\},$$

where, $B_S(n)$ denotes all elements in Δ of word length $\leq n$ with respect to the generating set S . We call a map $\mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ that can be realized for some field k and some (X_k, H, Δ, S) as such a function a *degree sequence*.

Note that our definition of degree sequences includes in particular degree sequences that are given by finitely generated groups of birational transformations $\Gamma \subset \text{Bir}(X_k)$.

In this paper we show that there exist only countably many degree sequences, display certain constraints on their growth and give some new examples.

1.2. Countability of degree sequences. In [BF00], Bonifant and Fornaess proved that the set of sequences $\{d_n\}$ such that there exists a rational self map f of $\mathbb{P}_\mathbb{C}^d$ satisfying $\deg(f^n) = d_n$ for all n , is countable, which answered a question of Ghys. We generalize the result of Bonifant and Fornaess to all degree sequences over all smooth projective varieties, all fields, all polarizations and all finite generating sets S of finitely generated monoids of rational dominant maps:

Theorem 1.1. *The set of all degree sequences is countable.*

1.3. Previous results. In dimension 2 the degree growth of birational transformations is well understood and is a helpful tool to understand the group structure of $\text{Bir}(S_k)$ for projective surfaces S_k over a field k .

Theorem 1.2 (Gizatullin; Cantat; Diller and Favre). *Let k be an algebraically closed field, S_k a projective surface over k with a fixed polarization H and $f \in \text{Bir}(S_k)$. Then one of the following is true:*

- the set $\{\deg_H(f^n)\}$ is bounded;
- $\deg_H(f^n) \sim cn$ for some positive constant c and f preserves a rational fibration;

- $\deg_H(f^n) \sim cn^2$ for some positive constant c and f preserves an elliptic fibration;
- $\deg_H(f^n) \sim \lambda^n$, where λ is a Pisot or Salem number.

For more details on this rich subject and references to the proof of Theorem 1.2 see [Can12a].

In the case of polynomial automorphisms of the affine plane, the situation is even less complicated. Let $f \in \text{Aut}(\mathbb{A}_k^2)$. Then the sequence $\{\deg(f^n)\}$ is either bounded or it grows exponentially in n . See [Fur99] for this and more results on the degree growth in $\text{Aut}(\mathbb{A}_k^2)$.

In higher dimensions there are only few results on the degree growth of birational transformations. In particular, the following questions are open:

Question 1. Does there exist a birational transformation f of a projective variety X_k such that $\deg_H(f^n)$ is of intermediate growth, for instance $\deg_H(f^n) \sim e^{\sqrt{n}}$?

Question 2. Does there exist a birational transformation f such that $\deg_H(f^n)$ is unbounded, but grows "slowly"? For instance, can we have $\deg_H(f^n) \sim \sqrt{n}$? Or do unbounded degree sequences grow at least linearly?

Question 3. If there is a birational transformation f such that $\deg_H(f^n) \sim \lambda^n$, is λ always an algebraic number?

Question 4. Do birational transformations of polynomial growth always preserve some non-trivial rational fibration?

In [LB14] Lo Bianco treats the case of automorphisms of compact Kähler three-folds.

1.4. Degree sequences of polynomial automorphisms. A good place to start the examination of degree sequences seems to be the group of polynomial automorphisms of the affine d -space $\text{Aut}(\mathbb{A}_k^d)$. In Section 3.2 we will show the following observation (the proof of which can be found as well in [Dés16]):

Proposition 1.3. *Let k be a field and $f \in \text{Aut}(\mathbb{A}_k^d)$ a polynomial automorphism such that $\deg(f^d) = \deg(f)^d$, then $\deg(f^n) = \deg(f)^n$ for all $n \in \mathbb{Z}^+$.*

The monoid $\text{End}(\mathbb{A}_k^d)$ has the additional structure of a k -vector space, on which the degree function induces a filtration of finite dimensional vector spaces. This gives rise to a new technique, which we will use to prove that unbounded degree sequences of groups of polynomial automorphisms diverge and can not grow arbitrarily slowly:

Theorem 1.4. *Let $f \in \text{End}(\mathbb{A}_k^d)$ be an endomorphism such that the sequence $\{\deg(f^n)\}$ is unbounded. Then for all integers K*

$$\#\{m \mid \deg(f^m) \leq K\} < C_d \cdot K^d,$$

where $C_d = \frac{(1+d)^d}{(d-1)!}$. In particular, $\deg(f^n)$ diverges to ∞ .

By a result of Ol'shanskii ([Ols99]), Theorem 1.4 shows that an unbounded degree sequence of a polynomial automorphism behaves in some ways like a word length function.

The following corollary is immediate:

Corollary 1.5. *Let $\Gamma \subset \text{End}(\mathbb{A}_k^d)$ be a finitely generated monoid with generating system S . If $D_S(n) < C_d \cdot n^{1/d}$ for infinitely many n then Γ is of bounded degree.*

Unfortunately our methods to prove Theorem 1.4 do not work for arbitrary birational transformations of \mathbb{P}_k^d . However, if we assume the ground field to be finite, we obtain a similar result:

Theorem 1.6. *Let \mathbb{F}_q be a finite field with q elements and let $f \in \text{Rat}(\mathbb{P}_{\mathbb{F}_q}^d)$ such that the sequence $\{\deg(f^n)\}$ is unbounded. Then, for all integers K ,*

$$\#\{m \mid \deg(f^m) \leq K\} \leq q^{C(K,d)},$$

where $C(K,d) = (d+1) \cdot \binom{d+K}{K}$. In particular, $\deg(f^n)$ diverges to ∞ .

This implies the following for degree sequences in $\text{Rat}(\mathbb{P}_{\mathbb{F}_q}^d)$:

Corollary 1.7. *Let $\Gamma \subset \text{Rat}(\mathbb{P}_{\mathbb{F}_q}^d)$ be a finitely generated monoid with generating system S . There exists a positive constant $C_{d,q}$ such that if $D_S(n) < C_{d,q} \cdot \log(n)^{1/d}$ for all n , then Γ is of bounded degree.*

1.5. Types of degree growth.

Definition. Let X_k be a smooth projective variety with polarization H over a field k and let $f \in \text{Bir}(X_k)$. We denote the *order of growth* of $\deg_H(f^n)$ by

$$\text{dpol}(f) := \limsup_{n \rightarrow \infty} \frac{\log(\deg_H(f^n))}{\log(n)}.$$

The order of growth can be infinite.

By a result of Dinh and Sibony, the order of growth does not depend on the choice of polarization if we work over the field of complex numbers (see Section 2.4):

Proposition 1.8. *Let $X_{\mathbb{C}}$ be a smooth complex projective variety and let $f \in \text{Bir}(X_{\mathbb{C}})$. Then $\text{dpol}(f)$ does not depend on the choice of polarization.*

Let f be a birational transformation of a surface. As recalled above, in that case $\text{dpol}(f) = 0, 1, 2$ or ∞ . This gives rise to the following question:

Question 5. Does there exist a constant $C(d)$ depending only on d such that for all varieties X_k of dimension d we have $\text{dpol}(f) < C(d)$ for all $f \in \text{Bir}(X_k)$ with $\text{dpol}(f)$ finite?

We give some examples of degree sequences that indicate that the degree growth in higher dimensions is richer than in dimension 2.

First of all, note that elements in $\text{Aut}(\mathbb{A}_k^d)$ can have polynomial growth:

Example 1.9. Let k be any field and define $f, g, h \in \text{Aut}(\mathbb{A}_k^d)$ by $g = (x + yz, y, z)$, $h = (x, y + xz, z)$ and

$$f = g \circ h = (x + z(y + xz), y + xz, z).$$

One sees by induction that $\deg(f^n) = 2n + 1$; in particular, $\text{dpol}(f) = 1$.

Example 1.10. More generally, for all $l \leq d/2$ there exist elements $f_l \in \text{Aut}(\mathbb{A}_k^d)$ such that $\text{dpol}(f) = l$ (Section 3.4).

Other interesting examples of degree sequences of polynomial automorphisms and the dynamical behavior of the corresponding maps are described in [Dés16].

For birational transformations of \mathbb{P}_k^d we can obtain even faster growth (see [Lin12] for more details):

Example 1.11. The birational transformation $f = (x_1, x_1x_2, \dots, x_1x_2 \cdots x_n)$ of \mathbb{P}_k^d defined with respect to local affine coordinates (x_1, \dots, x_d) satisfies $\deg(f^n) = n^{d-1}$, i.e. $\text{dpol}(f) = d - 1$.

The following interesting observation is due to Serge Cantat:

Example 1.12. Define $\omega := \frac{-1+\sqrt{-3}}{2}$ and the elliptic curve $E_\omega := \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\omega)$. Let

$$X := E_\omega \times E_\omega \times E_\omega$$

and $s: X \rightarrow X$ the automorphism of finite order given by diagonal multiplication with $-\omega$. In [OT15] Oguiso and Truong prove that the quotient $Y := X/s$ is a rational threefold. Let $f: X \rightarrow X$ be the automorphism defined by $(x_1, x_2, x_3) \mapsto (x_1, x_1 + x_2, x_2 + x_3)$. Since f commutes with s , it induces an automorphism on Y , which we denote by \hat{f} . Let $\phi_1: \tilde{Y} \rightarrow Y$ be a resolution of the singularities of Y and define $\tilde{f} \in \text{Bir}(\tilde{Y})$ by

$$\tilde{f} := \phi_1^{-1} \circ \hat{f} \circ \phi_1.$$

We will show in Section 3.5 that $\text{dpol}(\tilde{f}) = 4$.

Remark 1. In [Dés16] Déserti shows that for all $l \leq d$ there exists a birational transformation $f \in \text{Bir}(\mathbb{P}_\mathbb{C}^d)$ such that $\text{dpol}(f) = l$.

1.6. Acknowledgements. I thank my advisors Jérémy Blanc and Serge Cantat for many interesting and helpful discussions and their constant support. I would also like to thank Junyi Xie for inspiring conversations and for giving me, together with Serge Cantat, some of the main ideas for the proof of Theorem 1.1, as well as Federico Lo Bianco for showing me Proposition 2.6. Many thanks also to Bac Dang and Julie Déserti for their comments and helpful references.

2. PRELIMINARIES

2.1. Monoids of rational dominant transformations. Let X_k be a projective variety over a field k . There is a one to one correspondence between rational dominant self maps of X_k and k -endomorphisms of the function field $k(X_k)$. The field $k(X_k)$ is the field of fractions of a k -algebra of finite type $k[x_1, \dots, x_n]/I$, where $I \subset k[x_1, \dots, x_n]$ is a prime ideal generated by elements $f_1, \dots, f_l \in k[x_1, \dots, x_n]$. A field extension $k \rightarrow k'$ induces a base change $X_{k'} \rightarrow X_k$. The function field of $X_{k'}$ is the field of fractions of the k' -algebra $k'[x_1, \dots, x_n]/I'$, where I' is the ideal generated by f_1, \dots, f_l . Note that $k(X_k) \subset k'(X_{k'})$. We say that a k' -endomorphism of $k'(X_{k'})$ is defined over k , if it restricts to a k -endomorphism of $k(X_k)$. Consider a k' -endomorphism G of $k'(X_{k'})$ sending generators (x_1, \dots, x_n) of $k'(X_{k'})$ to (g_1, \dots, g_n) , where $g_i \in k'(X_{k'})$. Then G is defined over k if and only if $g_i \in k(X_k)$ for all i . On the other hand, let $g_1, \dots, g_n \in k(X_k)$ and let $(x_1, \dots, x_d) \mapsto (g_1, \dots, g_d)$ be a k -endomorphism of $k(X_k)$. Then $(x_1, \dots, x_d) \mapsto (g_1, \dots, g_d)$ defines as well a k' -endomorphism of $k'(X_{k'})$. So a k -endomorphism $(x_1, \dots, x_n) \mapsto (g_1, \dots, g_n)$ of $k(X_k)$ extends uniquely to a k' -endomorphism of $k'(X_{k'})$. This yields the following observation:

Lemma 2.1. *Let X_k be a projective variety over a field k and $\varphi: k \rightarrow k'$ a homomorphism of fields. Then φ induces a natural injection of monoids $\Psi_\varphi: \text{Rat}(X_k) \rightarrow \text{Rat}(X_{k'})$.*

Recall that there are uncountably many finitely generated groups and thus in particular uncountably many finitely generated monoids. The following observation by de Cornulier shows that being a monoid of rational dominant transformations is in some sense a special property (cf. [Cor13] and [Can12b]).

Proposition 2.2. *There exist only countably many finitely generated isomorphism classes of monoids of rational dominant transformations.*

Proof. Let $\Delta \subset \text{Rat}(X_k)$ be a monoid of rational dominant transformations with a finite generating set $f_1, \dots, f_n \in \Delta$, where X_k is a projective variety defined over a field k . Denote by $S \subset k$ the finite set of coefficients necessary to define X_k and the rational dominant transformations f_i . Let k' be the field $\mathbb{F}_p(S)$, where $p = \text{char}(k)$, or $\mathbb{Q}(S)$ if $\text{char}(k) = 0$.

We consider the function field $k'(X_{k'})$ as a subfield of the function field $k(X_k)$. Note that the action of the elements of Δ on $k'(X_{k'})$ preserves $k'(X_{k'})$ and that $f_{i_1} f_{i_2} \cdots f_{i_{k-1}} = f_{i_k}$ in $\text{Rat}(X_k)$ if and only if $f_{i_1} f_{i_2} \cdots f_{i_{k-1}} = f_{i_k}$ in $\text{Rat}(X_{k'})$. So without loss of generality we may assume $\Delta \subset \text{Rat}(X_{k'})$, where k' is a finitely generated field extension of some \mathbb{F}_p or of \mathbb{Q} .

A rational dominant transformation of a given variety X_k is defined by finitely many coefficients in k . So the cardinality of the set of all finitely generated monoids of rational dominant transformations of a variety X_k is at most the cardinality of k .

Recall that the cardinality of the set of all finitely generated field extensions of \mathbb{F}_p and \mathbb{Q} is countable. Since a projective variety is defined by a finite set of coefficients, we obtain that there are only countably many isomorphism classes of projective varieties defined over a field k' that is a finitely generated field extension of \mathbb{F}_p or \mathbb{Q} . The claim follows. \square

2.2. Intersection form. Let X_k be a smooth projective variety of dimension d over an algebraically closed field k and let D be a Cartier divisor on X_k . The *Euler characteristic of D* is the integer

$$\chi(X_k, D) = \sum_{i=0}^{\infty} (-1)^i \dim_k H^i(X_k, D).$$

Let D_1, \dots, D_d be Cartier divisors on X_k . The function

$$(m_1, \dots, m_d) \mapsto \chi(X_k, m_1 D_1 + \cdots + m_d D_d)$$

is a polynomial in (m_1, \dots, m_d) . The *intersection number* of D_1, \dots, D_d is then defined to be the coefficient of the term $m_1 m_2 \cdots m_d$ in this polynomial and we denote it by $D_1 \cdot D_2 \cdots D_d$. One can show that intersection multiplicities are always integers and that the intersection form is symmetric and linear in all d arguments. Moreover, if D_1, \dots, D_d are effective and meet properly in a finite number of points, $D_1 \cdots D_d$ is the number of points in $D_1 \cap \cdots \cap D_d$ counted with multiplicities and the intersection form is the unique bilinear form with this property. For the details on this construction we refer to [Deb01] and the references in there. The intersection number is preserved under linear equivalence, therefore it is well defined on classes of Cartier divisors. Note as well that an isomorphism between algebraically closed

fields does not change the cohomology dimensions and hence that the intersection multiplicities are invariant under such an isomorphism.

Let X_k be a smooth projective variety of dimension d over an arbitrary field k and let D_1, \dots, D_d be Cartier divisors on X_k . Denote by \bar{k} an algebraic closure of k . We define the intersection multiplicity $D_1 \cdot D_2 \cdots D_d$ as the intersection multiplicity of D_1, D_2, \dots, D_d on $X_{\bar{k}}$ after base extension $k \rightarrow \bar{k}$. By the remark above, this does not depend on the choice of the algebraic closure \bar{k} . Every field isomorphism $k \rightarrow k'$ extends to an isomorphism between the algebraic closures of k and k' , hence the intersection number is invariant under field isomorphisms. Since the intersection form is unique, it also does not change under a base extension $k \rightarrow k'$ between algebraically closed fields.

We summarize these properties in the following proposition:

Proposition 2.3. *Let X_k be a smooth projective variety of dimension d over a field k . Then there exists a symmetric d -linear form on the group of divisors of X_k :*

$$\mathrm{Div}(X_k) \times \cdots \times \mathrm{Div}(X_k) \rightarrow \mathbb{Z}, (D_1, \dots, D_d) \mapsto D_1 \cdot D_2 \cdots D_d,$$

such that if D_1, \dots, D_d are effective and meet properly in a finite number of points, $D_1 \cdots D_d$ is the number of points in $D_1 \cap \cdots \cap D_d$ counted with multiplicity. Moreover, this intersection form is invariant under base change $k \rightarrow k'$ of fields.

2.3. Polarizations and degree functions. A polarization on a smooth projective variety X_k of dimension d is an ample divisor class H . This implies in particular that $Y \cdot H^{d-1} > 0$ for all effective divisors Y .

Let $f \in \mathrm{Rat}(X_k)$ and denote by $\mathrm{dom}(f)$ the maximal open subset of X_k on which f is defined. The graph Γ_f of f is the closure of $\{(x, f(x) \mid x \in \mathrm{dom}(f))\} \subset X_k \times X_k$. Let p_1 and $p_2: \Gamma_f \rightarrow X$ be the natural projections on the first respectively second factor, then $f = p_2 \circ p_1^{-1}$. Note that p_1 is birational. The total transform of a divisor D under f is the divisor $p_{1*}(p_2^*D)$, where p_2^*D is the pullback of D as a Cartier divisor and $p_{1*}(p_2^*D)$ is the pushforward of p_2^*D as a Weil divisor. Note that if k is algebraically closed and D is effective, then f^*D is the closure of all the points in $\mathrm{dom}(f)$ that are mapped to D by f .

2.4. Degrees in the case of complex varieties. Let $X_{\mathbb{C}}$ be a smooth complex projective variety of dimension d . The following constructions and results can be found in [Laz04]. Recall that two divisors D_1 and D_2 are numerically equivalent if $D_1 \cdot \gamma = D_2 \cdot \gamma$ for all curves γ on X . Denote by $N^1(X)$ the Neron-Severi group, which is the group of divisors modulo numerical equivalence. The intersection number of divisors D_1, \dots, D_d is invariant under numerical equivalence. By taking the first Chern class one can embed $N^1(X_{\mathbb{C}})$ into $H^2(X; \mathbb{Z})_{t.f.}$, which is $H^2(X_{\mathbb{C}}; \mathbb{Z})$ modulo its torsion part. It turns out that $N^1(X_{\mathbb{C}}) = H^2(X_{\mathbb{C}}; \mathbb{Z})_{t.f.} \cap H^{1,1}(X_{\mathbb{C}})$. Let D_1, \dots, D_d be divisors and $\omega_1, \dots, \omega_d$ the corresponding $(1, 1)$ -forms, then

$$D_1 \cdot D_2 \cdots D_d = \int_{X_{\mathbb{C}}} \omega_1 \wedge \omega_2 \wedge \cdots \wedge \omega_d.$$

A divisor is ample if and only if the corresponding $(1, 1)$ -form is cohomologous to a Kähler form. Hence a polarization of a smooth complex projective variety can be seen as a Kähler class. In this section we will take this point of view.

Let f be a birational transformation of X . With the help of currents, one can define the pullback $f^*\omega$ of a $(1, 1)$ -form ω (see [DN11]). If H is a divisor of X and ω the corresponding $(1, 1)$ -form then $f^*\omega$ is the form corresponding to f^*H .

If we fix a polarization $\omega_{X_{\mathbb{C}}}$ of smooth complex projective variety $X_{\mathbb{C}}$, then the degree of a birational map $f \in \text{Bir}(X_{\mathbb{C}})$ is defined by

$$\deg_{\omega_{X_{\mathbb{C}}}}(f) = \int_{X_{\mathbb{C}}} f^* \omega_{X_{\mathbb{C}}} \wedge \omega_{X_{\mathbb{C}}} \wedge \cdots \wedge \omega_{X_{\mathbb{C}}}.$$

So the degree does not depend on whether we look at an ample divisor class or its corresponding Kähler class.

In [DS05] the following is shown (see also [DN11]):

Proposition 2.4. *Let $X_{\mathbb{C}}$ be a smooth complex projective variety with polarization $\omega_{X_{\mathbb{C}}}$ and $f \in \text{Bir}(X_{\mathbb{C}})$. There exists a constant $A > 0$ such that*

$$A^{-1} \deg_{\omega_{X_{\mathbb{C}}}}(f^n) \leq \|(f^n)^*\| \leq A \deg_{\omega_{X_{\mathbb{C}}}}(f^n),$$

where $\|\cdot\|$ denotes any norm on $\text{End}(H^{1,1}(X_{\mathbb{C}}))$.

The following corollary follows directly from Proposition 2.4 and proves Proposition 1.8:

Corollary 2.5. *Let $X_{\mathbb{C}}$ be a smooth complex projective variety, then*

$$\text{dpol}(f) = \limsup_{n \rightarrow \infty} \frac{\log(\|(f^n)^*\|)}{\log(n)}.$$

In particular, $\text{dpol}(f)$ does not depend on the choice of polarization.

Proposition 2.6. *Let $X_{\mathbb{C}}$ and $Y_{\mathbb{C}}$ be complex smooth projective varieties of dimension d , let $f \in \text{Bir}(X_{\mathbb{C}})$, $g \in \text{Bir}(Y_{\mathbb{C}})$ and $\pi: X \rightarrow Y$ a dominant morphism of finite degree k , such that $\pi \circ f = g \circ \pi$. Then $\text{dpol}(f) = \text{dpol}(g)$.*

Proof. Let $\omega_{X_{\mathbb{C}}}$ be a Kähler form on $X_{\mathbb{C}}$ and $\omega_{Y_{\mathbb{C}}}$ be a Kähler form on $Y_{\mathbb{C}}$. We have

$$\begin{aligned} \deg_{\omega_{X_{\mathbb{C}}}}(f^n) &= \int_{X_{\mathbb{C}}} (f^n)^* \omega_{X_{\mathbb{C}}} \wedge \omega_{X_{\mathbb{C}}}^{d-1} = \deg(\pi) \int_{Y_{\mathbb{C}}} \pi_* (f^n)^* \omega_{X_{\mathbb{C}}} \wedge \pi_* \omega_{X_{\mathbb{C}}}^{d-1} \\ &= \deg(\pi) \int_{Y_{\mathbb{C}}} (g^n)^* \pi_* \omega_{X_{\mathbb{C}}} \wedge \pi_* \omega_{X_{\mathbb{C}}}^{d-1} \\ &\leq K_1 \deg(\pi) \cdot \|(g^n)^*\| \cdot \|\pi_* \omega_{X_{\mathbb{C}}}\| \cdot \|\pi_* \omega_{X_{\mathbb{C}}}^{d-1}\| \leq K_2 \cdot \|(g^n)^*\|, \end{aligned}$$

for some positive constants K_1, K_2 not depending on n . By Corollary 2.5, this yields $\text{dpol}(f) \leq \text{dpol}(g)$.

On the other hand, we have

$$\begin{aligned} \deg_{\omega_{Y_{\mathbb{C}}}}(g^n) &= \int_{Y_{\mathbb{C}}} (g^n)^* \omega_{Y_{\mathbb{C}}} \wedge \omega_{Y_{\mathbb{C}}}^{d-1} = \frac{1}{\deg(\pi)} \int_{X_{\mathbb{C}}} \pi^* (g^n)^* \omega_{Y_{\mathbb{C}}} \wedge \pi^* \omega_{Y_{\mathbb{C}}}^{d-1} \\ &= \frac{1}{\deg(\pi)} \int_{X_{\mathbb{C}}} (f^n)^* \pi^* \omega_{Y_{\mathbb{C}}} \wedge \pi^* \omega_{Y_{\mathbb{C}}}^{d-1} \\ &\leq K'_1 \frac{1}{\deg(\pi)} \|(f^n)^*\| \cdot \|\pi^* \omega_{Y_{\mathbb{C}}}\| \cdot \|\pi^* \omega_{Y_{\mathbb{C}}}^{d-1}\| \leq K'_2 \cdot \|(f^n)^*\|, \end{aligned}$$

for some positive constants K'_1, K'_2 not depending on n and therefore, by Corollary 2.5, $\text{dpol}(g) \leq \text{dpol}(f)$. □

3. PROOFS

3.1. Proof of Theorem 1.1. Let k be a field, X_k a smooth projective variety defined over k , H a polarization of X_k and $\Delta \subset \text{Rat}(X_k)$ a finitely generated monoid of rational dominant transformations with generating set T . Then X_k , H and T are defined by a finite set S of coefficients from k . Let $k' \subset k$ be the field $\mathbb{F}_p(S)$, where $p = \text{char}(k)$, or $k' = \mathbb{Q}(S)$ if $\text{char}(k) = 0$. By Proposition 2.3, the degree of elements in Γ considered as rational dominant transformations of X_k with respect to the polarization H is the same as the degree of elements in Δ considered as rational dominant transformations of $X_{k'}$. So without loss of generality, we may assume that Δ is a submonoid of $\text{Rat}(X_{k'})$, where k' is a finite field extension of \mathbb{F}_p or of \mathbb{Q} .

As in the proof of Proposition 2.2, we use the fact that there are only countably many isomorphism classes of such varieties. Polarizations and rational self maps are defined by a finite set of coefficients, so the cardinality of the set of all $(k', X_{k'}, H, T)$ up to isomorphism, where k' is a finitely generated field extension of \mathbb{F}_p or \mathbb{Q} , $X_{k'}$ a smooth projective variety over k' , H a polarization of $X_{k'}$ and a finite set of elements in $\text{Rat}(X_{k'})$ is countable. It follows in particular that the set of all degree sequences is countable.

3.2. Proof of Proposition 1.3. Let k be a field and $f \in \text{Bir}(\mathbb{P}_k^d)$ a birational map that is given by $[f_0 : \cdots : f_d]$, with respect to homogeneous coordinates $[x_0 : \cdots : x_d]$, where the f_i are homogenous polynomials of the same degree without common factors. There are two important closed subsets of \mathbb{P}_k^d associated to f , the indeterminacy locus $\text{Ind}(f)$, consisting of all the points where f is not defined and the exceptional divisor $\text{Exc}(f)$, the set of all the points where f is not a local isomorphism. If f is not an automorphism, the indeterminacy locus is a closed set of codimension ≥ 2 and the exceptional divisor a closed set of codimension 1. Note that $\text{Ind}(f)$ is exactly the set of points, where all the f_i vanish. Let $X \subset \mathbb{P}_k^d$ be an irreducible closed set that is not contained in $\text{Ind}(f)$. We denote by $f(X)$ the closure of $f(X \setminus \text{Ind}(f))$ and we say that f contracts X if $\dim(f(X)) < \dim(X)$.

The following lemma is well known (see for example [FS94]):

Lemma 3.1. *Let k be a field and $g, f \in \text{Bir}(\mathbb{P}_k^d)$. Then $\deg(f \circ g) \leq \deg(f) \deg(g)$ and $\deg(f \circ g) < \deg(f) \deg(g)$ if and only if g contracts a hypersurface to a subset of $\text{Ind}(f)$.*

Proof. Let $f = [f_0 : \cdots : f_d]$ and $g = [g_0 : \cdots : g_d]$. Then $\deg(f \circ g) < \deg(f) \deg(g)$ if and only if the polynomials $f_0(g_0, \dots, g_d), \dots, f_d(g_0, \dots, g_d)$ have a non constant common factor $h \in k[x_0, \dots, x_d]$. Let $M \subset \mathbb{P}_k^d$ be the hypersurface defined by $h = 0$. Then f is not defined at $g(M)$. This implies that the codimension of $g(M)$ is ≥ 2 and therefore that g contracts M to a subset of $\text{Ind}(f)$.

On the other hand, let M be an irreducible component of a hypersurface that is contracted by g to a subset $\text{Ind}(f)$. Assume that M is the zero set of an irreducible polynomial h . Since $g(M) \subset \text{Ind}(f)$, we obtain that $f_0(g_0, \dots, g_d), \dots, f_d(g_0, \dots, g_d)$ vanish all on M and therefore that h divides $f_0(g_0, \dots, g_d), \dots, f_d(g_0, \dots, g_d)$. This implies $\deg(f \circ g) < \deg(f) \deg(g)$. \square

In order to prove Proposition 1.3, we consider an element $f \in \text{Aut}(\mathbb{A}_k^d)$ as a birational transformation of \mathbb{P}_k^d whose exceptional divisor is the hyperplane at infinity $H := \mathbb{P}_k^d \setminus \mathbb{A}_k^d$ and whose indeterminacy points are contained in H .

Note that $\deg(f^d) = \deg(f)^d$ implies $\deg(f^l) = \deg(f)^l$ for $l = 1, \dots, d$. We look at f as an element of $\text{Bir}(\mathbb{P}_k^d)$. If f is an automorphism of \mathbb{P}_k^d , its degree is 1 and the claim follows directly. Otherwise, f contracts the hyperplane H . By Lemma 3.1, $\deg(f^{l+1}) = \deg(f^l) \deg(f)$ is equivalent to $f^l(H)$ not being contained in $\text{Ind}(f)$. In particular, if $\deg(f^l) = \deg(f)^l$ and $f^l(H) = f^{l+1}(H)$ for some l then $\deg(f^i) = \deg(f)^i$ for all $i \geq l$.

Let $1 \leq l \leq d$. By Lemma 3.1, $f^l(H)$ is not contained in $\text{Ind}(f)$. Observe that $f^l(H)$ is irreducible and $f^{l+1}(H) \subset f^l(H)$. This implies that $\dim(f^{l+1}(H)) \leq \dim(f^l(H))$ and $\dim(f^{l+1}(H)) = \dim(f^l(H))$ if and only if $f^{l+1}(H) = f^l(H)$. It follows that the chain $H \supset f(H) \supset f^2(H) \supset \dots$ becomes stationary within the first d iterations. In particular, $f^d(H) = f^{d+i}(H)$ for all i and therefore $\deg(f^i) = \deg(f)^i$ for all i .

3.3. Proof of Theorem 1.4 and Theorem 1.6. We start by proving Theorem 1.4. Let $f \in \text{End}(\mathbb{A}_k^d)$ be an endomorphism such that the sequence $\{\deg(f^n)\}$ is unbounded. Our first remark is that the elements of $\{f^n\}$ are linearly independent in the vector space of polynomial endomorphisms $\text{End}(\mathbb{A}_k^d)$. Indeed, if for any n we have

$$f^n = \sum_{l < n} c_l f^l,$$

for some $c_1, \dots, c_{n-1} \in k$. It follows by induction that $\deg(f^{n+i})$ is smaller or equal to $\max_{l < n} \deg(f^l)$ for all i .

Denote by $\text{End}(\mathbb{A}^d)_{\leq K}$ the k -vector space of polynomial endomorphisms of degree $\leq K$, which has dimension $d \cdot \binom{d+K}{K}$. One calculates

$$d \cdot \binom{d+K}{K} < \frac{1}{(d-1)!} (K+d)^d \leq \frac{(1+d/K)^d}{(d-1)!} K^d \leq C_d K^d,$$

where $C_d = \frac{(1+d)^d}{(d-1)!}$. Since the elements in $\{f^k\} \cap \text{End}(\mathbb{A}^d)_{\leq K}$ are linearly independent, the cardinality of $\{f^k\} \cap \text{End}(\mathbb{A}^d)_{\leq K}$ is at most $C_d K^d$.

For the proof of Theorem 1.6 note that in the case of finite fields, there are only finitely many birational transformations of a given degree. If $f^l = f$ for some $l > 1$, then $\{\deg(f^n)\}$ is bounded. There are $\binom{d+K}{K}$ monomials of degree $\leq K$. A birational transformation of degree $\leq K$ is given by $d+1$ polynomials of degree $\leq K$, so by $C(K, d) = (d+1) \binom{d+K}{K}$ coefficients from \mathbb{F}_q . Hence there are less than $q^{C(K, d)}$ birational transformations of degree $\leq K$. This proves the claim.

3.4. Proof of Example 1.10. Let d be an integer and $l = \lfloor d/2 \rfloor$. For $d = 3$ the automorphism $f_3 := (x+z(y+xz), y+xz, z)$ from Example 1.9 satisfies $\deg(f_3^n) \sim n$. Moreover, the first coordinate of f_3^n is the coordinate with highest degree. Assume now that $d \geq 5$ and that we are given an automorphism $f_{d-2} \in \text{Aut}(\mathbb{A}^{d-2})$ such that $\deg(f_{d-2}^n)$ grows like n^{l-1} and that the first coordinate of f_{d-2}^n is the entry with highest degree. Let

$$f_d := (x_1 + x_3(x_2 + x_1 x_3), x_2 + x_1 x_3, f_{d-2}(x_3, \dots, x_d)).$$

One sees by induction that the degree of f_d^n grows like n^l and that the first coordinate of f_d^n is the coordinate of f_{d-2}^n with highest degree.

3.5. Proof of Example 1.12. In this section we use the notation introduced in Example 1.12.

Let dz_1, dz_2, dz_3 be a basis of $H^{1,0}(X)$. Then $d\bar{z}_1, d\bar{z}_2, d\bar{z}_3$ is a basis of $H^{0,1}(X)$. The automorphism f^n induces an action on both $H^{1,0}(X)$ and $H^{0,1}(X)$ whose norm grows like n^2 . Since $H^{1,1}(X) = H^{1,0}(X) \otimes H^{0,1}(X)$, this implies that the norm of the induced action of f^n on $H^{1,1}(X)$ grows like n^4 . By Corollary 2.5, we obtain $\text{dpol}(f) = 4$.

Denote by $\pi: X \rightarrow Y = X/s$ the quotient map. Let \tilde{X} be a smooth projective variety, $\phi_2: \tilde{X} \rightarrow X$ a birational morphism and $\psi: \tilde{X} \rightarrow \tilde{Y}$ a dominant morphism such that the following diagram commutes:

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\psi} & \tilde{Y} \\ \downarrow \phi_2 & & \downarrow \phi_1 \\ X & \xrightarrow{\pi} & Y. \end{array}$$

Note that ψ is generically finite. By Proposition 2.6, we have $\text{dpol}(\phi_2^{-1} \circ f \circ \phi_2) = \text{dpol}(f) = 4$ and hence, again by Proposition 2.6, $\text{dpol}(\tilde{f}) = \text{dpol}(\psi \circ \phi_2^{-1} \circ f \circ \phi_2 \circ \psi^{-1}) = 4$, which proves the claim of Example 1.12.

4. REMARKS

4.1. Other degree functions. One can define more general degree functions. Let X_k be a smooth projective variety over a field k with polarization H and $1 \leq l \leq d-1$, then

$$\text{deg}_H^l(f) := (f^*H)^l \cdot H^{d-l}.$$

These degree functions play an important role in dynamics. In characteristic 0, we still have $\text{deg}_H^l(fg) \leq C \text{deg}_H^l(f) \text{deg}_H^l(g)$ for a constant C not depending on f and g (see [DS05] and [Tru15] for generalizations to fields of positive characteristic).

Our proof of Theorem 1.1 works as well if we replace the function deg_H by deg_H^l for any l . Let $\Gamma \subset \text{Bir}(X_k)$ be a finitely generated group of birational transformations with a finite symmetric set of generators S . We define $D_{S,H}^l: \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ by $D_{S,H}^l(n) := \max_{\gamma \in B_S(n)} \{\text{deg}_H^l(\gamma)\}$ and we call a map $\mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ that can be realized for some (X_k, H, Γ, S, l) as such a function a *general degree sequence*.

Theorem 4.1. *The set of all general degree sequences is countable.*

Proof. Analogous to the proof of Theorem 1.1. □

REFERENCES

- [BF00] Araceli M. Bonifant and John Erik Fornæss, *Growth of degree for iterates of rational maps in several variables*, Indiana Univ. Math. J. **49** (2000), no. 2, 751–778. MR 1793690 [1.2](#)
- [Can12a] Serge Cantat, *The cremona group in two variables*, Proceedings of the sixth European Congress of Math, 2012, pp. 211–225. [1.3](#)
- [Can12b] ———, *A remark on groups of birational transformations and the word problem (after yves de cornulier)*, <https://perso.univ-rennes1.fr/serge.cantat/Articles/wpic.pdf> (2012). [2.1](#)
- [Cor13] Yves Cornulier, *Sofic profile and computability of Cremona groups*, Michigan Math. J. **62** (2013), no. 4, 823–841. MR 3160544 [2.1](#)
- [Deb01] Olivier Debarre, *Higher-dimensional algebraic geometry*, Universitext, Springer-Verlag, New York, 2001. MR 1841091 [2.2](#)

- [Dés16] Julie Déserti, *Degree growth of polynomial automorphisms and birational maps: some examples*, <http://arxiv.org/abs/1602.04642> (2016). [1.4](#), [1.5](#), [1](#)
- [DN11] Tien-Cuong Dinh and Viêt-Anh Nguyễn, *Comparison of dynamical degrees for semi-conjugate meromorphic maps*, *Comment. Math. Helv.* **86** (2011), no. 4, 817–840. MR 2851870 [2.4](#)
- [DS05] Tien-Cuong Dinh and Nessim Sibony, *Une borne supérieure pour l'entropie topologique d'une application rationnelle*, *Ann. of Math. (2)* **161** (2005), no. 3, 1637–1644. MR 2180409 [1.1](#), [2.4](#), [4.1](#)
- [FS94] John Erik Fornæss and Nessim Sibony, *Complex dynamics in higher dimensions.*, Complex potential theory. Proceedings of the NATO Advanced Study Institute and Séminaire de mathématiques supérieures, Montréal, Canada, July 26 - August 6, 1993, Dordrecht: Kluwer Academic Publishers, 1994, pp. 131–186 (English). [3.2](#)
- [Fur99] Jean-Philippe Furter, *On the degree of iterates of automorphisms of the affine plane*, *Manuscripta Math.* **98** (1999), no. 2, 183–193. MR 1667603 [1.3](#)
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 48, Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series. MR 2095471 [2.4](#)
- [LB14] Federico Lo Bianco, *Bornes sur les degrés dynamiques d'automorphismes de variétés kählériennes de dimension 3*, *C. R. Math. Acad. Sci. Paris* **352** (2014), no. 6, 515–519. MR 3210135 [1.3](#)
- [Lin12] Jan-Li Lin, *Algebraic stability and degree growth of monomial maps*, *Math. Z.* **271** (2012), no. 1-2, 293–311. MR 2917145 [1.5](#)
- [Ols99] A. Yu. Olshanskii, *Distortion functions for subgroups*, *Geometric group theory down under (Canberra, 1996)*, de Gruyter, Berlin, 1999, pp. 281–291. MR 1714850 [1.4](#)
- [OT15] Keiji Oguiso and Tuyen Trung Truong, *Explicit examples of rational and Calabi-Yau threefolds with primitive automorphisms of positive entropy*, *J. Math. Sci. Univ. Tokyo* **22** (2015), no. 1, 361–385. MR 3329200 [1.12](#)
- [Tru15] Tuyen Trung Truong, *(relative) dynamical degrees of rational maps over an algebraic closed field*, <http://arxiv.org/abs/1501.01523> (2015). [1.1](#), [4.1](#)

MATHEMATISCHES INSTITUT, UNIVERSITÄT BASEL, 4051 BASEL, SWITZERLAND

IRMAR, UNIVERSITÉ DE RENNES 1, 35042 RENNES, FRANCE

E-mail address: christian.urech@unibas.ch

LATEST PREPRINTS

No.	Author: <i>Title</i>
2015-36	J. K. Canci <i>Good reduction for endomorphisms of the projective line in terms of the branch locus</i>
2015-37	J. K. Canci, L. Paladino <i>Preperiodic points for rational functions defined over a global field in terms of good reduction</i>
2015-38	J. K. Canci <i>Preperiodic points for rational functions defined over a rational function field of characteristic zero</i>
2015-39	J. K. Canci, S. Vishkautsan <i>Quadratic maps with a periodic critical point of period 2</i>
2015-40	H. Harbrecht, M. Peters <i>The second order perturbation approach for PDEs on random domains</i>
2015-41	H. Harbrecht, W. L. Wendland, N. Zorii <i>Minimal energy problems for strongly singular Riesz kernels</i>
2015-42	H. Harbrecht, M. Utzinger <i>On adaptive wavelet boundary element methods</i>
2016-01	S. Iula, G. Mancini <i>Extremal functions for singular Moser-Trudinger embeddings</i>
2016-02	J. K. Canci, L. Paladino <i>On preperiodic points of rational functions defined over $\mathbb{F}_p(t)$</i>
2016-03	H. Harbrecht, R. Schneider <i>A note on multilevel based error estimation</i>
2016-04	C. Urech <i>On homomorphisms between Cremona groups</i>
2016-05	A. P. Choudhury, H. Heck <i>Stability of the inverse boundary value problem for the biharmonic operator: Logarithmic estimates</i>
2016-06	M. Dambrine, I. Greff, H. Harbrecht, B. Puig <i>Numerical solution of the homogeneous Neumann boundary value problem on domains with a thin layer of random thickness</i>

LATEST PREPRINTS

No.	Author:	Title
2016-07	G. Alberti, G. Crippa, A. L. Mazzucato	<i>Exponential self-similar mixing by incompressible flows</i>
2016-08	M. Bainbridge, P. Habegger, M. Möller	<i>Teichmüller curves in genus three and just likely intersections in $G_m^n \times G_a^n$</i>
2016-09	Gabriel A. Dill	<i>Effective approximation and Diophantine applications</i>
2016-10	J. Blanc, S. Zimmermann	<i>Topological simplicity of the Cremona groups</i>
2016-11	I. Hedén, S. Zimmermann	<i>The decomposition group of a line in the plane</i>
2016-12	J. Ballani, D. Kressner, M. Peters	<i>Multilevel tensor approximation of PDEs with random data</i>
2016-13	M. J. Grote, M. Kray, U. Nahum	<i>Adaptive eigenspace method for inverse scattering problems in the frequency domain</i>
2016-14	H. Harbrecht, M. Peters, M. Schmidlin	<i>Uncertainty quantification for PDEs with anisotropic random diffusion</i>
2016-15	F. Da Lio, L. Martinazzi	<i>The nonlocal Liouville-type equation in R and conformal immersions of the disk with boundary singularities</i>
2016-16	A. Hyder	<i>Conformally Euclidean metrics on R^n with arbitrary total Q-curvature</i>
2016-17	G. Mancini, L. Martinazzi	<i>The Moser-Trudinger inequality and its extremals on a disk via energy estimates</i>
2016-18	R. N. Gantner, M. D. Peters	<i>Higher order quasi-Monte Carlo for Bayesian shape inversion</i>
2016-19	C. Urech	<i>Remarks on the degree growth of birational transformations</i>