



# Activation of Primary and Secondary Benzylic and Tertiary Alkyl (sp<sup>3</sup>)C-F Bonds Inside a Self-Assembled Molecular Container

Jesper M. Köster<sup>1</sup>, Daniel Häussinger<sup>1</sup> and Konrad Tiefenbacher<sup>1,2\*</sup>

<sup>1</sup> Department of Chemistry, University of Basel, Basel, Switzerland, <sup>2</sup> Department of Biosystems Science and Engineering, ETH Zürich, Basel, Switzerland

Alkyl fluorides are generally regarded as chemically inert. However, several literature examples describe the activation of alkyl (sp<sup>3</sup>)C-F bonds via strong Brønsted or Lewis acids under harsh conditions. We here report that catalytic amounts of the self-assembled resorcinarene capsule are able to activate alkyl (sp<sup>3</sup>)C-F bonds under mild conditions (40°C, no strong Brønsted or Lewis acid present). Kinetic measurements display a sigmoidal reaction progress after an initial induction period. Control experiments indicate that the presence of the supramolecular capsule is required for an efficient reaction acceleration.

**Keywords:** supramolecular catalysis, molecular capsules, supramolecular chemistry, acid catalysis, elimination

## OPEN ACCESS

### Edited by:

Pablo Ballester,  
Catalan Institution for Research and  
Advanced Studies, Spain

### Reviewed by:

Yevgen Karpichev,  
Tallinn University of Technology,  
Estonia  
Guzman Gil-Ramirez,  
University of Lincoln, United Kingdom  
Joost Reek,  
UvA Minds, Netherlands

### \*Correspondence:

Konrad Tiefenbacher  
konrad.tiefenbacher@unibas.ch;  
tkonrad@ethz.ch

### Specialty section:

This article was submitted to  
Supramolecular Chemistry,  
a section of the journal  
Frontiers in Chemistry

**Received:** 26 June 2018

**Accepted:** 07 December 2018

**Published:** 04 January 2019


### Citation:

Köster JM, Häussinger D and  
Tiefenbacher K (2019) Activation of  
Primary and Secondary Benzylic and  
Tertiary Alkyl (sp<sup>3</sup>)C-F Bonds Inside a  
Self-Assembled Molecular Container.  
Front. Chem. 6:639.  
doi: 10.3389/fchem.2018.00639

## INTRODUCTION

Catalysis inside closed binding pockets, much like inside enzymes, has been a challenging research area but in recent decades active supramolecular systems have been reported (Marchetti and Levine, 2011; Wiester et al., 2011; Raynal et al., 2014; Brown et al., 2015; Leenders et al., 2015; Zarra et al., 2015; Catti et al., 2016a,b; Cullen et al., 2016; Jans et al., 2016; Kuijpers et al., 2016; Levin et al., 2016; Otte, 2016; Wang et al., 2016, 2017; Bräuer et al., 2017; Ueda et al., 2017; Catti and Tiefenbacher, 2018; La Manna et al., 2018a). Drawing upon a multitude of structural classes such as coordination cages, metal-organic frameworks, and non-covalent assemblies, a whole series of reactions were successfully catalyzed. One of the catalytically active supramolecular containers is the hexameric resorcinarene capsule **I** (see **Figure 1B**), which self-assembles in non-polar solvents (e.g., chloroform or benzene) from six molecules of resorcinarene (**1**, **Figure 1A**) and eight water molecules (MacGillivray and Atwood, 1997; Avram and Cohen, 2002). It possesses an internal volume of ~1,400 Å<sup>3</sup> and is readily accessible in a one-step condensation from resorcinol and dodecanal. It was demonstrated to serve as a mild Brønsted acid (pK<sub>A</sub> ~ 5.5–6) and functions as a non-nucleophilic counterion due to the extensive delocalization of the negative charge (Zhang and Tiefenbacher, 2013).

The supramolecular capsule **I** was used as a catalyst in the hydrolysis of acetals (Zhang and Tiefenbacher, 2013), the tail-to-head cyclization of terpenes (Zhang and Tiefenbacher, 2015), the hydroalkoxylation of alkenes (Catti and Tiefenbacher, 2015), the cyclodehydration of alcohols (Catti et al., 2017), the carbonyl-olefin metathesis (Catti and Tiefenbacher, 2018), the hydration of isonitriles (Bianchini et al., 2013), [2+3]-cycloaddition reactions (Giust et al., 2015; La Manna et al., 2018a), the Meinwald rearrangement of epoxides (Caneva et al., 2016), the hydration of alkynes (La Sorella et al., 2016a), the oxidation of thioethers (La Sorella et al., 2016b), and Friedel-Crafts reactions (La Manna et al., 2018b).



**FIGURE 1 |** (A) C<sub>11</sub>-resorcinarene (1); (B) hexameric capsule I, C<sub>11</sub>-feet omitted; (C) competitive guest TBAB (2).

Recently, we investigated “inert” fluorinated acyclic terpenes as probes to quantify substrate encapsulation via <sup>19</sup>F-NMR. Surprisingly, the C-F bond of tertiary fluoride **10** (Table 1) was activated in the presence of catalytic amounts of capsule (10 mol%). This observation prompted us to investigate this reaction more closely and here we present the results.












Alkyl halides, known for their broad spectrum of reactivity (substitution, elimination, metalation), are a cornerstone of organic chemistry. In contrast to iodides, bromides, and chlorides alkyl fluorides display a general lack of reactivity under typical conditions for halide chemistry. Contrary to intuition, the exceptionally high electronegativity of fluorine does not facilitate nucleophilic attacks on the adjacent carbon but instead strengthens the (sp<sup>3</sup>)C-F bond (105.4 kcal/mol, C-H: 98.8 kcal/mol, C-Cl: 78.5 kcal/mol), due to substantial electrostatic attraction between the highly polarized bonding partners (O’Hagan, 2008). Strategies that work for alkyl chlorides, bromides, or iodides seem to fail to sufficiently activate fluorides for further reactions. Nevertheless, some successful approaches for C-F bond activation have been developed (Burdeniuc et al., 1997; Amii and Uneyama, 2009; Kuehnel et al., 2013; Shen et al., 2015). Exploiting fluorine’s high affinity toward boron, and aluminum, most examples cover activation by Lewis acids. Activation by Brønsted acids, known since 1949, when Le Fave found that heating trifluorotoluene in concentrated sulfuric acid facilitated the hydrolysis to benzoic acid (Fave, 1949), has received relatively little attention by contrast. Wang et al. recently published an improved take on this approach to activate trifluorotoluyll substrates by employing an excess of triflic acid

at ambient temperature (Wang and Hu, 2009). The benzyl acylium generated by this method were then converted to benzophenone derivatives by adding arene nucleophiles to promote a Friedel-Crafts acylation. The inert nature of the CF<sub>3</sub> group requires the use of superstoichiometric amounts of strong acids, which limits the functional group tolerance considerably.

Although the topic of organic fluorides as hydrogen bond acceptors is controversial, increasing evidence indicates a weak interaction (Champagne et al., 2015a). A mild method of activation for benzylic fluorides via hydrogen bonding was recently developed by Paquin et al. (Champagne et al., 2014, 2015b). Hexafluoroisopropanol (HFIP) serves as hydrogen bond donor to the fluorine atom, which destabilizes the C-F bond to then generate a benzylic cation, which can be intercepted by an arene nucleophile in a Friedel-Crafts reaction. HF is liberated after the conversion and replaces HFIP as the catalytically active species because of its higher hydrogen bond donating ability (Rosenberg, 2012; Champagne et al., 2014).

Activation of non-benzylic fluorides remains challenging under Brønsted acidic conditions. Under harsh Brønsted acidic conditions (HCl, 80–110°C) elimination of tertiary alkyl fluorides was reported (Figures 2A,B). Primary and secondary fluorides show no reactivity due to the poor stabilization of the respective intermediate carbocations (Wenckens et al., 1998; Bellezza et al., 2005). Interestingly, the enzyme pocket of isopenicillin N synthase facilitated the elimination of a tertiary fluoride under mild conditions (Figure 2C).

**TABLE 1** | Substrate overview with yields; product denominations: **a**, alcohol; **e**, cyclic ether; **i**, internal alkene; **t**, terminal alkene; all reactions run in filtered CDCl<sub>3</sub>.

| Entry | Substrate   | Products   | Background   |
|-------|---|--|--|
| 1     |    | <b>3i</b> 69% <sup>a</sup> (16 h)<br><b>3t</b> 13% <sup>a</sup>                                | 0 <sup>b</sup><br>0 <sup>c</sup>   |
| 2     |    | <b>4i</b> 45% <sup>a</sup> (16 h)<br><b>4a</b> 35% <sup>a</sup><br><b>4t</b> 16% <sup>a</sup>  | 0 <sup>b</sup><br>0 <sup>c</sup>   |
| 3     |    | <b>5i</b> 60% <sup>a</sup> (20 h)<br><b>5i</b> 29% <sup>a</sup><br><b>5t</b> 3% <sup>a</sup>   | <br><b>5i</b><br>0 <sup>b</sup><br>0 <sup>c</sup>                   |
| 4     |    | <b>6e</b> 81% <sup>a</sup> (21 h)<br><b>6i</b> 11% <sup>a</sup>                                | <br><b>6e</b><br>0 <sup>b</sup><br>0 <sup>c</sup>                   |
| 5     |    | <b>7t</b> 98% <sup>a</sup> (2 h)   | <b>7t</b> 98% <sup>b</sup> (20 h)<br>0 <sup>c</sup>  |
| 6     |   | <b>8e</b> 72% <sup>a</sup> (4 h)   | <br><b>8e</b><br><b>8e</b> 71% <sup>b</sup> (7d)<br>0 <sup>c</sup> |
| 7     |  | <b>9i</b> 65% <sup>a</sup> (4 h)<br><b>9t</b> 6% <sup>a</sup>                                  | <b>9i</b> 5% <sup>b</sup> (7d)<br>0 <sup>c</sup>   |
| 8     |  | <b>10i</b> 60% <sup>a</sup> (4 h)<br><b>10a</b> 6% <sup>a</sup><br><b>10t</b> 16% <sup>a</sup> | <b>10i</b> 52% <sup>b</sup> (7d)<br><b>10a</b> 6% <sup>b</sup><br><b>10t</b> 13% <sup>a</sup><br>0 <sup>c</sup>  |

All reactions run in CDCl<sub>3</sub> (filtered over basic Al<sub>2</sub>O<sub>3</sub> prior to use) at 40°C; <sup>a</sup>determined via <sup>1</sup>H NMR <sup>b</sup>addition of TBAB (1.5 eq. relative to capsule I), reaction run for 7 d at 40°C <sup>c</sup>addition of HOAc (10 mol%), no hexamer present, 7d, 40°C.

## RESULTS AND DISCUSSION

Similarly to the enzyme pocket (see above), catalytic amounts of capsule **I** (10 mol%) also facilitated the activation of tertiary alkyl fluorides. In light of our recent investigations (Köster and Tiefenbacher, 2018) concerning the role of HCl as a cocatalyst for some reactions catalyzed by capsule **I**, it is important to note that the activation of tertiary fluorides reported here does not require an acid cocatalyst.

The aryl-substituted substrates **3** and **4** (Table 1, entries 1 and 2) were mainly transformed to the corresponding homostyrene **3i** and styrene **4i**, respectively. In both cases the formation of

the trisubstituted alkene was favored and the terminal alkenes **3t** and **4t** were observed to a smaller degree. Brønsted-acid catalyzed isomerization of homostyrene **3i** to the corresponding styrene was not observed. Furthermore, an intramolecular Friedel-Crafts alkylation could not be observed. Considerable amounts of homobenzylic alcohol **4a** were formed as a side product. It is important to note that no background reaction was observed when the capsule was blocked with inhibitor **2** (see Figure 1C, 1.5 eq. tetrabutylammonium bromide per capsule) or when the capsule was omitted and replaced with an acid of similar acidity (acetic acid). These control experiments indicate that the reaction takes place inside the binding pocket of capsule **I**.

The main product of the reaction of ester **5** was found to be lactone **5l**, presumably resulting from a nucleophilic attack of the ester carbonyl oxygen on the tertiary carbocation. The elimination pathway also led to the formation of the tri-substituted alkene **5i** and of terminal alkene **5t** in trace amounts. Again, no background reactions were observed.




When the unsaturated fluoride **6** was investigated, the main product was found to be cyclic ether **6e** (see **Table 1**). Nucleophilic interception of the initially formed tertiary carbocation by water (from the hexameric assembly or the bulk solution) presumably leads to the formation of an alkenol. The ensuing cyclization is known to be catalyzed by capsule **I** (Catti and Tiefenbacher, 2015). As a side reaction, elimination to the trisubstituted alkene was observed.


Camphene-derived fluoride **7** was converted to the alkene **7t** within 2 h quantitatively. The addition of inhibitor **2** did not result in suppression of the reaction but only slowed down the reaction (completion after 20 h). The complete selectivity toward elimination may be explained by the steric constraints of the carbocation, with a quaternary center and a bridgehead atom in close proximity, which effectively shield the cation from the attack of any nucleophiles.

The fluoroalcohol **8** was converted to the seven-membered cyclic ether with good selectivity. The addition of inhibitor **2** did not prevent the reaction but slowed it down significantly (7 d instead of 4 h). The benzyl-protected analog **9** was transformed mainly to the corresponding trisubstituted alkene **9i** because the cyclization pathway was blocked. Experiments with a blocked cavity revealed only marginal conversion after 7 d. The  $\beta$ -Citronellol-derived fluoride **10** showed complete conversion within 4 h with the major product being the trisubstituted alkene. The formation of the corresponding eight-membered cyclic ether could not be observed. As observed with **8**, substantial conversion could be observed in presence of **2**, although at reduced rates (7 d instead of 4 h).

A possible explanation for the conversion of substrates in the presence of inhibitor may lie in the ability of capsule **I** to bind substrates via H-bonds from the outside. However, the significantly reduced reaction rates point toward an acceleration of the reaction on the inside of **I**. All substrates remained intact upon prolonged exposure to 10 mol% acetic acid ( $pK_A = 4.75$ ), indicating that the hexamer's ( $pK_A \sim 5.5-6$ ) mode of activation is likely based on hydrogen bonding to the substrate and stabilization of the resulting cation via cation- $\pi$  interactions instead of Brønsted acidity alone.

After having demonstrated the ability of capsule **I** to activate tertiary alkyl fluorides, we wondered whether the Friedel-Crafts alkylation with benzylic fluoride electrophiles was also possible inside the hexameric capsule **I**. Starting with 4-tert-butyl benzyl fluoride (**11**) and *p*-xylene (**Scheme 1**), a reaction to the expected product could be observed, albeit in low yield (17%). Since it is known that unfunctionalized guests





**FIGURE 3** | Reaction progress for the elimination of HF from **3** in presence of capsule **I**; 30 °C (□); 40 °C (●), 10 mol% HF, no capsule present, 40 °C (Δ).

are taken up poorly by container **I**, we employed 1,3,5-trimethoxybenzene as arene nucleophile to take advantage of the efficient uptake of oxygenated species by capsule **I**. This indeed resulted in an increase in yield to 51%. These results clearly indicate that the activation of benzylic fluorides is also possible and further corroborate capsule **I**'s ability to accelerate Friedel-Crafts reactions as was recently demonstrated by La Manna et al. (2018b). In the absence of suitable nucleophiles, oligomerization was observed with primary and secondary benzyl fluorides. When benzylic di- and trifluorides were investigated, no reaction could be observed. Attempts to use tertiary alkyl fluorides as electrophiles for Friedel-Crafts reactions were unsuccessful, leading only to the respective elimination products.

Following the findings of Paquin et al. (Champagne et al., 2014, 2015a,b) we propose an autocatalytic cycle starting with the encapsulation of a substrate molecule (see **Scheme 2**). The fluorine substituent likely is interacting *via* a hydrogen bond to a water molecule of the assembly. Subsequent cleavage of the activated C-F bond then leads to formation of a carbocation and hydrogen bonded fluoride<sup>1</sup>. After elimination of a proton and subsequent recombination with fluoride to form HF, the alkene is released from the inner cavity. With its superior hydrogen bond donating ability (Chęcinska and Grabowski, 2006; Champagne et al., 2014) hydrogen fluoride acts as catalytically active species in ensuing cycles.

<sup>1</sup>If X = F, the ion is known as bifluoride, however, more equivalents of HF could be involved.

This mechanistic proposal was underlined by kinetic measurements of the reaction of **3** inside container **I**. After an induction period a sigmoidal reaction progress can be observed at 40 °C, a typical indication of an autocatalytic reaction (see **Figure 3**) (Anslyn and Dougherty, 2006).

Interestingly, the addition of HF did not initiate a rapid autocatalytic conversion in the absence of capsule **I**. Conversion of the starting fluoride after addition of catalytic amounts of hydrogen fluoride was sluggish, reaching only 2% after 8 h. A probable cause of this observation might be the fast equilibration of hydrogen fluoride with silicon of the glass reaction vessel walls, which greatly diminishes the local concentration of the catalytically active species. Although HF has been shown to exist in equilibrium with glass walls (Chapman and Levy, 1952), only minuscule amounts of HF may exist at any given time in solution. Interestingly, experiments in plastic vials (see **SI chapter 5.2**) indicated that this is not true. Addition of HF in the absence of hexamer **I** did not lead to a significant conversion even when the reaction was run in plastic vials. However, in the presence of hexamer **I** the addition of HF initiated the reaction immediately (no induction period observed). This strongly suggests a synergistic catalysis mediated by HF and capsule **I**. Control reactions with a large substrate (see **SI chapter 5.5**) that is unable to enter the cavity of **I**, showed no conversion within 4 days under standard reaction conditions (40 °C, acid-free CDCl<sub>3</sub>, 10 mol% **I**). However, a competition experiment together with benzyl ether **9** resulted in the full conversion of both substrates within 4 h (see **SI-Figure 6**). It is clearly visible that the smaller substrate **9** initiated the reaction and subsequently the larger substrate reacted, obviously outside of capsule **I**. In summary, this

suggests that: (1) capsule **I** is necessary to activate the substrate, (2) HF is only catalytically active in conjunction with capsule **I** (3) if HF has been liberated large substrates can also react outside of capsule **I**, in accordance with the proposed mechanism (see **Scheme 2**).

Kinetic experiments with monodeuterated substrate **3d** (see **SI chapter 5.3**) to elucidate the reaction mechanism resulted in a KIE of  $1.16 \pm 0.03$ . This is indicative of a secondary deuterium KIE arising from hyperconjugative stabilization of the formed cation, thereby indicating fluoride abstraction as rate-limiting (Anslyn and Dougherty, 2006). This result is in agreement with earlier observations made in the acid-catalyzed solvolysis of tertiary alkyl fluorides (Chapman and Levy, 1952).

In conclusion, we have demonstrated the ability of **I** to activate tertiary alkyl fluorides and primary and secondary benzylic fluorides. The hexameric assembly **I** presumably not only functions as a hydrogen bond donor to activate the C-F bond but is also able to stabilize the intermediate carbocations *via* cation- $\pi$  interactions. The stabilized intermediates underwent several transformations inside the cavity, ranging from elimination, cyclization to intermolecular nucleophilic attack in the case of benzyl fluorides. The displayed ability of hexamer **I** to activate aliphatic C-F bonds is noteworthy since the reaction conditions are

unusually mild: 40°; no strong Brønsted or Lewis acids present.

## AUTHOR CONTRIBUTIONS

KT conceived and supervised the project. KT and JK planned the project. JK carried out all the experiments. DH performed kinetic NMR measurements. JK and KT compiled the first draft of the manuscript. All authors contributed to the final version of the manuscript.

## ACKNOWLEDGMENTS

This work was supported by funding from the European Research Council Horizon 2020 Programme [ERC Starting Grant 714620-TERPENECAT], the Swiss National Science Foundation as part of the NCCR Molecular Systems Engineering and the Bayerische Akademie der Wissenschaften (Junges Kolleg).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fchem.2018.00639/full#supplementary-material>

## REFERENCES

- Amii, H., and Uneyama, K. (2009). C-F bond activation in organic synthesis. *Chem. Rev.* 109, 2119–2183. doi: 10.1021/cr800388c
- Anslyn, E. V., and Dougherty, D. A. (2006). *Modern Physical Organic Chemistry*. Sausalito, CA: University Science Books.
- Avram, L., and Cohen, Y. (2002). The role of water molecules in a resorcinarene capsule as probed by NMR diffusion measurements. *Org. Lett.* 4, 4365–4368. doi: 10.1021/ol0271077
- Bellezza, F., Cipiciani, A., Ricci, G., and Ruzziconi, R. (2005). On the enzymatic hydrolysis of methyl 2-fluoro-2-arylpropionates by lipases, *Tetrahedron* 61, 8005–8012. doi: 10.1016/j.tet.2005.06.007
- Bianchini, G., La Sorella, G. L., Canever, N., Scarso, A., and Strukul, G. (2013). Efficient isonitrile hydration through encapsulation within a hexameric self-assembled capsule and selective inhibition by a photo-controllable competitive guest. *Chem. Commun.* 49, 5322–5324. doi: 10.1039/c3cc42233j
- Bräuer, T. M., Zhang, Q., and Tiefenbacher, K. (2017). Iminium catalysis inside a self-assembled supramolecular capsule: scope and mechanistic studies. *J. Am. Chem. Soc.* 139, 17500–17507. doi: 10.1021/jacs.7b08976
- Brown, C. J., Toste, F. D., Bergman, R. G., and Raymond, K. N. (2015). Supramolecular catalysis in metal-ligand cluster hosts. *Chem. Rev.* 115, 3012–3035. doi: 10.1021/cr4001226
- Burdeniuc, J., Jedicka, B., and Crabtree, R. H. (1997). Recent advances in C-F bond activation. *Chem. Ber.* 130, 145–154. doi: 10.1002/cber.19971300203
- Caneva, T., Sporni, L., Strukul, G., and Scarso, A. (2016). Efficient epoxide isomerization within a self-assembled hexameric organic capsule. *RSC Adv.* 6, 83505–83509. doi: 10.1039/C6RA20271C
- Catti, L., Pöthig, A., and Tiefenbacher, K. (2017). Host-catalyzed cyclodehydration–rearrangement cascade reaction of unsaturated tertiary alcohols. *Adv. Synth. Catal.* 359, 1331–1338. doi: 10.1002/adsc.201601363
- Catti, L., and Tiefenbacher, K. (2015). Intramolecular hydroalkoxylation catalyzed inside a self-assembled cavity of an enzyme-like host structure. *Chem. Commun.* 51, 892–894. doi: 10.1039/C4CC08211G
- Catti, L., and Tiefenbacher, K. (2018). Brønsted acid-catalyzed carbonyl-olefin metathesis inside a self-assembled supramolecular host. *Angew. Chem. Int. Ed.* 57, 14589–14592. doi: 10.1002/anie.201712141
- Catti, L., Zhang, Q., and Tiefenbacher, K. (2016a). Self-assembled supramolecular structures as catalysts for reactions involving cationic transition states. *Synthesis* 48, 313–328. doi: 10.1055/s-0035-1560362
- Catti, L., Zhang, Q., and Tiefenbacher, K. (2016b). Advantages of catalysis in self-assembled molecular capsules. *Chem. Eur. J.* 22, 9060–9066. doi: 10.1002/chem.201600726
- Champagne, P. A., Benhassine, Y., Desroches, J., and Paquin, J. F. (2014). Friedel–Crafts reaction of benzyl fluorides: selective activation of C-F bonds as enabled by hydrogen bonding. *Angew. Chem.* 126, 14055–14059. doi: 10.1002/ange.201406088
- Champagne, P. A., Desroches, J., and Paquin, J.-F. (2015a). Organic fluorine as a hydrogen-bond acceptor: recent examples and applications. *Synthesis* 47, 306–322. doi: 10.1055/s-0034-1379537
- Champagne, P. A., Drouin, M., Legault, C. Y., Audubert, C., and Paquin, J.-F. (2015b). Revised mechanistic explanation for the alcohol-promoted amination of benzylic fluorides under highly concentrated conditions: computational and experimental evidence on a model substrate. *J. Fluor. Chem.* 171, 113–119. doi: 10.1016/j.jfluchem.2014.08.018
- Chapman, N. B., and Levy, J. L. (1952). Nucleophilic displacement of fluorine from organic compounds. Part III. Acid-catalyzed solvolysis of alkyl fluorides in aqueous ethanol. *J. Chem. Soc.* 1677–1682. doi: 10.1039/jr9520001677
- Checinska, L., and Grabowski, S. J. (2006). F-H...F-C hydrogen bonds—The influence of hybridization of carbon atom connected with F-acceptor on their properties. *Chem. Phys.* 327, 202–208. doi: 10.1016/j.chemphys.2006.04.011
- Cullen, W., Misuraca, M. C., Hunter, C. A., Williams, N. H., and Ward, M. D. (2016). Highly efficient catalysis of the Kemp elimination in the cavity of a cubic coordination cage. *Nat. Chem.* 8:231. doi: 10.1038/nchem.2452
- Fave, G. M. L. (1949). Some reactions of the trifluoromethyl group in the benzotrifluoride series. I. Hydrolysis. *J. Am. Chem. Soc.* 71, 4148–4149. doi: 10.1021/ja01180a507
- Giust, S., La Sorella, G., Sporni, L., Fabris, F., Strukul, G., and Scarso, A. (2015). Supramolecular catalysis in the synthesis of substituted 1 H-tetrazoles from isonitriles by a self-assembled hexameric capsule. *Asian J. Org. Chem.* 4, 217–220. doi: 10.1002/ajoc.201402229
- Grether, U., Kimbara, A., Nettekoven, M., Ricklin, F., Roever, S., Rogers-Evans, M., et al. (2013). WO/2014/086805.

- Grummitt, A. R., Rutledge, P. J., Clifton, I. J., and Baldwin, J. E. (2004). Active-site-mediated elimination of hydrogen fluoride from a fluorinated substrate analogue by isopenicillin N synthase. *Biochem. J.* 382, 659–666. doi: 10.1042/BJ20040529
- Jans, A. C., Gómez-Suárez, A., Nolan, S. P., and Reek, J. N. (2016). A Switchable gold catalyst by encapsulation in a self-assembled cage. *Chem. Eur. J.* 22, 14836–14839. doi: 10.1002/chem.201603162
- Köster, J., and Tiefenbacher, K. (2018). Elucidating the importance of hydrochloric acid as a cocatalyst for resorcinarene-capsule-catalyzed reactions. *ChemCatChem* 10, 2941–2944. doi: 10.1002/cctc.201800326
- Kuehnel, M. F., Lentz, D., and Braun, T. (2013). Synthesis of fluorinated building blocks by transition-metal-mediated hydrodefluorination reactions. *Angew. Chem. Int. Ed.* 52, 3328–3348. doi: 10.1002/anie.201205260
- Kuijpers, P. F., Otte, M., Dürr, M., Ivanović-Burmazović, I., Reek, J. N. H., and de Bruin, B. (2016). A self-assembled molecular cage for substrate-selective epoxidation reactions in aqueous media. *ACS Catal.* 6, 3106–3112. doi: 10.1021/acscatal.6b00283
- La Manna, P., De Rosa, M., Talotta, C., Gaeta, C., Soriente, A., Floresta, G., et al. (2018a). The hexameric resorcinarene capsule as an artificial enzyme: ruling the regio and stereochemistry of a 1,3-dipolar cycloaddition between nitrones and unsaturated aldehydes. *Org. Chem. Front.* 5, 827–837. doi: 10.1039/C7QO00942A
- La Manna, P., Talotta, C., Floresta, G., De Rosa, M., Soriente, A., Rescifina, A. et al. (2018b). Mild Friedel-Crafts reactions inside a hexameric resorcinarene capsule: C-Cl bond activation through hydrogen bonding to bridging water molecules. *Angew. Chem. Int. Ed.* 57, 5423–5428. doi: 10.1002/anie.2018101642
- La Sorella, G., Sporni, L., Ballester, P., Strukul, G., and Scarso, A. (2016a). Hydration of aromatic alkynes catalyzed by a self-assembled hexameric organic capsule. *Catal. Sci. Technol.* 6, 6031–6036. doi: 10.1039/C6CY00307A
- La Sorella, G., Sporni, L., Strukul, G., and Scarso, A. (2016b). Supramolecular activation of hydrogen peroxide in the selective sulfoxidation of thioethers by a self-assembled hexameric capsule. *Adv. Synth. Catal.* 358, 3443–3449. doi: 10.1002/adsc.201600430
- Leenders, S., Gramage-Doria, R., de Bruin, B., and Reek, J. N. H. (2015). Transition metal catalysis in confined spaces. *Chem. Soc. Rev.* 44, 433–448. doi: 10.1039/C4CS00192C
- Levin, M. D., Kaphan, D. M., Hong, C. M., Bergman, R. G., Raymond, K. N., and Toste, F. D. (2016). Scope and mechanism of cooperativity at the intersection of organometallic and supramolecular catalysis. *J. Am. Chem. Soc.* 138, 9682–9693. doi: 10.1021/jacs.6b05442
- MacGillivray, L. R., and Atwood, J. L. (1997). A chiral spherical molecular assembly held together by 60 hydrogen bonds. *Nature* 389, 469. doi: 10.1038/38985
- Marchetti, L., and Levine, M. (2011). Biomimetic catalysis. *ACS Catal.* 1, 1090–1118. doi: 10.1021/cs200171u
- O'Hagan, D. (2008). Understanding organofluorine chemistry. An introduction to the C-F bond. *Chem. Soc. Rev.* 37, 308–319. doi: 10.1039/B711844A
- Otte, M. (2016). Size-selective molecular flasks. *ACS Catal.* 6, 6491–6510. doi: 10.1021/acscatal.6b01776
- Raynal, M., Ballester, P., Vidal-Ferran, A., and van Leeuwen, P. W. (2014). Supramolecular catalysis. Part 2: artificial enzyme mimics. *Chem. Soc. Rev.* 43, 1734–1787. doi: 10.1039/C3CS60037H
- Rosenberg, R. E. (2012). Does fluoromethane form a hydrogen bond with water? *J. Phys. Chem. A* 116, 10842–10849. doi: 10.1021/jp308533b
- Shen, Q., Huang, Y.-G., Liu, C., Xiao, J.-C., Chen, Q.-Y., and Guo, Y. (2015). Review of recent advances in CF bond activation of aliphatic fluorides. *J. Fluor. Chem.* 179, 14–22. doi: 10.1016/j.jfluchem.2015.07.007
- Ueda, Y., Ito, H., Fujita, D., and Fujita, M. (2017). Permeable self-assembled molecular containers for catalyst isolation enabling two-step cascade reactions. *J. Am. Chem. Soc.* 139, 6090–6093. doi: 10.1021/jacs.7b02745
- Wang, F., and Hu, J. (2009). Brønsted Acid Mediated (sp<sup>3</sup>)carbon-fluorine bond activation: inter- and intramolecular arylation of trifluoromethylated arenes. *Chin. J. Chem.* 27, 93–98. doi: 10.1002/cjoc.200990032
- Wang, Q.-Q., Gonell, S., Leenders, S. H., Dürr, M., Ivanović-Burmazović, I., and Reek, J. N. (2016). Self-assembled nanospheres with multiple endohedral binding sites pre-organize catalysts and substrates for highly efficient reactions. *Nat. Chem.* 8:225. doi: 10.1038/nchem.2425
- Wang, X., Nurttila, S. S., Dzik, W. I., Becker, R., Rodgers, J., and Reek, J. N. H. (2017). Tuning the porphyrin building block in self-assembled cages for branched-selective hydroformylation of propene. *Chem. Eur. J.* 23, 14769–14777. doi: 10.1002/chem.201703135
- Wenkens, M., Grønvald, F., and Hansen, J. B. (1998). Synthesis of meiosis-activating sterols containing fluorine. *Acta Chem. Scand.* 52, 503–507. doi: 10.3891/acta.chem.scand.52-0503
- Wiester, M. J., Ulmann, P. A., and Mirkin, C. A. (2011). Enzyme mimics based upon supramolecular coordination chemistry. *Angew. Chem. Int. Ed.* 50, 114–137. doi: 10.1002/anie.201000380
- Zarra, S., Wood, D. M., Roberts, D. A., and Nitschke, J. R. (2015). Molecular containers in complex chemical systems. *Chem. Soc. Rev.* 44, 419–432. doi: 10.1039/C4CS00165F
- Zhang, Q., and Tiefenbacher, K. (2013). Hexameric resorcinarene capsule is a Brønsted acid: investigation and application to synthesis and catalysis. *J. Am. Chem. Soc.* 135, 16213–16219. doi: 10.1021/ja4080375
- Zhang, Q., and Tiefenbacher, K. (2015). Terpene cyclization catalysed inside a self-assembled cavity. *Nat. Chem.* 7, 197–202. doi: 10.1038/nchem.2181

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Köster, Häussinger and Tiefenbacher. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.