


Direct Synthesis of Cyclopropanes from *gem*-Dialkyl Groups through Double C–H Activation

Antonin Clemenceau, Pierre Thesmar, Maxime Gicquel, Alexandre Le Flohic, and Olivier Baudoin*

ABSTRACT: Cyclopropanes are important structural motifs found in numerous bioactive molecules, and a number of methods are available for their synthesis. However, one of the simplest cyclopropanation reactions involving the intramolecular coupling of two C–H bonds on *gem*-dialkyl groups has remained an elusive transformation. We demonstrate herein that this reaction is accessible using aryl bromide or triflate precursors and the 1,4-Pd shift mechanism. The use of pivalate as the base was found to be crucial to divert the mechanistic pathway toward the cyclopropane instead of the previously obtained benzocyclobutene product. Stoichiometric mechanistic studies allowed the identification of aryl- and alkylpalladium pivalates, which are in equilibrium via a five-membered palladacycle. With pivalate, a second C(sp³)–H activation leading to the four-membered palladacycle intermediate and the cyclopropane product is favored. A catalytic reaction was developed and showed a broad scope for the generation of diverse arylcyclopropanes, including valuable bicyclo[3.1.0] systems. This method was applied to a concise synthesis of lemborexant, a recently approved anti-insomnia drug.



INTRODUCTION

Cyclopropanes are widespread structural motifs in natural products and bioactive molecules (see Figure 1 for examples).¹




Figure 1. Examples of drugs or drug candidates containing an arylcyclopropane motif relevant to this study.

Indeed, the cyclopropane ring can improve the pharmacological properties of active substances by reducing their lipophilicity, increasing their metabolic stability, or modifying their acido-basicity.² Although a great number of methods have been developed to synthesize cyclopropanes,³ the cycloaddition of alkenes and carbenes or carbenoids largely prevails for applications to complex molecules.⁴ In this context, one of the simplest ways to construct cyclopropanes, through C–C


bond formation from two geminal alkyl groups (Scheme 1a), has remained an elusive transformation to date. In 2006, Yu and co-workers reported the synthesis of cyclopropanes via a two-step sequence involving palladium(II)-catalyzed, oxazoline-directed C(sp³)–H diiodination of *gem*-dimethyl groups followed by radical cyclization. (Scheme 1b).⁵ Besides, cyclopropanes have been generated through domino reactions including olefin carbopalladation and C(sp³)–H activation, leading to putative 4-membered palladacyclic intermediates (Scheme 1c).^{6,7}

The current work reports the first single-step synthesis of cyclopropanes via coupling between two geminal methyl groups or one methyl and one activated methylene group (Scheme 1d). This reaction is based on the propension of the arylpalladium halide, generated in the initial oxidative addition step, to undergo 1,4-Pd shift to generate a σ -alkylpalladium intermediate.⁸ Since seminal observations by Dyker,⁹ such alkylpalladium intermediates have been exploited in various transformations: β -H elimination to generate olefins,¹⁰ trapping with boronic acids¹¹ and anilines,¹² or intramolecular dearomatizing carbopalladation, furnishing spiroannulation products.¹³ Recently, we showed that such organopalladium species are also able to perform the cleavage of an activated

Received: May 30, 2020

Published: August 12, 2020

Scheme 1. Synthesis of Cyclopropanes by Pd-Catalyzed C(sp³)-H Activation




C(sp³)-H bond (i. e., benzylic and adjacent to a carbonyl group) to generate oxygen and nitrogen heterocycles.¹⁴ The extension of this mechanism to the formation of cyclopropanes is reported herein.

RESULTS AND DISCUSSION

Initial Work and Mechanistic Study. The origin of this work stems from our initial studies on the generation of

Scheme 2. Influence of the Base on the Formation of Benzocyclobutene vs Cyclopropane



^a¹H NMR yield using CH₂Br₂ as internal reference. ^bUsing Pd₂dba₃ (5 mol %)/P(*t*-Bu)₃ (20 mol %) instead of Pd(PPh₃)₄.

benzocyclobutenes (BCBs) by Pd⁰-catalyzed C(sp³)-H activation.¹⁵ Recent reinvestigations pointed to a spectacular influence of the base on the reaction outcome (Scheme 2). As initially reported, using potassium carbonate as the base in the intramolecular C-H arylation of substrate **1a** under the shown conditions provided BCB **2a** as the sole observable C-H activation product (entry 1), together with the protodehalogenated byproduct (not shown). Replacing carbonate with pivalate led to a complete change of selectivity, with cyclopropane **3a** becoming the sole C-H activation product (entry 2). The ligand also had a noticeable effect on the selectivity, with the bulky P(*t*-Bu)₃ favoring the BCB product, as previously reported (entry 3).¹⁵

Following these initial observations, and inspired from a study by Martin and co-workers,¹⁶ we decided to reinvestigate the reaction mechanism by following each step through stoichiometric experiments (see the Supporting Information

for details). A tentative mechanism, represented in qualitative energy profile format, is displayed in Figure 2. The oxidative addition of prototypical aryl bromide **1b** to Pd(PPh₃)₄ gives rise to complex **A**, which undergoes bromide substitution with pivalate, producing complex **B**¹. The latter is in equilibrium with complex **B**², containing one phosphine ligand and pivalate coordinated in the κ² mode,^{17c} which undergoes base-mediated C(sp³)-H activation via the concerted metalation-deprotonation mechanism^{15,17} to give five-membered palladacycle **C**¹. Reductive elimination from **C**¹ provides the BCB product **2b**. Alternatively, protonation of **C**¹ with pivalic acid^{15,17e} furnishes alkylpalladium complex **D**¹. The sequence **B**¹ → **C**¹ → **D**¹ results in an overall 1,4-Pd shift.^{8,14} A second base-induced C(sp³)-H activation from **D**¹ leads to the rare four-membered palladacycle **E**^{1,6,7,18} already suggested by Martin,¹⁶ which reductively eliminates to generate cyclopropane **3b**.

The oxidative addition of the bulky aryl bromide **1b** was followed by ¹H and ³¹P NMR and was found to be slow using PPh₃ as the ligand, with only 26% of complex **A** formed at 80 °C after 16 h, and 72% at 120 °C together with 20% remaining starting material (Scheme 3a). Complex **A** was independently synthesized in 88% yield from **1b** and Pd(PPh₃)₄ (toluene, 120 °C), and its three-dimensional structure was confirmed by X-ray diffraction analysis (Scheme 3). Consistent with results displayed in Scheme 2, heating **A** to 100 °C in the presence of K₂CO₃ furnished BCB **2b** in 65% yield (Scheme 3b). The substitution step was then analyzed by mixing complex **A** with KOPiv (Scheme 3b). This step was relatively easy, with σ-aryl-palladium pivalate **B**¹ being slowly formed at room temperature (25% after 16 h, not shown) and in 82% yield at 60 °C. Raising the temperature to 80 and 100 °C led to the formation of cyclopropane **3b** at the expense of complex **B**¹. This result shows that the formation of the cyclopropane product is faster than the oxidative addition and, therefore, that the latter is the slowest step of the reaction. The reaction of the known σ-alkylpalladium complex (PhCMe₂CH₂)Pd(COD)Cl **D**² (COD = cyclooctadiene)¹⁹ in the presence of KOPiv and PPh₃ in toluene at room temperature led to alkylpalladium pivalate **D**¹ in 88% yield (Scheme S1, see Figure 2 or Scheme 3 for the structure of **D**¹). In contrast, at 60 °C arylpalladium pivalate **B**¹ was formed from **D**², thus showing that **B**¹ is more stable than **D**¹. The thermal decomposition of **B**¹ was next analyzed at various temperatures (Scheme 3c). At 80 °C, both **D**¹ and cyclopropane **3b** were formed, and the proportion of cyclopropane markedly increased at 100 °C. The reactivity of alkylpalladium pivalate **D**¹ was also analyzed (Scheme 3d). At 80 °C, a new complex, which was assigned to arylpalladium species **B**² containing one PPh₃ ligand and the κ²-coordinated pivalate, was formed together with degradation products. Adding PPh₃ (1 equiv) led to a cleaner transformation, with both mono- and bis-ligated complexes **B**¹-**B**² and cyclopropane **3b** being formed in comparable amounts. Heating this mixture to 100 °C furnished a higher proportion of **B**² and cyclopropane **3b**, and upon addition of 1 equiv of KOPiv the cyclopropane was formed exclusively. These results show that aryl- and alkylpivalate complexes **B**¹, **B**² and **D**¹ are in equilibrium from 80 °C and are all competent intermediates en route to cyclopropane **3b**. Finally, since we did not observe the formation of five-membered palladacycles such as **C**¹ (Figure 2) during the above studies, likely because they are higher-energy intermediates, we decided to prepare the known²⁰ five-membered COD-stabilized palladacycle **C**² and study its reactivity (Scheme 3e). In the presence of KOPiv, PivOH,





Figure 2. Qualitative energy profile of the reaction of substrate **1b** based on experimental observations. The relative stabilities of **B²** and **D¹**, **C¹** and **E¹**, and **2b** and **3b** were calculated by DFT (see the [Supporting Information](#) for details). L = PPh₃. TS = transition state; OA = oxidative addition; Sub = substitution; CH = C–H activation; PROT = protonation; RE = reductive elimination.

Scheme 3. Stepwise Study of the Reaction Mechanism through Stoichiometric Experiments^a




^a¹H NMR yields using CH₂Br₂ as internal reference. All reactions were performed in C₆D₆ for 16 h. ^bThermal ellipsoids at 50% probability, H atoms omitted for clarity. COD = 1,5-cyclooctadiene.

and PPh₃ at room temperature, complex **D¹** was formed as major product, together with minor amounts of **B¹**, thereby showing that the protonation with PivOH is kinetically favored on the aromatic vs the alkyl ligand. Heating to 80 °C inverted the ratio between **D¹** and **B¹**, thus confirming the higher thermodynamic stability of **B¹**, and produced a minor amount


of cyclopropane **3b**. At 100 °C, only the latter was formed (61% yield). In contrast, heating palladacycle **C²** to 80 °C in the presence of PPh₃ only provided **2b** in good yield.¹⁶

Taken together, these experimental observations allow to deduce the following relative stabilities for organopalladium complexes: **B¹** > **B²** > **D¹** > **C¹**. In addition, DFT calculations


Scheme 4. Reaction Scope^a



a) disubstituted cyclopropanes ($R^2 = H$)



b) trisubstituted cyclopropanes ($R^2 \neq H$)



^aReactions were performed on a 0.2 mmol scale. The aryl bromide precursor was employed unless otherwise noted. Yields refer to the isolated product or mixture of diastereoisomers. Diastereomeric ratios (d.r.) were determined by ¹H NMR of the crude mixture. Relative configurations were determined by NOESY NMR. ^bPerformed at 120 °C without DMSO. ^c¹H NMR yield using CH₂Br₂ as internal reference. ^dPerformed on a 1 mmol scale using *o*-xylene as the solvent.



indicated that the four-membered palladacycle **E**¹ is less stable than the five-membered isomer **C**¹ by ca. 7 kcal mol⁻¹, hence establishing the stability order **B**¹ > **B**² > **D**¹ > **C**¹ > **E**¹. Furthermore, cyclopropane **3b** was found to be more stable than BCB **2b** by 1.7 kcal mol⁻¹. Figure 2 provides a qualitative overview of the reaction profile summarizing all experimental results and calculations. The oxidative addition is the highest kinetic barrier, which arises from the fact that the relatively electronically neutral PPh₃ was employed as ligand. In the presence of carbonate and the absence of pivalate, reductive elimination from five-membered palladacycle **C**¹ occurs preferentially to give BCB product **2b**. In contrast, pivalate promotes the formation of cyclopropane **3b** via C(sp³)-H activation of a second methyl group producing the four-membered palladacycle intermediate **E**¹. Both bicarbonate²¹ and pivalic acid are able to open up palladacycle **C**¹ to give alkylpalladium intermediates, and therefore the unique ability of pivalate to promote the second C(sp³)-H activation event leading to cyclopropanation remains unclear and will be the object of later studies.

Reaction Optimization and Scope. Taking advantage of the above mechanistic study, a catalytic version of this transformation was successfully developed (see the Supporting

Information for details). Unsubstituted cyclopropane **3b** was obtained from both aryl bromide and triflate precursors in good yield using 10 mol % Pd(PPh₃)₄ and 2 equiv potassium pivalate in toluene at 120 °C (Scheme 4a). However, this volatile product was best isolated from the triflate precursor in 72% yield. For functionalized cyclopropanes, a temperature of 140 °C was essential to ensure complete conversion. Besides, the addition of DMSO (5% vol.) was found to diminish the formation of the proto-dehalogenated side-product, presumably by increasing the solubility of pivalate.

With these standard conditions in hand, the versatility of this double C(sp³)-H activation-based cyclopropanation was inspected. Substrates bearing an ester (**3a** and **3c**), amide (**3d**), protected amine (**3e**), or protected alcohol (**3i** and **3m**) on the quaternary benzylic carbon worked well under the optimized conditions (72–85% yields). Of note, the reaction producing cyclopropane **3a** was successfully scaled up to 1 mmol using *o*-xylene as the solvent for practical reasons. In addition, substrate **1f** bearing a substituted linear alkyl chain provided a mixture of the desired cyclopropane **3f** (50% yield) and olefin **4f** arising from methylene C-H activation and β-H elimination, consistent with previous work.^{10a,17e} Reactants containing various electron-donating groups on the aromatic

Scheme 5. Synthesis of Cyclopropane-Fused Lactones and Lactams




^aConditions: $\text{Pd}(\text{PPh}_3)_4$ (10 mol %), KOiPr (2 equiv), toluene/DMSO (95:5), 140 °C, 16 h. ^bNMR yield. ^cYield of the isolated product. ^dYield of the isolated minor diastereoisomer. ^eRatio determined by ¹H NMR of the crude mixture, relative configuration determined by NOESY NMR.

ring such as dimethoxy (**3g–i**), methylenedioxy (**3j**), or TIPS-protected phenol (**3k**) gave the desired cyclopropanes in good yields (65–81%). Likewise, electron-withdrawing groups such as fluoro (**3l**, **3m**, and **3o**), nitro (**3n**), and trifluoromethyl (**3p**) were found to be compatible. The position of the substituent on the aromatic ring (meta or para) did not have a significant influence on the reaction outcome. In addition, the pyridinylcyclopropane **3q** was generated in 70% yield, hence indicating the compatibility of heteroarenes. Moreover, substrates functionalized in ortho position to the quaternary carbon (**1r**) or to the bromine atom (**1s** and **1t**) provided the corresponding cyclopropane products **3r–t** in good yields (69–78%). Remarkably, a 2-fold cyclopropanation reaction was successfully carried out, allowing formation of bis-cyclopropane **3u** in 92% yield via two consecutive cyclopropanations including four $\text{C}(\text{sp}^3)\text{--H}$ activation steps.

As expected, substrates bearing benzyl instead of methyl groups did not undergo $\text{C}(\text{sp}^3)\text{--H}$ activation and cyclopropanation, but rather direct $\text{C}(\text{sp}^2)\text{--H}$ arylation, as illustrated with the 6-membered ring product **5** (Scheme 4b). In contrast, in the case of substrates bearing a quaternary carbon with one methyl and one methylene adjacent to an ester or nitrile group, the construction of trisubstituted cyclopropanes was possible. First, cyclopropanes **3aa** and **3ab** containing two esters were synthesized in average yields and good diastereoselectivity, probably arising from thermodynamic control. The positional isomers **3ac** and **3ad** containing nitrile groups were also produced, albeit with a reduced diastereoselectivity. Finally, the trisubstituted cyclopropane **3ae** was generated in 86% combined yield and with a weak, but opposite diastereoselectivity compared to **3ab**.

Molecules containing an aryl-substituted [3.1.0] bicyclic system show interesting biological activities,² but only a few methods have been developed for their synthesis^{22–26} The access to these attractive scaffolds was investigated by applying

Scheme 6. Application to the Synthesis of Lemborexant^a



^aConditions: 1. MeOH, H_2SO_4 (0.1 equiv), 50 °C; 2. LDA (1.2 equiv), MeI (1.05 equiv), -78 to $+25$ °C; 3. LiHMDS (3 equiv), allyl bromide (5 equiv), THF, 0 to 25 °C (72% for 3 steps); 4. DIBAL-H (2.1 equiv), Et_2O , -78 to $+25$ °C (82%); 5. RuCl_3 (5 mol %), NaIO_4 (6 equiv), MeCN/ H_2O 10:1, 25 °C; 6. $\text{Pd}(\text{PPh}_3)_4$ (10 mol %), KOiPr (2 equiv), *o*-xylene/DMSO 95:5, 140 °C; 7. LiHMDS (3 equiv), **14** (3 equiv), toluene, 25 °C (89%); 8. **16** (1.5 equiv), PPh_3 (1.5 equiv), DIAD (1.5 equiv), THF, 25 °C (68%).

our newly developed Pd-catalyzed double $\text{C}(\text{sp}^3)\text{--H}$ activation reaction (Scheme 5). Starting from lactone **6a**, the cyclopropane-fused product **7a** was formed in low yield and the major product was the butenolide **8a** resulting from β -hydrogen elimination (Scheme 5a). A similar result was obtained from lactam **6b**. To suppress the olefin formation, the reaction was conducted with γ -lactones and lactams **9** bearing the quaternary carbon in β position to the carbonyl group (Scheme 5b). Gratifyingly, this modification led to the cyclopropane-fused lactones **10a–b** and lactam **10c** in 65–79% yields. To further increase the complexity, a more substituted substrate bearing a $\text{CH}_2\text{CO}_2\text{Me}$ instead of the CH_3 group was employed. A separable 3:1 diastereomeric mixture of fused γ -lactams was obtained in 79% combined yield, with the major diastereoisomer **10c** having the aryl and ester groups in cis relationship. This example shows that our methodology can also be employed to couple two activated methylene groups for the synthesis of tetrasubstituted cyclopropanes.

Lemborexant **11** (Dayvigo, Figure 1) is a dual antagonist of the orexin OX_1 and OX_2 receptors, which was recently approved for the treatment of insomnia.²⁷ The application of the current method to a short synthesis of this bioactive target is depicted in Scheme 6. Esterification of commercially available aryl bromide **12** and sequential alkylation and allylation led to methyl ester **13**. The latter was subjected to ester reduction and oxidative cleavage of the alkene, which

directly provided γ -butyrolactone **9e**.²⁸ The key cyclopropanation of **9e** was performed under the optimized conditions, to furnish the fused lactone **10e** in good yield (79%). Direct amidation of the latter with aminopyridine **14** and LiHMDS according to Szostak and co-workers,²⁹ followed by Mitsunobu reaction with pyrimidinol **16** furnished racemic lemborexant **11** in 28% overall yield over 8 steps.

CONCLUSIONS

The elusive intramolecular cyclopropanation reaction cleaving two C–H bonds and forging a C–C bond has been developed by taking advantage of the 1,4-Pd shift mechanism. Stoichiometric studies indicated that oxidative addition is the rate-limiting step with the employed substrates and catalyst. The characterization of intermediate Pd complexes allowed the delineation of a mechanistic pathway including σ -aryl- and σ -alkylpalladium pivalates, which are in equilibrium via the five-membered diorganopalladacycle. With pivalate as the base and PPh₃ as the ligand, the reductive elimination from this palladacycle is disfavored and the activation of a second C(sp³)–H bond is favored, leading to a high-energy four-membered palladacycle which reductively eliminates to give the cyclopropane product. A catalytic version was developed, which enabled the generation of an array of arylcyclopropanes in good yields, including fused ring systems, from simple aryl bromide or triflate precursors. The generation of fused cyclopropanes was applied to a straightforward synthesis of the anti-insomnia drug lemborexant. This work further illustrates that tuning the catalyst and base in Pd-catalyzed C–H activation reactions may allow to divert mechanistic pathways toward the formation of high value-added products.

ASSOCIATED CONTENT

Full optimization tables, crystallographic, procedural, and spectral data (PDF)

Crystallographic data of a compound as discussed in the text (CIF)

DFT-optimized structures of the compounds as discussed in the text (XYZ)

AUTHOR INFORMATION

Corresponding Author

Olivier Baudoin – Department of Chemistry, University of Basel, CH-4056 Basel, Switzerland; orcid.org/0000-0002-0847-8493; Email: olivier.baudoin@unibas.ch

Authors

Antonin Clemenceau – Department of Chemistry, University of Basel, CH-4056 Basel, Switzerland

Pierre Thesmar – Department of Chemistry, University of Basel, CH-4056 Basel, Switzerland

Maxime Gicquel – Oril Industrie, 76210 Bolbec, France

Alexandre Le Flohic – Oril Industrie, 76210 Bolbec, France

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was financially supported by Oril Industrie, affiliated to Les Laboratoires Servier, and the University of Basel. We thank Dr. Lucile Vaysse-Ludot, Dr. R. Tamion, and Dr. J. Fournier, ORIL Industrie, for fruitful discussions and constant support; Dr. D. Häussinger, University of Basel, for NMR experiment; Dr. Alessandro Prescimone, University of Basel, for Xray diffraction analysis; and Dr. M. Pfeffer, University of Basel, for MS analyses.

REFERENCES

- (1) Faust, R. Fascinating Natural and Artificial Cyclopropane Architectures. *Angew. Chem., Int. Ed.* **2001**, *40*, 2251–2253.
- (2) (a) Talele, T. T. The "Cyclopropyl Fragment" is a Versatile Player that Frequently Appears in Preclinical/Clinical Drug Molecules. *J. Med. Chem.* **2016**, *59*, 8712–8756. (b) Ćasar, Z. Synthetic Approaches to Contemporary Drugs that Contain the Cyclopropyl Moiety. *Synthesis* **2020**, *52*, 1315–1345.
- (3) (a) Roy, M.-N.; Lindsay, V. N. G.; Charette, A. B. Cyclopropanation Reactions. In *Science of Synthesis*; De Vries, J. G., Molander, G. A., Evans, P. A., Eds.; Thieme: Stuttgart, 2011; Vol. 1, pp 731–817. (b) Kulinkovich, O. G. *Cyclopropanes in Organic Synthesis*; John Wiley & Sons, Inc.: Hoboken, NJ, 2015.
- (4) (a) Chen, D. Y.-K.; Pouwer, R. H.; Richard, J.-A. Recent Advances in the Total Synthesis of Cyclopropane-Containing Natural Products. *Chem. Soc. Rev.* **2012**, *41*, 4631–4642. (b) Ebner, C.; Carreira, E. M. Cyclopropanation Strategies in Recent Total Syntheses. *Chem. Rev.* **2017**, *117*, 11651–11679.
- (5) Giri, R.; Wasa, M.; Breazzano, S. P.; Yu, J.-Q. Converting gem-Dimethyl Groups into Cyclopropanes via Pd-Catalyzed Sequential C–H Activation and Radical Cyclization. *Org. Lett.* **2006**, *8*, 5685–5688.
- (6) (a) Liron, F.; Knochel, P. Reactivity of Stable Neopentyl-Pd Intermediates in the Absence of Nucleophile. *Tetrahedron Lett.* **2007**, *48*, 4943–4946. (b) Kim, H. S.; Gowrisankar, S.; Kim, S. H.; Kim, J. N. Synthesis of 6-Oxacyclopropa[a]indene Derivatives Starting from Baylis-Hillman Adducts via Pd-Mediated C(sp³)–H Activation. *Tetrahedron Lett.* **2008**, *49*, 3858–3861. (c) Huang, Q.; Larock, R. C. Synthesis of Cyclopropanes by Pd-Catalyzed Activation of Alkyl C–H Bonds. *Tetrahedron Lett.* **2009**, *50*, 7235–7238. (d) Mao, J.; Zhang, S.-Q.; Shi, B.-F.; Bao, W. Palladium(0)-Catalyzed Cyclopropanation of Benzyl Bromides via C(sp³)–H Bond Activation. *Chem. Commun.* **2014**, *50*, 3692–3694. (e) Du, W.; Gu, Q.; Li, Z.; Yang, D. Palladium(II)-Catalyzed Intramolecular Tandem Aminoalkylation via Divergent C(sp³)–H Functionalization. *J. Am. Chem. Soc.* **2015**, *137*, 1130–1135. (f) Chung, D. S.; Lee, J. S.; Ryu, H.; Park, J.; Kim, H.; Lee, J. H.; Kim, U. B.; Lee, W. K.; Baik, M.-H.; Lee, S.-g. Palladium-Catalyzed Divergent Cyclopropanation by Regioselective Solvent-Driven C(sp³)–H Bond Activation. *Angew. Chem., Int. Ed.* **2018**, *57*, 15460–15464.
- (7) For the construction of cyclopropanes via C–C activation and C(sp³)–H activation: Cao, J.; Chen, L.; Sun, F.-N.; Sun, Y.-L.; Jiang, K.-Z.; Yang, K.-F.; Xu, Z.; Xu, L.-W. Pd-Catalyzed Enantioselective Ring Opening/Cross-Coupling and Cyclopropanation of Cyclobutanones. *Angew. Chem., Int. Ed.* **2019**, *58*, 897–901.
- (8) (a) Ma, S.; Gu, Z. 1,4-Migration of Rhodium and Palladium in Catalytic Organometallic Reactions. *Angew. Chem., Int. Ed.* **2005**, *44*, 7512–7517. (b) Shi, F.; Larock, R. C. Remote C–H Activation via Through-Space Palladium and Rhodium Migrations. *Top. Curr. Chem.* **2009**, *292*, 123–164. (c) Rahim, A.; Feng, J.; Gu, Z. 1,4-Migration of Transition Metals in Organic Synthesis. *Chin. J. Chem.* **2019**, *37*, 929–945.
- (9) (a) Dyker, G. Palladium-Catalyzed C–H Activation of Methoxy Groups: A Facile Synthesis of Substituted 6H-Dibenzo[b,d]pyrans. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 1023–1025. (b) Dyker, G. Palladium-Catalyzed C–H Activation of *tert*-Butyl Groups: A Simple Synthesis of 1,2-Dihydrocyclobutabenzene Derivatives. *Angew. Chem.,*

- Int. Ed. Engl.* **1994**, *33*, 103–105. (c) Dyker, G. Palladium-Catalyzed C–H Activation at Methoxy Groups: Regiochemistry of the Domino Coupling Process. *Chem. Ber.* **1994**, *127*, 739–742.
- (10) (a) Baudoin, O.; Herrbach, A.; Guéritte, F. The Palladium-Catalyzed C–H Activation of Benzylic *gem*-Dialkyl Groups. *Angew. Chem., Int. Ed.* **2003**, *42*, 5736–5740. (b) Hitce, J.; Retailliau, P.; Baudoin, O. Palladium-Catalyzed Intramolecular C(sp³)–H Functionalization: Catalyst Development and Synthetic Applications. *Chem. - Eur. J.* **2007**, *13*, 792–799. (c) Motti, E.; Catellani, M. Catalytic Dehydrogenation of *o*-Alkylated or *o*-Alkoxyated Iodoarenes with Concomitant Hydrogenolysis. *Adv. Synth. Catal.* **2008**, *350*, 565–569. (d) Bheeter, C. B.; Jin, R.; Bera, J. K.; Dixneuf, P. H.; Doucet, H. Palladium-Catalyzed Dehydrogenative sp³ C–H Bonds Functionalisation into Alkenes: A Direct Access to *N*-Alkylbenzenesulfonamides. *Adv. Synth. Catal.* **2014**, *356*, 119–124.
- (11) Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. Catalysts for Suzuki–Miyaura Coupling Processes: Scope and Studies of the Effect of Ligand Structure. *J. Am. Chem. Soc.* **2005**, *127*, 4685–4696.
- (12) Pan, J.; Su, M.; Buchwald, S. L. Palladium(0)-Catalyzed Intermolecular Amination of Unactivated C_{sp³}–H Bonds. *Angew. Chem., Int. Ed.* **2011**, *50*, 8647–8651.
- (13) Tan, B.; Bai, L.; Ding, P.; Liu, J.; Wang, Y.; Luan, X. Palladium-Catalyzed Intermolecular [4 + 1] Spiroannulation by C(sp³)–H Activation and Naphthol Dearomatization. *Angew. Chem., Int. Ed.* **2019**, *58*, 1474–1478.
- (14) Rocaboy, R.; Anastasiou, I.; Baudoin, O. Redox-Neutral Coupling between Two C(sp³)–H Bonds Enabled by 1,4-Palladium Shift for the Synthesis of Fused Heterocycles. *Angew. Chem., Int. Ed.* **2019**, *58*, 14625–14628.
- (15) Chaumontet, M.; Piccardi, R.; Audic, N.; Hitce, J.; Peglion, J.-L.; Clot, E.; Baudoin, O. Synthesis of Benzocyclobutenes by Palladium-Catalyzed C–H Activation of Methyl Groups: Method and Mechanistic Study. *J. Am. Chem. Soc.* **2008**, *130*, 15157–15166.
- (16) Gutiérrez-Bonet, A.; Juliá-Hernández, F.; de Luis, B.; Martín, R. Pd-Catalyzed C(sp³)–H Functionalization/Carbenoid Insertion: All Carbon Quaternary Centers via Multiple C–C Bond Formation. *J. Am. Chem. Soc.* **2016**, *138*, 6384–6387.
- (17) (a) Lafrance, M.; Gorelsky, S. I.; Fagnou, K. High-Yielding Palladium-Catalyzed Intramolecular Alkane Arylation: Reaction Development and Mechanistic Studies. *J. Am. Chem. Soc.* **2007**, *129*, 14570–14571. (b) Kefalidis, C. E.; Baudoin, O.; Clot, E. DFT Study of the Mechanism of Benzocyclobutene Formation by Palladium-Catalyzed C(sp³)–H Activation: Role of the Nature of the Base and the Phosphine. *Dalton Trans.* **2010**, *39*, 10528–10535. (c) Rousseaux, S.; Gorelsky, S. I.; Chung, B. K. W.; Fagnou, K. Investigation of the Mechanism of C(sp³)–H Bond Cleavage in Pd(0)-Catalyzed Intramolecular Alkane Arylation Adjacent to Amides and Sulfonamides. *J. Am. Chem. Soc.* **2010**, *132*, 10692–10705. (d) Rousseaux, R.; Davi, M.; Sofack-Kreutzer, J.; Pierre, C.; Kefalidis, C. E.; Clot, E.; Fagnou, K.; Baudoin, O. Intramolecular Palladium-Catalyzed Alkane C–H Arylation from Aryl Chlorides. *J. Am. Chem. Soc.* **2010**, *132*, 10706–10716. (e) Kefalidis, C. E.; Davi, M.; Holstein, P. M.; Clot, E.; Baudoin, O. Mechanistic Study on the Selectivity of Olefin versus Cyclobutene Formation by Palladium(0)-Catalyzed Intramolecular C(sp³)–H Activation. *J. Org. Chem.* **2014**, *79*, 11903–11910.
- (18) (a) McNally, A.; Haffemayer, B.; Collins, B. S. L.; Gaunt, M. J. Palladium-Catalyzed C–H Activation of Aliphatic Amines to Give Strained Nitrogen Heterocycles. *Nature* **2014**, *510*, 129–133. (b) He, C.; Gaunt, M. J. Ligand-Enabled Catalytic C–H Arylation of Aliphatic Amines by a Four-Membered-Ring Cyclopalladation Pathway. *Angew. Chem., Int. Ed.* **2015**, *54*, 15840–15844. (c) Su, B.; Bunescu, A.; Qiu, Y.; Zuend, S. J.; Ernst, M.; Hartwig, J. F. Palladium-Catalyzed Oxidation of β-C(sp³)–H Bonds of Primary Alkylamines through a Rare Four-Membered Palladacycle Intermediate. *J. Am. Chem. Soc.* **2020**, *142*, 7912–7919.
- (19) Gutiérrez, E.; Nicasio, M. C.; Paneque, M.; Ruiz, C.; Salazar, V. Synthesis and Reactivity of New Palladium Alkyl Complexes Containing PMe₃ Ligands: Insertion Reactions and Formation of Bis(pyrazolyl)borate Derivatives. *J. Organomet. Chem.* **1997**, *549*, 167–176.
- (20) Cámpora, J.; López, J. A.; Palma, P.; del Rio, D.; Carmona, E.; Valerga, P.; Graiff, C.; Tiripicchio, A. Synthesis and Insertion Reactions of the Cyclometalated Palladium–Alkyl Complexes Pd(CH₂CMe₂-*o*-C₆H₄)L₂. Observation of a Pentacoordinated Intermediate in the Insertion of SO₂. *Inorg. Chem.* **2001**, *40*, 4116–4126.
- (21) The opening of five-membered palladacycle C¹ under carbonate-mediated conditions was indirectly observed through the formation BCB isomers and deuterium incorporation on the aryl ring.¹⁵
- (22) 2-Aryl cyclopropane-fused γ-lactones: (a) Milewska, M. J.; Gdaniec, M.; Polonski, T. Synthesis, Stereochemistry, and Chiroptical Spectra of Cyclopropyl Lactones and Thionolactones. *Tetrahedron: Asymmetry* **1996**, *7*, 3169–3180. (b) Muraoka, O.; Tanabe, G.; Momose, T. Accentuation of the Di-*p*-Methane Reactivity by Central Carbon Substitution in the 4-(Phenylmethyl)-2(5H)-furanone. *Heterocycles* **1990**, *31*, 1589–1592.
- (23) (a) Shuto, S.; Ono, S.; Hase, Y.; Kamiyama, N.; Matsuda, A. Synthesis of (+)- and (–)-milnaciprans and their conformationally restricted analogs. *Tetrahedron Lett.* **1996**, *37*, 641–644. (b) Doyle, M. P.; Hu, W. A New Enantioselective Synthesis of Milnacipran and an Analogue by Catalytic Asymmetric Cyclopropanation. *Adv. Synth. Catal.* **2001**, *343*, 299–302.
- (24) 1-Aryl cyclopropane-fused γ-lactams: Wang, K.-B.; Ran, R.-Q.; Xiu, S.-D.; Li, C.-Y. Synthesis of 3-Aza-bicyclo[3.1.0]hexan-2-one Derivatives via Gold-Catalyzed Oxidative Cyclopropanation of *N*-Allylynamides. *Org. Lett.* **2013**, *15*, 2374–2377.
- (25) Pedroni, J.; Cramer, N. Enantioselective C–H Functionalization–Addition Sequence Delivers Densely Substituted 3-Azabicyclo[3.1.0]hexanes. *J. Am. Chem. Soc.* **2017**, *139*, 12398–12401.
- (26) Zhang, Z.-Z.; Han, Y.-Q.; Zhan, B.-B.; Wang, S.; Shi, B.-F. Synthesis of Bicyclo[*n*.1.0]alkanes by a Cobalt-Catalyzed Multiple C(sp³)–H Activation Strategy. *Angew. Chem., Int. Ed.* **2017**, *56*, 13145–13149.
- (27) Yoshida, Y.; Naoe, Y.; Terauchi, T.; Ozaki, F.; Doko, T.; Takemura, A.; Tanaka, T.; Sorimachi, K.; Beuckmann, C. T.; Suzuki, M.; Ueno, T.; Ozaki, S.; Yonaga, M. Discovery of (1*R*,2*S*)-2-[(2,4-Dimethylpyrimidin-5-yl)oxy]methyl]-2-(3-fluorophenyl)-*N*-(5-fluoropyridin-2-yl)cyclopropanecarboxamide (E2006): A Potent and Efficacious Oral Orexin Receptor Antagonist. *J. Med. Chem.* **2015**, *58*, 4648–4664.
- (28) Gao, R.; Fan, R.; Canney, D. J. Synthesis of β,β-Disubstituted γ-Butyrolactones by Chemoselective Oxidation of 1,4-Diols and γ-Hydroxy Olefins with RuCl₃/NaIO₄. *Synlett* **2015**, *26*, 661–665.
- (29) Li, G.; Ji, C.-L.; Hong, X.; Szostak, M. Highly Chemoselective, Transition-Metal-Free Transamidation of Unactivated Amides and Direct Amidation of Alkyl Esters by *N*–C/O–C Cleavage. *J. Am. Chem. Soc.* **2019**, *141*, 11161–11172.