



CONCISE REVIEW

Therapeutic vascularization in regenerative medicine

Roberto Gianni-Barrera¹ | Nunzia Di Maggio¹ | Ludovic Melly^{1,2} |
Maximilian G. Burger^{1,3} | Edin Mujagic^{1,4} | Lorenz Gürke⁴ | Dirk J. Schaefer³ |
Andrea Banfi^{1,3,4} 

¹Cell and Gene Therapy, Department of Biomedicine, Basel University Hospital and University of Basel, Basel, Switzerland

²Cardiac, Vascular, and Thoracic Surgery, CHU UCL Namur, Yvoir, Belgium

³Plastic and Reconstructive Surgery, Department of Surgery, Basel University Hospital and University of Basel, Basel, Switzerland

⁴Vascular Surgery, Department of Surgery, Basel University Hospital and University of Basel, Basel, Switzerland

Correspondence

Andrea Banfi, MD, Department of Biomedicine, Basel University Hospital, Hebelstrasse 20, 4031 Basel, Switzerland. Email: andrea.banfi@usb.ch

Funding information

Department of Surgery of Basel University Hospital; Swiss Nanoscience Institute; European Union H2020 Program, Grant/Award Numbers: 801159, 646075; Swiss National Science Foundation, Grant/Award Number: 182357

Abstract

Therapeutic angiogenesis, that is, the generation of new vessels by delivery of specific factors, is required both for rapid vascularization of tissue-engineered constructs and to treat ischemic conditions. Vascular endothelial growth factor (VEGF) is the master regulator of angiogenesis. However, uncontrolled expression can lead to aberrant vascular growth and vascular tumors (angiomas). Major challenges to fully exploit VEGF potency for therapy include the need to precisely control in vivo distribution of growth factor dose and duration of expression. In fact, the therapeutic window of VEGF delivery depends on its amount in the microenvironment around each producing cell rather than on the total dose, since VEGF remains tightly bound to extracellular matrix (ECM). On the other hand, short-term expression of less than about 4 weeks leads to unstable vessels, which promptly regress following cessation of the angiogenic stimulus. Here, we will briefly overview some key aspects of the biology of VEGF and angiogenesis and discuss their therapeutic implications with a particular focus on approaches using gene therapy, genetically modified progenitors, and ECM engineering with recombinant factors. Lastly, we will present recent insights into the mechanisms that regulate vessel stabilization and the switch between normal and aberrant vascular growth after VEGF delivery, to identify novel molecular targets that may improve both safety and efficacy of therapeutic angiogenesis.

KEYWORDS

extracellular matrix, genetic therapy, ischemia, neovascularization, tissue engineering, vascular endothelial growth factor

1 | INTRODUCTION

Blood vessel growth is an integral process in regenerative medicine. Vascular regulation is also key for the repair of naturally avascular tissues such as cartilage, where inhibition of angiogenesis has been

shown to favor the spontaneous chondrogenic differentiation of progenitors in vivo.¹ Focusing only on the situations where new vascular induction is required, two main areas of therapeutic interest can be conceptually distinguished: (a) the expansion of pre-existing vascular networks in ischemic tissues to restore blood flow and salvage function (therapeutic angiogenesis); and (b) the rapid de novo vascular invasion of engineered grafts to enable progenitor survival and differentiation.

Roberto Gianni-Barrera and Nunzia Di Maggio contributed equally to the study.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. STEM CELLS TRANSLATIONAL MEDICINE published by Wiley Periodicals, Inc. on behalf of AlphaMed Press

Ischemia is caused by an inadequate blood supply for tissue demand. The most prevalent cause is progressive atherosclerotic stenosis of cardiac or limb arteries (coronary or peripheral artery disease, respectively; CAD and PAD), which can be compounded by microvascular dysfunction in metabolic syndromes such as diabetes. The consequences of CAD range from chest pain upon exertion (*angina*) to myocardial infarction and end-stage heart failure, whereas PAD leads to progressive muscle pain with walking (*claudicatio*), skin ulcerations, and necrosis, necessitating amputation. Impaired wound healing in diabetic patients, especially of the foot, can have dire consequences through infection and necrosis. Together with their high prevalence in western-style societies, cardiovascular ischemic diseases pose a heavy social and economic burden, due to early mortality, impaired quality of life with pain and loss of mobility, and recurrent hospitalization.² Current treatments for CAD and PAD mainly include endovascular procedures such as balloon angioplasty/stenting and bypass surgery or possibly exercise training for limb ischemia. However, many patients with end stage CAD or PAD are not candidates for such procedures due to a lack of target vessels. On the other hand, standard treatment of diabetic foot ulcers is purely supportive.³ Therapeutic angiogenesis, that is, the generation of new vascular networks through delivery of specific growth factors, is an attractive strategy to restore perfusion to ischemic tissues and fill this unmet clinical need.

Another area of regenerative medicine in which vascular growth plays a key role is the vascularization of tissue-engineered grafts.⁴ In particular, bone replacement is required in several situations due to trauma, surgery, or idiopathic conditions such as avascular necrosis of small bones, where spontaneous regeneration is insufficient. Bone tissue-engineering holds promise for the generation of osteogenic grafts, combining osteo-progenitors with bio-compatible scaffolds.^{5,6} However, for defects of clinically relevant size, the lack of rapid vascularization *in vivo* causes severe ischemia and progenitor death in the graft core deeper than 1 to 2 mm.⁷ Similar biological principles apply to the vascularization of both ischemic tissue and engineered grafts, but with different translational considerations.

2 | BIOLOGICAL BASES OF THERAPEUTIC VASCULARIZATION

After the discovery of vascular endothelial growth factor (VEGF), first as a permeability factor in 1983⁸ and then as an endothelial mitogen in 1989,^{9,10} intense investigations began into the new concept of therapeutic angiogenesis, aiming at restoring the blood supply in ischemic tissues by growing new blood vessels with VEGF and later other angiogenic factors. Positive preclinical and early clinical evidence seemed to indicate great potential for this strategy. However, a decade of subsequent controlled clinical studies showed that the simple delivery of the VEGF protein or gene to ischemic tissue has no clear efficacy at safe doses.¹¹ The disappointing clinical results are a stark contrast to the fundamental biological role of VEGF as the

Significance statement

The promotion of blood vessel growth for therapeutic purposes remains a challenge both for the treatment of ischemic conditions and the generation of functional tissue-engineered grafts. Physiological angiogenesis is a complex and highly concerted process. A fine understanding of the cellular and molecular mechanisms of vascular growth needs to provide the biological basis for the design of rational therapeutic approaches.

master regulator of vascular growth. In fact, VEGF kicks off the complex cascade of cellular and molecular events leading to the orderly assembly of new endothelial structures (morphogenesis), their association with mural cells/pericytes (maturation), and subsequent ability to persist indefinitely in the absence of further growth factor signaling (stabilization), to form fully functional vascular networks.¹² Therefore, a better understanding of the physiological mechanisms of vascular growth is important to exploit its therapeutic potential. Here, we will address some key aspects of the biology of VEGF and angiogenesis, and their therapeutic implications.

2.1 | VEGF and its gradients

The mammalian VEGF family comprises five main ligands (VEGF-A, -B, -C, and -D and placenta-derived growth factor, PlGF) and three receptors (VEGF-R1, -R2, and -R3). Although the principal role of VEGF-C and -D is to stimulate lymphatic angiogenesis through VEGF-R3, blood vessel growth is mostly coordinated by the signaling of VEGF-A and -B and PlGF through R1 and R2 (for a comprehensive review, see reference 13). Despite the multiplicity of players, the molecular target for therapeutic angiogenesis is essentially VEGF-A signaling, as VEGF-B and PlGF play more accessory or tissue-specific roles.¹⁴ We will refer to VEGF-A simply as VEGF throughout the manuscript.

An important feature of VEGF function, with widespread therapeutic implications, is its interaction with extracellular matrix (ECM), which dictates its spatial localization in tissues and regulates the outcome of the angiogenic process. In fact, alternative mRNA splicing of the *Vegfa* transcript gives rise to three major isoforms with different degrees of affinity for the ECM.¹⁵ These comprise 120, 164, and 188 residues in rodents (or 121, 165, and 189 in humans, respectively) due to the presence or absence of heparin-binding domains that interact with ECM proteoglycans, so that matrix affinity is very low in the shortest isoform and increases with molecular size.¹⁶ The isoforms also display differences in their signaling. In fact, VEGF_{164/165} binds the coreceptor Neuropilin-1 (Nrp1), enhancing activation of VEGF-R2 and endothelial proliferation and migration, whereas VEGF_{120/121} does not.¹⁷ Furthermore, a distal splice site in the last exon of the VEGF gene can give rise to a second set of “b” isoforms, which differ only in the sequence of the

last six residues and are therefore named VEGF_{xxx**b**}. However, contrary to the classic isoforms, the b variants are antiangiogenic and provide a further layer of regulation to the angiogenic balance in tissues.¹⁸ As a consequence of differential matrix binding, VEGF_{120/121} is highly diffusible in tissues, VEGF_{188/189} remains extremely localized at the site of secretion and VEGF_{164/165} instead generates intermediate gradients of concentration around the producing cells. The importance of differential matrix affinity of VEGF isoforms was shown elegantly in transgenic mice selectively producing only one isoform from the endogenous locus, so that regulation of expression was not altered.¹⁹ Diffusible VEGF₁₂₀ induced malformed vessels, which were aberrantly enlarged and lacked branching, whereas vessels generated by sticky VEGF₁₈₈ showed opposite defects, with very small diameters and hyper-branching. VEGF₁₆₄ was the only isoform capable of inducing physiological vascular networks in the absence of the other ones, thanks to its intermediate matrix affinity. It should be noted that the key requirement for physiological VEGF function is a balance between diffusibility and binding, rather than a specific isoform. In fact, normal vascular morphogenesis also took place in the absence of VEGF₁₆₄, as long as VEGF₁₂₀ and VEGF₁₈₈ were both expressed. The importance of balanced matrix affinity therefore makes VEGF_{164/165} the isoform of choice for therapeutic delivery.

2.2 | Cellular mechanisms: Sprouting and intussusception

Sprouting is the best characterized cellular mechanism of angiogenesis and is the primary process by which new vessels grow out of pre-existing ones to invade surrounding tissue, for example, during embryonic development, endochondral ossification, menstrual decidua regeneration, or tumor vascularization. Sprouting entails the specification of endothelium into two functionally distinct phenotypes, that is, tip and stalk cells, and is guided by the formation of VEGF concentration gradients.²⁰ The first endothelial cell reacting to VEGF becomes a tip, which extends numerous thin filopodia from the basal side into the surrounding matrix to sense the gradient and migrates toward its source (Figure 1A). Each tip cell instructs its neighboring cells to acquire the stalk phenotype and these proliferate to form the new vessel trunk (Figure 1B). Interestingly, while tip cells respond to the gradient of VEGF distribution, stalk cell proliferation is regulated by its absolute concentration.²⁰ Finally, to form a functional network the tip cells of two vascular sprouts first make filopodia contacts with each other, then fuse and finally form a new continuous perfused lumen through a complex series of cellular rearrangements²¹ (Figure 1C). Tip cell fusion can also be promoted by macrophages, which act as chaperones between the contacting filopodia²² (Figure 1B). The balanced formation of tip and stalk cells is finely regulated by Dll4/Notch1 signaling, whereby the first endothelial cell sensing the VEGF gradient becomes a tip by default and upregulates the ligand Dll4. This activates Notch1 on the neighboring cells, which are so instructed to downregulate both VEGF-R2 and Dll4 expression and acquire a stalk phenotype, through a mechanism of lateral inhibition.²³

Disturbance of Notch1 signaling causes excessive tip cell formation and increased sprouting, which, however, is accompanied by impaired stalk cell generation and therefore leads to nonfunctional endothelial structures that have no lumen and are not perfused.²³

Vascular network expansion can take place also by the alternative mechanism of intussusception (from the Latin meaning "growth within itself"), also referred to as splitting angiogenesis.²⁴ Intussusception can be initiated very rapidly by increased blood flow and shear stress in the absence of growth factors,²⁵ but it can also follow VEGF upregulation.²⁶ It is still unclear what determines whether VEGF induces sprouting or angiogenesis, but its distribution in matrix is likely to play a role and the absence of a concentration gradient appears to favor intussusception. For example, both outcomes have been described in skeletal muscle: spontaneous upregulation of VEGF by ischemia leads to sprouting,²⁷ but its over-expression at significantly higher and therapeutic levels, which saturate the little matrix between fibers and abrogate local gradients, causes angiogenesis by intussusception.²⁶ Contrary to sprouting, no tip cells are formed and activated endothelial cells respond exclusively by proliferation without migration, leading first to circumferential enlargement of the vessel (Figure 1D,E), which then splits longitudinally into new daughter vessels. Splitting requires the formation of tissue pillars across the vascular lumen. These can form through two alternative processes: (a) a vascular wall invagination that creates a contact between the opposite endothelial cells²⁸ (Figure 1F), or (b) the extension and fusion of intraluminal filopodial-like protrusion from the endothelium²⁹ (Figure 1G,J,K). Subsequently, the endothelial junctions reorganize and myofibroblast invade the core, stabilizing the structures into mature transluminal tissue pillars (Figure 1H,L,M). Finally, these align along the length of the vessel, fuse together, and divide the affected vascular segment longitudinally (Figure 1I).

In recent years, it has become increasingly clear that intussusception is a therapeutically important mechanism of angiogenesis.^{12,26,30} The molecular regulation of intussusception is still poorly understood and likely quite different from that of sprouting. Inhibition of Notch signaling in the absence of growth factor delivery has been described to stimulate intussusceptive vascular expansion in the chicken chorio-allantoic membrane³¹ and in the mouse liver.³² On the other hand, recent data show that the outcome of VEGF-induced intussusceptive angiogenesis can be modulated by pericyte recruitment by platelet-derived growth factor-BB (PDGF-BB)³³ and stimulation of endothelial EphB4 signaling by pericyte-expressed ephrinB2.³⁴

2.3 | PDGF-BB and the pericyte-endothelium crosstalk

After endothelial assembly, new microvascular networks need to undergo maturation by associating with pericytes. This important cell type has exquisitely regulatory functions and provides signals that switch off endothelial proliferation and permeability and make the new vessels stable, that is, independent of continued VEGF stimulation and able to persist indefinitely after the angiogenic signals subside. Pericytes are recruited by PDGF-BB secreted by activated




FIGURE 1 Sprouting and intussusception: two alternative modes of angiogenesis. Schematic representation of the processes generating new vascular structures by sprouting (A–C) or by intussusception (vascular splitting; D–I). J–M, Immunofluorescence images of vessels undergoing intussusception after vascular endothelial growth factor delivery in murine skeletal muscle, stained for endomucin (endothelial cells, green), laminin (basal lamina, red), and with DAPI (nuclei, blue). Circumferentially enlarged vessels displayed: (a) no degradation of the basement membrane; (b) intraluminal filopodia-like protrusions from the endothelial layer (white arrowheads in high-magnification panels K and M); and (c) mature intraluminal tissue pillars (white arrows in high-magnification panel M). * represents vascular lumen; scale bars = 20 μ m in all panels

endothelium and interference with this process is incompatible with embryonic development, leading to sustained endothelial proliferation, vessel wall fragility, and lethal bleeding.^{35,36} PDGF-BB function

also critically relies on its interaction with ECM and gradient formation. In fact, removal of its matrix-binding domain leads to pericyte detachment from nascent vessels and severe vascular dysfunction.³⁷

Pericytes regulate endothelial function through both paracrine and cell contact-dependent signals. The principal pathways mediating this molecular crosstalk are transforming-growth factor- β 1 (TGF- β 1), angiotensin-1 and -2 (Ang-1 and Ang-2), and ephrinB2/EphB4.³⁸ TGF- β 1 is produced in a latent form and its activation requires cell-to-cell contact between endothelial and mural cells through proteolytic cleavage of the latency-associated peptide by plasmin at the vessel wall interface.³⁹ TGF- β 1 regulates both the endothelium and pericytes and can promote contrasting functions, depending on alternative Alk1 and Alk5 receptor stimulation and downstream SMAD pathway activation.⁴⁰ In fact, activation of Alk5-SMAD2/3 specifically promotes vascular stabilization by inducing endothelial quiescence and stimulating the production of ECM and basement membrane proteins.⁴¹ In contrast, activation of Alk1-SMAD1/5 promotes endothelial cell proliferation and migration as well as endothelial-to-mesenchymal transition.⁴¹

Angiotensins are the ligands of the endothelium-specific tyrosine kinase receptor Tie2. Ang1 is expressed by mural cells and activates Tie2: it facilitates the further recruitment and association of pericytes with newly formed vascular structures, acts as a survival signal for endothelial cells and inhibits VEGF-induced vascular leakage.⁴² On the other hand, Ang2 is a context-dependent partial agonist of the Tie2 receptor, mostly acting as an inhibitor of its activation. It is stored preformed in Weibel–Palade bodies of endothelial cells, from which it is rapidly released upon VEGF stimulation. It plays a key role to initiate the angiogenic process by promoting the dissociation of pericytes and endothelial cells and allowing their formation of new vascular structures. However, it also promotes endothelial cell apoptosis and vascular regression if VEGF signaling is disrupted at this crucial stage.⁴²

Eph receptors and ephrin ligands are both membrane-bound and, therefore, binding and activation of Eph and ephrins requires cell–cell contact. Characteristically, engagement of the Eph–ephrin system elicits bidirectional signaling, that is, “forward signaling” in the cell expressing the Eph receptor and “reverse signaling” in the cell expressing the ephrin ligand.⁴³ In tumor angiogenesis, activation of reverse signaling via the EphB4 ligand ephrinB2 was shown to stimulate pericyte association to blood vessels and inhibit their permeability through the activation of the Ang1/Tie2 pathway.⁴⁴ Conversely, in a tissue-specific mutant model, ephrinB2-deficient pericytes failed to establish proper cell-to-cell contacts with microvascular endothelium, leading to hemorrhaging and perinatal mortality.⁴⁵ EphrinB2 has also been shown to directly modulate VEGF-R2 and VEGF-R3 activation in sprouting tip cells by controlling their internalization, necessary for receptor activation of downstream signaling.^{46,47} Recently, we also found that EphB4 activation also regulates splitting angiogenesis independently of VEGF-R2 internalization or phosphorylation, but rather by modulating the activation of ERK1/2 downstream of the receptor and the speed of endothelial proliferation.³⁴

2.4 | Circulating monocytes

Besides endothelial and mural cells, different populations of circulating myeloid cells play a crucial role in both induction and maturation of new vessels.⁴⁸ CXCR4+ monocytes are recruited by VEGF

expression, do not incorporate into new vessels (unlike circulating endothelial progenitors), but are retained in a perivascular position through Stromal Cell-Derived Factor-1 (SDF-1)/CXCR4 signaling and sustain vascular growth through further secretion of pro-angiogenic factors.⁴⁹ On the other hand, a similar CXCR4+ population of circulating monocytes has been recently shown to promote intussusceptive vascular growth during liver regeneration by participating in the formation and stabilization of intraluminal pillars.⁵⁰ Tissue macrophages also aid the completion of sprouting angiogenesis by promoting tip cell fusion and anastomosis, acting as cellular chaperones.²² These macrophages expressed both the surface receptors Tie2 and Neuropilin1 (Nrp1) and therefore were phenotypically similar to the family of pro-angiogenic Tie2-expressing macrophages (TEM). TEM have been described as tumor-infiltrating myeloid cells that convey pro-angiogenic signals and can provide a mechanism of resistance to anti-angiogenic drugs, promoting cancer progression.⁵¹ A population of circulating Neuropilin1-expressing monocytes (NEM) has been shown to also stimulate vascular maturation and smooth muscle cell recruitment. These cells coexpress CD11b and Nrp-1, which is a receptor for both VEGF and Semaphorin-3A (Sema3A), are recruited at sites of VEGF expression and promote the formation of arteries by the paracrine actions of several factors such as Ang1, TGF- β , and PBGF-BB.⁵² We recently found that NEM are specifically recruited to angiogenic sites by Sema3A produced by activated endothelium and accelerate stabilization of newly induced microvascular networks, that is, their ability to survive independently of VEGF signaling and persist indefinitely.⁵³

3 | THERAPEUTIC CONSIDERATIONS FOR VEGF DELIVERY

3.1 | The control of dose distribution

Therapeutic use of VEGF requires careful dose control, as uncontrolled delivery has been shown to cause aberrant vascular growth and angioma-like vascular tumors in muscle, heart and other tissues.^{54–57} The physiological affinity of VEGF for ECM has significant therapeutic implications for its dosing and delivery strategies. In fact, upon VEGF gene delivery, it remains tightly localized in the microenvironment around each producing cell and different levels of expression do not average with each other. Therefore, the therapeutic outcome between safety and efficacy of factor delivery is determined by the distribution of microenvironmental concentrations in tissue rather than simply the total dose delivered. This concept is exemplified by experiments in which the average and microenvironmental dose of VEGF could be controlled independently by cell-based gene delivery.⁵⁸ Different VEGF doses were expressed in skeletal muscle by populations of genetically modified muscle progenitor cells (myoblasts). Random genomic integration of the viral vector leads to different expression levels in each transduced cell. Implantation of this heterogeneous population always leads to aberrant angioma-like vascular growth, even if the total dose is reduced several-fold by dilution




FIGURE 2 Functional outcomes of vascular endothelial growth factor (VEGF) dose distribution in tissue. A, Heterogeneous dose distributions (eg, by gene therapy vectors) lead to hotspots of excessive expression that remains localized in the microenvironment around producing cells (red spots, upper right panel) and lead to toxic effects. Reducing the total dose does not completely avoid toxic hotspots even if therapeutic levels are achieved in some areas (blue spots, upper middle panel), until the total dose is so low that mostly ineffective levels are achieved (gray spots, upper left panel). B, Homogeneous distribution of the total dose allows therapeutic levels (blue spots, upper middle panel) to be achieved and to harness the therapeutic window of VEGF delivery

with nonexpressing cells. On the other hand, by isolating and expanding single myoblasts it is possible to obtain monoclonal populations in which every cell produces the same level. Under these conditions, which ensure a homogeneous microenvironmental distribution of the dose, it is possible to induce only normal and physiological angiogenesis across a wide range of doses, as well as to effectively restore functional blood flow in limb ischemia.⁵⁹ In contrast, the same total doses are toxic when delivered by dilutions of the heterogeneous population, because rare hotspots of excessive production cannot be avoided.⁵⁸

The need to control the distribution of microenvironmental doses is key when addressing the therapeutic window of VEGF delivery. In fact, when heterogeneous expression levels are generated in the tissue, such as by direct delivery of gene therapy vectors, effective doses in some microenvironments are accompanied also by ineffective and by toxic levels in other areas. In order to ensure safety, the total vector dose must be limited, but this increases the frequency of ineffective levels, thereby jeopardizing efficacy (Figure 2A). As a consequence, the usable therapeutic window of VEGF gene delivery appears to be very narrow.⁵⁵ On the other hand, VEGF does not have an intrinsically very steep dose-response curve and its biological therapeutic window is much larger,^{58,59} but it is key to control the distribution of doses in the microenvironment *in vivo* in order to exploit the therapeutic potential of VEGF delivery, ensuring both safety and efficacy (Figure 2B).

Translating this concept poses a fresh set of challenges. In fact, gene therapy is the preferred approach for therapeutic angiogenesis, thanks to a very robust delivery potential and standardized GMP vector production, but it is extremely difficult to avoid heterogeneous expression levels *in vivo* with uncontrolled vectors. Several groups are actively developing new generations of vectors, for example, ones that may self-regulate in the tissue,⁶⁰ and investigating other improvements in the gene therapy approaches: an excellent coverage of the latest discoveries can be found in a recent review by Ylä-Herttua and Baker.⁶¹

3.2 | Cell-based gene delivery

The distribution of expression levels *in vivo* can be controlled by a cell-based gene therapy approach. As detailed above, transduction of a progenitor population with an integrating viral vector leads to stable but heterogeneous expression levels in different cells. We developed a fluorescence-activated cell sorting (FACS)-based technology to predict the level of transgene expression in single live cells and to purify populations homogeneously expressing specific levels.⁶² The transgene of interest (eg, VEGF) is linked in a bicistronic viral vector to a nonfunctional truncated version of the lymphocyte-specific marker CD8a, which acts simply as a membrane-localized reporter protein. In this way, changes in the level of expression of VEGF are reflected by a parallel change in cell-surface expression of CD8a, which can be detected and quantified on live cells by FACS, enabling the purification of specific populations homogeneously expressing different VEGF levels (Figure 3). This technology can be used with different progenitor classes⁶³ for different applications. FACS purification of VEGF-expressing myoblasts was shown to yield only normal and functional angiogenesis in skeletal muscle,^{62,64} while purification of VEGF-expressing adipose-derived stromal cells ensured controlled vascular growth in the normal myocardium⁶⁵ and prevented deterioration of cardiac function in a model of myocardial infarction by limiting fibrotic scarring.⁶⁶

Cell-based VEGF gene delivery has also been investigated to promote the rapid vascularization of tissue-engineered grafts, for example, for cardiac patches or bone regeneration. Differently from the ischemic conditions described above, in these strategies, suitable progenitors are seeded on appropriate scaffolds and need to rapidly attract a vascular supply after *in vivo* implantation in order to differentiate and function. It is therefore desirable to link the angiogenic signal to the presence of the seeded cells by genetically engineering them to produce VEGF. Cardiac patches engineered with FACS-purified




FIGURE 3 Fluorescence-activated cell sorting (FACS)-purification of genetically modified progenitor populations expressing homogeneous vascular endothelial growth factor (VEGF) levels. A, Retroviral vector carrying a bicistronic cassette, in which the sequence of VEGF is linked to that of the membrane-bound reporter CD8 through an Internal Ribosome Entry Site (IRES) sequence. B, Progenitors of interest are transduced with this retroviral vector. C, After integration in the cell chromatin, the IRES sequence enables cotranslation of both proteins from the same mRNA molecule. Therefore, the amount of CD8 on the cell membrane reflects that of secreted VEGF regardless of their absolute level of expression. D, Transduced cells in the primary population express heterogeneous VEGF levels depending on their viral copy number and the transcriptional activity of the chromatin integration sites. E, These can be FACS-sorted into homogeneous subpopulations stably expressing specific VEGF levels based on the intensity of membrane CD8 staining

VEGF-expressing myoblasts⁶⁷ or adipose-derived stromal cells⁶⁸ could drive efficient vascularization of the patch itself and also provided controlled VEGF release to induce extrinsic angiogenesis in the underlying myocardium.

VEGF-expressing human bone marrow osteoprogenitors could effectively increase the vascularization of engineered bone grafts by threefold, inducing physiologically structured vascular networks with both conductance vessels and capillaries.⁶⁹ However, sustained and uncontrolled VEGF expression unexpectedly caused a global reduction in the quantity of bone, posing a challenge to its clinical application. Progenitor engraftment or differentiation was not impaired, but rather the recruitment of osteoclasts was strongly increased, thereby disrupting bone homeostasis toward excessive resorption. Current research is aiming at decoding the crosstalk between angiogenesis and osteogenesis, as well as ensuring that the two processes are therapeutically coupled.

3.3 | Recombinant factor engineering for matrix decoration

It is desirable to independently control the dose and duration of angiogenic factor delivery in grafts. This is technically challenging with gene transfer, while protein delivery suffers from short half-life *in vivo*. Extensive efforts have generated many strategies to ensure

sustained release of recombinant growth factors from natural and synthetic biomaterials. These approaches have been recently reviewed by Browne and Pandit.⁷⁰ However, it has also been possible to create modified versions of growth factors by protein engineering, so that they can be used to decorate natural matrices and be presented to cells in their physiological context during tissue regeneration.⁷¹ The most ubiquitous and abundant ECM protein is collagen, which is also employed as a basis for a variety of biomaterials. Therefore, recombinant growth factors have been engineered with specific collagen-binding domains, allowing their use to decorate collagen-based biomaterials and prolonging their bioavailability and efficacy (reviewed by Addi et al⁷²). On the other hand, tissue regeneration after damage starts in all cases with the deposition of a fibrin-based matrix rich in growth factors,⁷³ which provides ideal conditions for cell migration, vascular invasion and progenitor differentiation. Therefore, significant efforts have been directed at mimicking ECM decoration to ensure physiological presentation of morphogens.

Growth factors have been incorporated into fibrin matrix exploiting the coagulation process itself (Figure 4A, top). For example, murine VEGF₁₆₄ was fused to the octapeptide NQEQVSPL, which is the substrate of the transglutaminase coagulation factor XIIIa, allowing its covalent crosslinking into fibrin hydrogels and release only by enzymatic cleavage.⁷⁴ Further addition of the fibrinolysis inhibitor aprotinin, also engineered with the same technology, could finely tune the hydrogel degradation rate and therefore independently control




FIGURE 4 Strategies for matrix decoration with engineered recombinant factors. A, Fibrin matrices can be decorated with engineered growth factors to mimic extracellular matrix (ECM) functions. Taking advantage of the coagulation cascade, an octapeptide substrate of the TransGlutaminase factor XIII (TG hook) can be fused to growth factors (GFs), enabling their covalent crosslinking to fibrin. Specific domains of ECM proteins (eg, fibronectin) can also be incorporated through a TG hook to exploit their natural affinity for different GFs. B, Endogenous ECM can be decorated with therapeutic GFs engineered to exhibit super-affinity to a broad range of ECM components

the duration of factor release. Controlled VEGF delivery to skeletal muscle through this optimized platform was shown to yield exclusively normal, stable, and functional angiogenesis, over a wide range of easily controllable doses, and restored blood flow to ischemic tissues.⁷⁵

The natural affinity of different ECM proteins for growth factors can also be exploited (Figure 4A, bottom). For example, fibronectin has binding sites for fibrin, integrins, and growth factors. By engineering a recombinant fibronectin fragment comprising all these sites, it has been possible to codeliver the morphogens bone morphogenetic protein-2 and PDGF-BB within a fibrin matrix to promote bone regeneration.⁷⁶ Interestingly, presentation of the factors in their physiological matrix context greatly increased their functional efficacy, ensuring robust bone regeneration at very low and otherwise ineffective doses. Taking a reverse strategy, a short domain found in placenta-derived growth factor-2 was found to mediate broad binding to a wide variety of ECM proteins. Engineering of any growth factor with this peptide endowed them with super-affinity for ECM, thereby enabling the in situ decoration of endogenous matrix with exogenously provided therapeutic proteins⁷⁷ (Figure 4B). The increased efficacy of the modified factors avoided the need to deliver supra-physiological doses of VEGF, thereby increasing safety.

3.4 | Modulation of dose-dependent outcomes

An alternative class of approaches may help reduce the need to precisely control the distribution of VEGF doses in vivo. VEGF-A is a very powerful activator of VEGF-R2 signaling, but alternative ligands have been found to have milder activation profiles and therefore may require less stringent dose control. The properties of VEGF-B might particularly benefit applications to the cardiac muscle. In fact, VEGF-B is specifically active in the myocardium, while it is poorly angiogenic in other tissues.⁷⁸ Furthermore, VEGF-B does not bind to VEGF-R2, but rather activates it indirectly, by displacing inactive VEGF-A from

reservoir sites. Therefore, even robust over-expression of VEGF-B does not easily lead to excessive VEGF-R2 signaling, because it is limited to the endogenous levels of VEGF-A for its action.^{79,80} Another example is VEGF-D: in its native form it stimulates lymphatic angiogenesis through VEGF-R3, but it can also generate a shorter form by proteolytic cleavage of both the N- and C-termini, therefore named VEGF-D^{ΔNΔC}. VEGF-D^{ΔNΔC} shifts its affinity to VEGF-R2 and stimulates blood angiogenesis instead. However, because it has no heparin-binding domain, it leads to a more diffuse distribution through ECM and induces more physiological vascular growth than VEGF-A over-expression.⁸¹ Clinical trials of therapeutic angiogenesis by adenoviral delivery of VEGF-D^{ΔNΔC} are ongoing.^{82,83}

The outcome of VEGF over-expression may also be significantly modified by promoting pericyte recruitment. Studies in skeletal muscle showed that the transition between normal and aberrant angiogenesis is not determined exclusively by VEGF dose, but rather by the balance between endothelial activation by VEGF and pericyte recruitment by PDGF-BB.⁸⁴ Codelivery of VEGF₁₆₄ and PDGF-BB at a fixed relative ratio, achieved through coexpression from a single bicistronic vector, ensured the generation and long-term stability of exclusively normal and functional microvascular networks regardless of absolute VEGF dose,^{84,85} by limiting endothelial proliferation.³³ Although PDGF-BB alone does not induce vascular growth either in normal or ischemic skeletal muscle,^{84,86} the beneficial effects of the VEGF/PDGF-BB combination have been described in a variety of preclinical settings, including gene delivery with adenoviral⁸⁶ or adeno-associated viral vectors,⁸⁷ sustained release of the recombinant factors from polymeric biomaterials⁸⁸ or treatment with modified proteins engineered for super-affinity to the ECM.⁷⁷

As described above, pericytes exchange a complex molecular crosstalk with endothelium and these signaling pathways can offer more specific targets to regulate VEGF effects. For example, stimulation of the Tie2 receptor by Ang1 has been shown to significantly reduce the detrimental hyperpermeability and vascular leakage that accompany VEGF stimulation.⁸⁹ This therapeutic benefit can be

further compounded by the contemporaneous inhibition of Ang2.⁹⁰ On the other hand, activation of endothelial EphB4 by ephrinB2 finely tunes ERK1/2 phosphorylation downstream of VEGF-R2, thereby limiting the rate of endothelial proliferation induced by VEGF.³⁴ Therefore, systemic treatment with recombinant ephrinB2-Fc was shown to prevent aberrant angiogenesis by robust and uncontrolled VEGF expression, without interfering with efficient normal microvascular network formation.³⁴

Although VEGF is the principal target for therapeutic angiogenesis, the delivery of other factors has shown promising preclinical results, such as the fibroblast growth factor (FGF) family (FGF-1, FGF-2, and FGF-4) or hepatocyte growth factor, and clinical studies are currently underway to evaluate their potential in coronary or peripheral artery disease (CAD or PAD).⁶¹

3.5 | Duration of delivery and vessel stabilization

Another important consideration for the therapeutic effectiveness of newly induced vasculature is its long-term persistence. In fact, while new vascular networks are rapidly formed in a matter of few days after VEGF delivery, they are initially unstable and will regress if VEGF stimulation is withdrawn too early. Several lines of evidence in models of inducible transgene expression^{91,92} or pharmacologic blockade⁵⁸ support a need to sustain VEGF stimulation for about 4 weeks before new vessels are stabilized and persist independently. This requirement impairs the efficacy of transient VEGF gene delivery, for example, by adenoviral vectors that have been widely used in clinical trials and afford robust expression, but that are also cleared by the immune system in about 10 days.⁹³

More recently, we found that VEGF dose-dependently impairs the kinetics of vascular stabilization, that is, vessels induced by lower levels of VEGF stabilize faster. However, pericyte recruitment is not affected. Rather, VEGF negatively regulates the production of Sema3A by activated endothelium, which in turn impairs the recruitment of Nrp1-expressing monocytes, activation of TGF- β 1-SMAD2/3 signaling and the induction of endothelial quiescence.⁵³ In a therapeutic perspective, treatment with recombinant Sema3A was shown to accelerate the stabilization of newly induced angiogenesis and allow its persistence despite transient VEGF expression.⁵³ Interestingly, the stabilization of newly induced vasculature can also be promoted by the mechanical properties (low stiffness) of a biomaterial environment, through the recruitment of a novel population of mechano-sensitive Piezo1+ monocytes.⁹⁴

4 | CONCLUSIONS AND PERSPECTIVES

The biology of angiogenesis is complex and a better understanding of its cellular and molecular mechanisms is necessary for the design of rational and more effective therapeutic strategies. Therapeutic angiogenesis in ischemic tissues and de novo vascularization of engineered

grafts present rather different requirements and therefore might be best achieved by different approaches.

Tissue-engineered grafts are avascular upon implantation and need to attract vascular in-growth. In order to rapidly guide sprouting of new vessels and their migration toward the graft core, it is desirable that the graft matrix presents an optimized microenvironment of angiogenic cues. This can be achieved by predecorating a suitable material (such as fibrin or collagen) with optimized doses and combinations of engineered factors, as described above, or also by employing decellularized ECM, enriched in morphogens by suitable progenitor cell lines.⁹⁵ The process can also be accelerated by prevascularization, that is, preseeding with endothelial cells to generate self-assembling vascular structures inside the construct that can anastomose with the penetrating host vessels.

On the other hand, ischemic tissues are already vascularized and proangiogenic therapy aims at expanding the microvascular networks to promote collateral artery remodeling and restore physiological blood flow.¹³ For this, it is key to treat large volumes of target tissue and gene delivery by viral vectors has a clearly superior efficacy compared with both protein and cell-based approaches. For example, adenoviral vectors combine two clinically desirable features, namely robustness of expression and transient duration, limited by the immune response. However, these same features compromise safety (by uncontrolled distribution of high levels) and efficacy (by regression of unstable vessels). Based on recent advances in elucidating the molecular regulation of the switch between normal and aberrant angiogenesis and of vascular stabilization, as described above, it can be envisioned that local delivery of VEGF-expressing adenoviral vectors might be complemented by systemic treatment with drugs targeting ancillary pathways (eg, EphB4 or Sema3A signaling), to ensure a safe and effective outcome while exploiting the clinically attractive features of VEGF gene therapy.

ACKNOWLEDGMENTS

A.B. is supported by the Swiss National Science Foundation (182357), the European Union H2020 Program (646075 and 801159), the Swiss Nanoscience Institute (KOKORO), and intramural grants from the Department of Surgery of Basel University Hospital.

CONFLICT OF INTEREST

The authors declare no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

R. G.-B., N.D.M.: manuscript writing, figure preparation, final approval of manuscript; L.M., M.G.B., E.M.: manuscript writing, final approval of manuscript; L.G., D.J.S.: manuscript writing, financial support, final approval of manuscript; A.B.: conception and design, financial support, manuscript writing, final approval of manuscript.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Andrea Banfi  <https://orcid.org/0000-0001-5737-8811>

REFERENCES

- Marsano A, Medeiros da Cunha CM, Ghanaati S, et al. Spontaneous in vivo Chondrogenesis of bone marrow-derived mesenchymal progenitor cells by blocking vascular endothelial growth factor signaling. *STEM CELLS TRANSLATIONAL MEDICINE*. 2016;5:1730-1738.
- Benjamin EJ, Muntner P, Alonso A, et al. Heart disease and stroke statistics-2019 update: a report from the American Heart Association. *Circulation*. 2019;139:e56-e528.
- Everett E, Mathioudakis N. Update on management of diabetic foot ulcers. *Ann N Y Acad Sci*. 2018;1411:153-165.
- Rouwkema J, Rivron NC, van Blitterswijk CA. Vascularization in tissue engineering. *Trends Biotechnol*. 2008;26:434-441.
- Garcia JR, Garcia AJ. Biomaterial-mediated strategies targeting vascularization for bone repair. *Drug Deliv Transl Res*. 2016;6:77-95.
- Lopes D, Martins-Cruz C, Oliveira MB, Mano JF. Bone physiology as inspiration for tissue regenerative therapies. *Biomaterials*. 2018;185:240-275.
- Scheufler O, Schaefer DJ, Jaquiere C, et al. Spatial and temporal patterns of bone formation in ectopically pre-fabricated, autologous cell-based engineered bone flaps in rabbits. *J Cell Mol Med*. 2008;12:1238-1249.
- Senger DR, Galli SJ, Dvorak AM, Perruzzi C, Harvey V, Dvorak H. Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science*. 1983;219:983-985.
- Keck PJ, Hauser SD, Krivi G, et al. Vascular permeability factor, an endothelial cell mitogen related to PDGF. *Science*. 1989;246:1309-1312.
- Leung DW, Cachianes G, Kuang WJ, Goeddel D, Ferrara N. Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science*. 1989;246:1306-1309.
- Gupta R, Tongers J, Losordo DW. Human studies of angiogenic gene therapy. *Circ Res*. 2009;105:724-736.
- Carmeliet P, Jain RK. Molecular mechanisms and clinical applications of angiogenesis. *Nature*. 2011;473:298-307.
- Annex BH. Therapeutic angiogenesis for critical limb ischaemia. *Nat Rev Cardiol*. 2013;10:387-396.
- Korpisalo P, Yla-Herttuala S. Stimulation of functional vessel growth by gene therapy. *Integr Biol*. 2010;2:102-112.
- Tischer E, Mitchell R, Hartman T, et al. The human gene for vascular endothelial growth factor. Multiple protein forms are encoded through alternative exon splicing. *J Biol Chem*. 1991;266:11947-11954.
- Ferrara N. Binding to the extracellular matrix and proteolytic processing: two key mechanisms regulating vascular endothelial growth factor action. *Mol Biol Cell*. 2010;21:687-690.
- Lamproulou A, Ruhrberg C. Neuropilin regulation of angiogenesis. *Biochem Soc Trans*. 2014;42:1623-1628.
- Harper SJ, Bates DO. VEGF-A splicing: the key to anti-angiogenic therapeutics? *Nat Rev Cancer*. 2008;8:880-887.
- Ruhrberg C, Gerhardt H, Golding M, et al. Spatially restricted patterning cues provided by heparin-binding VEGF-A control blood vessel branching morphogenesis. *Genes Dev*. 2002;16:2684-2698.
- Gerhardt H, Golding M, Fruttiger M, et al. VEGF guides angiogenic sprouting utilizing endothelial tip cell filopodia. *J Cell Biol*. 2003;161:1163-1177.
- Betz C, Lenard A, Belting HG, Affolter M. Cell behaviors and dynamics during angiogenesis. *Development*. 2016;143:2249-2260.
- Fantini A, Vieira JM, Gestri G, et al. Tissue macrophages act as cellular chaperones for vascular anastomosis downstream of VEGF-mediated endothelial tip cell induction. *Blood*. 2010;116:829-840.
- Hellstrom M, Phng LK, Hofmann JJ, et al. Dll4 signalling through Notch1 regulates formation of tip cells during angiogenesis. *Nature*. 2007;445:776-780.
- Gianni-Barrera R, Bartolomeo M, Vollmar B, Djonov V, Banfi A. Split for the cure: VEGF, PDGF-BB and intussusception in therapeutic angiogenesis. *Biochem Soc Trans*. 2014;42:1637-1642.
- Egginton S, Zhou AL, Brown MD, Hudlická O. Unorthodox angiogenesis in skeletal muscle. *Cardiovasc Res*. 2001;49:634-646.
- Gianni-Barrera R, Trani M, Fontanellaz C, et al. VEGF over-expression in skeletal muscle induces angiogenesis by intussusception rather than sprouting. *Angiogenesis*. 2013;16:123-136.
- Al Haj Zen A, Oikawa A, Bazan-Peregrino M, et al. Inhibition of delta-like-4-mediated signaling impairs reparative angiogenesis after ischemia. *Circ Res*. 2010;107:283-293.
- Makanya AN, Hlushchuk R, Djonov VG. Intussusceptive angiogenesis and its role in vascular morphogenesis, patterning, and remodeling. *Angiogenesis*. 2009;12:113-123.
- Egginton S. Invited review: activity-induced angiogenesis. *Pflugers Arch*. 2009;457:963-977.
- De Spiegelaere W, Casteleyn C, Van den Broeck W, et al. Intussusceptive angiogenesis: a biologically relevant form of angiogenesis. *J Vasc Res*. 2012;49:390-404.
- Dimova I, Hlushchuk R, Makanya A, et al. Inhibition of notch signaling induces extensive intussusceptive neo-angiogenesis by recruitment of mononuclear cells. *Angiogenesis*. 2013;16:921-937.
- Dill MT, Rothweiler S, Djonov V, et al. Disruption of Notch1 induces vascular remodeling, intussusceptive angiogenesis, and angiosarcomas in livers of mice. *Gastroenterology*. 2012;142:967.e962-977.e962.
- Gianni-Barrera R, Butschkau A, Uccelli A, et al. PDGF-BB regulates splitting angiogenesis in skeletal muscle by limiting VEGF-induced endothelial proliferation. *Angiogenesis*. 2018;21:883-900.
- Groppa E, Brkic S, Uccelli A, et al. EphrinB2/EphB4 signaling regulates non-sprouting angiogenesis by VEGF. *EMBO Rep*. 2018;19:e45054.
- Bjarnegard M, Enge M, Norlin J, et al. Endothelium-specific ablation of PDGFB leads to pericyte loss and glomerular, cardiac and placental abnormalities. *Development*. 2004;131:1847-1857.
- Lindahl P, Johansson BR, Leveen P, et al. Pericyte loss and microaneurysm formation in PDGF-B-deficient mice. *Science*. 1997;277:242-245.
- Lindblom P, Gerhardt H, Liebner S, et al. Endothelial PDGF-B retention is required for proper investment of pericytes in the microvessel wall. *Genes Dev*. 2003;17:1835-1840.
- Armulik A, Genove G, Betsholtz C. Pericytes: developmental, physiological, and pathological perspectives, problems, and promises. *Dev Cell*. 2011;21:193-215.
- Sato Y, Rifkin DB. Inhibition of endothelial cell movement by pericytes and smooth muscle cells: activation of a latent transforming growth factor-beta 1-like molecule by plasmin during co-culture. *J Cell Biol*. 1989;109:309-315.
- Goumans MJ, Valdimarsdottir G, Itoh S, Rosendahl A, Sideras P, ten Dijke P. Balancing the activation state of the endothelium via two distinct TGF-beta type I receptors. *EMBO J*. 2002;21:1743-1753.
- van Meeteren LA, ten Dijke P. Regulation of endothelial cell plasticity by TGF-beta. *Cell Tissue Res*. 2012;347:177-186.
- Fagiani E, Christofori G. Angiopoietins in angiogenesis. *Cancer Lett*. 2013;328:18-26.
- Pasquale EB. Eph receptors and ephrins in cancer: bidirectional signaling and beyond. *Nat Rev Cancer*. 2010;10:165-180.
- Erber R, Eichelsbacher U, Powajob V, et al. EphB4 controls blood vascular morphogenesis during postnatal angiogenesis. *EMBO J*. 2006;25:628-641.
- Foo SS, Turner CJ, Adams S, et al. Ephrin-B2 controls cell motility and adhesion during blood-vessel-wall assembly. *Cell*. 2006;124:161-173.

46. Sawamiphak S, Seidel S, Essmann CL, et al. Ephrin-B2 regulates VEGFR2 function in developmental and tumour angiogenesis. *Nature*. 2010;465:487-491.
47. Wang Y, Nakayama M, Pitulescu ME, et al. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. *Nature*. 2010;465:483-486.
48. Murdoch C, Muthana M, Coffelt SB, Lewis CE. The role of myeloid cells in the promotion of tumour angiogenesis. *Nat Rev Cancer*. 2008;8:618-631.
49. Grunewald M, Avraham I, Dor Y, et al. VEGF-induced adult neovascularization: recruitment, retention, and role of accessory cells. *Cell*. 2006;124:175-189.
50. Dimova I, Karthik S, Makanya A, et al. SDF-1/CXCR4 signalling is involved in blood vessel growth and remodelling by intussusception. *J Cell Mol Med*. 2019;23:3916-3926.
51. De Palma M, Naldini L. Tie2-expressing monocytes (TEMs): novel targets and vehicles of anticancer therapy? *Biochim Biophys Acta*. 2009;1796:5-10.
52. Zacchigna S, Pattarini L, Zentilin L, et al. Bone marrow cells recruited through the neuropilin-1 receptor promote arterial formation at the sites of adult neoangiogenesis in mice. *J Clin Invest*. 2008;118:2062-2075.
53. Groppa E, Brkic S, Bovo E, et al. VEGF dose regulates vascular stabilization through Semaphorin3A and the Neuropilin-1+ monocyte/TGF-beta1 paracrine axis. *EMBO Mol Med*. 2015;7:1366-1384.
54. Lee RJ, Springer ML, Blanco-Bose WE, Shaw R, Ursell PC, Blau HM. VEGF gene delivery to myocardium: deleterious effects of unregulated expression. *Circulation*. 2000;102:898-901.
55. Schwarz ER, Speakman MT, Patterson M, et al. Evaluation of the effects of intramyocardial injection of DNA expressing vascular endothelial growth factor (VEGF) in a myocardial infarction model in the rat—angiogenesis and angioma formation. *J Am Coll Cardiol*. 2000;35:1323-1330.
56. Springer ML, Chen AS, Kraft PE, et al. VEGF gene delivery to muscle: potential role for vasculogenesis in adults. *Mol Cell*. 1998;2:549-558.
57. Sundberg C, Nagy JA, Brown LF, et al. Glomeruloid microvascular proliferation follows adenoviral vascular permeability factor/vascular endothelial growth factor-164 gene delivery. *Am J Pathol*. 2001;158:1145-1160.
58. Ozawa CR, Banfi A, Glazer NL, et al. Microenvironmental VEGF concentration, not total dose, determines a threshold between normal and aberrant angiogenesis. *J Clin Invest*. 2004;113:516-527.
59. von Degenfeld G, Banfi A, Springer ML, et al. Microenvironmental VEGF distribution is critical for stable and functional vessel growth in ischemia. *FASEB J*. 2006;20:2657-2659.
60. Boden J, Lassance-Soares RM, Wang H, et al. Vascular regeneration in ischemic Hindlimb by adeno-associated virus expressing conditionally silenced vascular endothelial growth factor. *J Am Heart Assoc*. 2016;5:e001815.
61. Ylä-Herttuala S, Baker AH. Cardiovascular gene therapy: past, present, and future. *Mol Ther*. 2017;25:1095-1106.
62. Misteli H, Wolff T, Fuglistaler P, et al. High-throughput flow cytometry purification of transduced progenitors expressing defined levels of vascular endothelial growth factor induces controlled angiogenesis in vivo. *STEM CELLS*. 2010;28:611-619.
63. Helmrich U, Marsano A, Melly L, et al. Generation of human adult mesenchymal stromal/stem cells expressing defined xenogenic vascular endothelial growth factor levels by optimized transduction and flow cytometry purification. *Tissue Eng Part C Methods*. 2012;18:283-292.
64. Wolff T, Mujagic E, Gianni-Barrera R, et al. FACS-purified myoblasts producing controlled VEGF levels induce safe and stable angiogenesis in chronic hind limb ischemia. *J Cell Mol Med*. 2012;16:107-117.
65. Melly LF, Marsano A, Frobert A, et al. Controlled angiogenesis in the heart by cell-based expression of specific vascular endothelial growth factor levels. *Hum Gene Ther Methods*. 2012;23:346-356.
66. Melly L, Cerino G, Frobert A, et al. Myocardial infarction stabilization by cell-based expression of controlled vascular endothelial growth factor levels. *J Cell Mol Med*. 2018;22:2580-2591.
67. Marsano A, Maidhof R, Luo J, et al. The effect of controlled expression of VEGF by transduced myoblasts in a cardiac patch on vascularization in a mouse model of myocardial infarction. *Biomaterials*. 2013;34:393-401.
68. Boccardo S, Gaudiello E, Melly L, et al. Engineered mesenchymal cell-based patches as controlled VEGF delivery systems to induce extrinsic angiogenesis. *Acta Biomater*. 2016;42:127-135.
69. Helmrich U, Di Maggio N, Guven S, et al. Osteogenic graft vascularization and bone resorption by VEGF-expressing human mesenchymal progenitors. *Biomaterials*. 2013;34:5025-5035.
70. Browne S, Pandit A. Engineered systems for therapeutic angiogenesis. *Curr Opin Pharmacol*. 2017;36:34-43.
71. Martino MM, Brkic S, Bovo E, et al. Extracellular matrix and growth factor engineering for controlled angiogenesis in regenerative medicine. *Front Bioeng Biotechnol*. 2015;3:45.
72. Addi C, Murschel F, De Crescenzo G. Design and use of chimeric proteins containing a collagen-binding domain for wound healing and bone regeneration. *Tissue Eng Part B Rev*. 2017;23:163-182.
73. Bao P, Kodra A, Tomic-Canic M, Golinko MS, Ehrlich HP, Brem H. The role of vascular endothelial growth factor in wound healing. *J Surg Res*. 2009;153:347-358.
74. Zisch AH, Schenk U, Schense JC, Sakiyama-Elbert SE, Hubbell JA. Covalently conjugated VEGF—fibrin matrices for endothelialization. *J Control Release*. 2001;72:101-113.
75. Sacchi V, Mittermayr R, Hartinger J, et al. Long-lasting fibrin matrices ensure stable and functional angiogenesis by highly tunable, sustained delivery of recombinant VEGF164. *Proc Natl Acad Sci USA*. 2014;111:6952-6957.
76. Martino MM, Tortelli F, Mochizuki M, et al. Engineering the growth factor microenvironment with fibronectin domains to promote wound and bone tissue healing. *Sci Transl Med*. 2011;3:100ra189.
77. Martino MM, Briquez PS, Guc E, et al. Growth factors engineered for super-affinity to the extracellular matrix enhance tissue healing. *Science*. 2014;343:885-888.
78. Li X, Tjwa M, Van Hove I, et al. Reevaluation of the role of VEGF-B suggests a restricted role in the revascularization of the ischemic myocardium. *Arterioscler Thromb Vasc Biol*. 2008;28:1614-1620.
79. Kivela R, Bry M, Robciuc MR, et al. VEGF-B-induced vascular growth leads to metabolic reprogramming and ischemia resistance in the heart. *EMBO Mol Med*. 2014;6:307-321.
80. Lahteenvuo JE, Lahteenvuo MT, Kivela A, et al. Vascular endothelial growth factor-B induces myocardium-specific angiogenesis and arteriogenesis via vascular endothelial growth factor receptor-1- and neuropilin receptor-1-dependent mechanisms. *Circulation*. 2009;119:845-856.
81. Rissanen TT, Markkanen JE, Gruchala M, et al. VEGF-D is the strongest angiogenic and lymphangiogenic effector among VEGFs delivered into skeletal muscle via adenoviruses. *Circ Res*. 2003;92:1098-1106.
82. Health USNIo. Adenovirus Vascular Endothelial Growth Factor D (AdvVEGF-D) Therapy for Treatment of Refractory Angina Pectoris (ReGenHeart). <https://clinicaltrials.gov/ct2/show/NCT03039751>. Accessed September 24, 2019.
83. Hartikainen J, Hassinen I, Hedman A, et al. Adenoviral intramyocardial VEGF-D/DeltaNDeltaC gene transfer increases myocardial perfusion reserve in refractory angina patients: a phase I/IIa study with 1-year follow-up. *Eur Heart J*. 2017;38:2547-2555.

84. Banfi A, von Degenfeld G, Gianni-Barrera R, et al. Therapeutic angiogenesis due to balanced single-vector delivery of VEGF and PDGF-BB. *FASEB J*. 2012;26:2486-2497.
85. Gianni-Barrera R, Burger M, Wolff T, et al. Long-term safety and stability of angiogenesis induced by balanced single-vector co-expression of PDGF-BB and VEGF164 in skeletal muscle. *Sci Rep*. 2016;6:21546.
86. Korpisalo P, Karvinen H, Rissanen TT, et al. Vascular endothelial growth factor-a and platelet-derived growth factor-B combination gene therapy prolongs angiogenic effects via recruitment of interstitial mononuclear cells and paracrine effects rather than improved pericyte coverage of angiogenic vessels. *Circ Res*. 2008;103:1092-1099.
87. Kupatt C, Hinkel R, Pfosser A, et al. Cotransfection of vascular endothelial growth factor-a and platelet-derived growth factor-B via recombinant adeno-associated virus resolves chronic ischemic malperfusion. Role of vessel maturation. *J Am Coll Cardiol*. 2010;56:414-422.
88. Richardson TP, Peters MC, Ennett AB, Mooney DJ. Polymeric system for dual growth factor delivery. *Nat Biotechnol*. 2001;19:1029-1034.
89. Thurston G, Suri C, Smith K, et al. Leakage-resistant blood vessels in mice transgenically overexpressing angiopoietin-1. *Science*. 1999;286:2511-2514.
90. Han S, Lee SJ, Kim KE, et al. Amelioration of sepsis by TIE2 activation-induced vascular protection. *Sci Transl Med*. 2016;8:335ra355.
91. Dor Y, Djonov V, Abramovitch R, et al. Conditional switching of VEGF provides new insights into adult neovascularization and pro-angiogenic therapy. *EMBO J*. 2002;21:1939-1947.
92. Tafuro S, Ayuso E, Zacchigna S, et al. Inducible adeno-associated virus vectors promote functional angiogenesis in adult organisms via regulated vascular endothelial growth factor expression. *Cardiovasc Res*. 2009;83:663-671.
93. Dai Y, Schwarz EM, Gu D, Zhang WW, Sarvetnick N, Verma IM. Cellular and humoral immune responses to adenoviral vectors containing factor IX gene: tolerization of factor IX and vector antigens allows for long-term expression. *Proc Natl Acad Sci USA*. 1995;92:1401-1405.
94. Forget A, Gianni-Barrera R, Uccelli A, et al. Mechanically defined microenvironment promotes stabilization of microvasculature, which correlates with the enrichment of a novel Piezo-1(+) population of circulating CD11b(+)/CD115(+) monocytes. *Adv Mater*. 2019;31:e1808050.
95. Bourguin PE, Gaudiello E, Pippenger B, et al. Engineered extracellular matrices as biomaterials of tunable composition and function. *Adv Funct Mater*. 2017;27:1605486.

How to cite this article: Gianni-Barrera R, Di Maggio N, Melly L, et al. Therapeutic vascularization in regenerative medicine. *STEM CELLS Transl Med*. 2020;9:433-444. <https://doi.org/10.1002/sctm.19-0319>