



Review Article

Membraneless organelles: phasing out of equilibrium

Author(s):

Hondele, Maria; Heinrich, Stephanie; De Los Rios, Paolo; Weis, Karsten

Publication Date:

2020-12

Permanent Link:

<https://doi.org/10.3929/ethz-b-000454411> →

Originally published in:

Emerging Topics in Life Sciences 4(3), <http://doi.org/10.1042/ETLS20190190> →

Rights / License:

[In Copyright - Non-Commercial Use Permitted](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).

Membraneless organelles: phasing out of equilibrium

Authors/Affiliations:

Maria Hondele^{1*}, Stephanie Heinrich¹, Paolo De Los Rios², Karsten Weis^{1*}

¹ - Institute of Biochemistry, Department of Biology, ETH Zurich, Zurich, CH-8093, Switzerland

² - Institute of Physics, School of Basic Sciences, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

*Correspondence: maria.hondele@bc.biol.ethz.ch and karsten.weis@bc.biol.ethz.ch

ABSTRACT

Over the past years, liquid-liquid phase separation has emerged as a ubiquitous principle of cellular organisation implicated in many biological processes ranging from gene expression to cell division. The formation of biological condensates, like the nucleolus or stress granules, by liquid-liquid phase separation is at its core a thermodynamic equilibrium process. However, life does not operate at equilibrium, and cells have evolved multiple strategies to keep condensates in a non-equilibrium state. In this review, we discuss how these non-equilibrium drivers counteract solidification and potentially detrimental aggregation, and at the same time enable biological condensates to perform work and control the flux of substrates and information in a spatial and temporal manner.

Cells are amazing mini-reactors, crowded with DNA, RNA, proteins, lipids and metabolites. To avoid chaos and to make intracellular reactions efficient, cells have evolved several strategies to compartmentalise and organize their content. In eukaryotic cells, intracellular membranes form specialised organelles that separate, for example, transcription in the nucleus from translation in the cytosol, or sequester respiratory chain enzymes in mitochondria. In addition, the cytoskeleton provides 'highways' that allow for the directed transport of, e.g., RNA or vesicles to the distal end of cells, and positions membrane-bounded organelles such as the endoplasmic reticulum.

In recent years, it has become clear that there is yet another organising principle: the formation of membraneless organelles by liquid-liquid phase separation (LLPS) (1,2). In brief, LLPS describes the phenomenon that some biomolecules, such as proteins or nucleic acids, undergo dynamic self-association to form condensates within the surrounding cyto- or nucleoplasm. Examples include the nucleolus, nuclear speckles, stress granules or centrosomes. The role of LLPS in organising cellular content is ubiquitous, and critical for many biological processes in all cells, from bacteria to humans. This suggests that the process of biomolecular condensation is evolutionary ancient, and that phase-separated structures might have organized cellular components since the beginning of life (3-5).

LLPS-competent molecules have a certain solubility limit, and if this limit is exceeded, they reversibly condensate. Such behaviour is favoured because separating the cellular milieu into two regions, one of high component concentration and one diluted, minimizes the free energy of the system (6,7). In this respect, LLPS is, at its core, a thermodynamic equilibrium phenomenon. Yet, at equilibrium, the second principle of thermodynamics forbids any form of activity, such as directional molecular processing. Thus, in the absence of external energy input, condensates cannot perform any cellular work. In fact, they might even pose a danger since LLPS-mediated structures are prone to form irreversible, potentially detrimental aggregates (8). However, life does not operate at equilibrium and energy-consuming processes have evolved that keep condensates away from equilibrium. In this review, we discuss how cells exploit such non-equilibrium drivers not only as a protective mechanism but also to perform work, e.g., to regulate the flux of substrates and information, or to control biochemical reactions in a spatial or temporal manner. Thus, these non-equilibrium processes endow condensates with biological functions beyond a mere role in organizing a cell's content.

Physical principles of condensation

Interactions between molecules, e.g., between proteins, or proteins and nucleic acids, are characterised by the affinity or 'strength' of individual interactions, and the 'valency', which describes the number of interactions one molecule can undergo with its partners. The formation of many biomolecular condensates by LLPS is based on weak, multivalent interactions: components need to have a valency of at least three to establish a branched, interconnected meshwork (Figure 1) (9,10), and interactions generally have to be weak (in the high nanomolar to low micromolar range) to keep the condensates readily reversible and liquid-like. Interactions can involve globular domains or unstructured protein regions harbouring low amino acid sequence complexity (e.g., unusually rich in glutamine, asparagine, glycine, serine etc.), blocks of

opposite charge, or cation- π interactions (between aromatic amino acids such as tyrosine and cations like arginine) (1,11). Theoretically, condensates can also form at valencies below three, for example when entropy favours dense packing of large, inert entities accompanied by the un-packing of smaller molecules, a phenomenon referred to as 'depletion interaction' (12). However, since depletion interactions of colloidal particles are not specific, it is unclear whether they play a significant role in biology.

Condensates assemble when the concentration of at least one of their key components crosses its critical saturation limit. The result is the formation of liquid-like 'droplets' where the components can concentrate over their surrounding environment by several orders of magnitude. Following the law of mass action, an "initial equilibrium" that partitions components within the droplet and in the soluble phase outside the droplet is established rapidly, until the soluble pool is again below the solubility limit (7,13). Once an equilibrium is reached, there is no longer any net flux of components into or out of the droplet, although – due to the weak, short-lived nature of the interactions - passive exchange of components between the droplet phase and the soluble phase can still occur.

This "initial equilibrium" state persists if all components keep their original conformations and interactions. However, most biological LLPS systems display additional 'ripening' behaviours (Figure 2): after the two liquid phases are phase-separated, often nucleated by a multitude of sites, the system still tries to minimize the interface between phases, and thus the free energy associated with the surface tension. To do so, smaller droplets can either fuse, or undergo a process called 'Ostwald ripening' where smaller droplets dissolve and their components associate to larger ones until, ideally, a single droplet remains (4,14). Another ripening behaviour is a kinetically slow solidification process that turns liquid-like droplets over time into irreversible aggregates (15,16). While the structural principles underlying this droplet aging remain poorly understood, one possibility is that in the initial liquid-like state the condensate components are not yet in an optimal sterical conformation and only a fraction of all possible inter-molecular interactions are formed. Over time, aligning - or deforming - them into more favourable conformations could then promote the formation of additional and stronger interactions (17). We are only beginning to understand what protein features (18) or factors in the cellular environment could promote these transformations, however, an increase in the percentage of occupied interaction sites would lead to a solidification, which confers rigidity to the denser meshwork limiting exchange of its components. A "final equilibrium" - or minimal free-energy state - is reached when a maximum of interactions is established without generating strain. Solidification has been observed in several biological systems, including centrosomes (19,20), germ granules (21,22), enhancer condensates (23) and heterochromatin (24). For some of these systems, a gel-like material state seems to be beneficial and presumably supports biological function by providing mechanical stability, by increasing component retention times (25) or as a means of cellular adaptation (26). But in many cases the benefit of solidification is unclear, and the process might be rather detrimental. In this light, the aging behaviour of phase-separated condensates has gained a lot of attention as it might underly the aggregation phenomena seen in a multitude of age-dependent aggregation diseases, including a large group of neurodegenerative diseases (8,27,28). For example, alpha-synuclein

(29) or TDP-43 (30) and FUS (31), critical drivers of Parkinson's disease or amyotrophic lateral sclerosis (ALS), respectively, can undergo LLPS and over time transform into irreversible aggregates *in vitro*.

Biological condensates are heterogenous

Biological condensates, or membraneless organelles, are built from different types of macromolecules, and contain a large number of distinct components. For example, the stress granule proteome was estimated to consist of at least several hundred different proteins and a large number of distinct RNAs (32-35). Among them, 'primary' components (also called 'scaffolds') have the ability to undergo phase separation whereas 'secondary' factors (or 'clients') cannot phase-separate by themselves, but nonetheless can be selectively recruited to condensates via interaction with the primary components, and in consequence enrich significantly in the organelle (2,9). Given the compositional complexity of membraneless organelles such a classification is obviously a simplification. Secondary factors can regulate, enhance or decrease phase separation and thereby dramatically influence the overall condensation result (6,36-38). Importantly, the heterogeneity of condensates can also lead to non-intuitive differences in the behaviour of their constituents. For example, both primary and secondary components can have very different exchange rates, ranging from rapid exchange and liquid-like behaviour to very slow turnover, as if they were 'solid' components of a given organelle (38-40). Likely these behaviours depend on the strength and valency of interactions of a given component with its partners, but potentially also the size and diffusional ability of a component within the LLPS meshwork.

Differences in biochemical component properties can also lead to further un-mixing and formation of 'droplets within droplets'. This can induce the establishment of distinct sub-compartments with different material and thermodynamic characteristics. Gel-like or even fibrillar cores are, for instance, found in the nucleolus (41) or in stress granules (32) and surrounded by more liquid compartments. The inverted situation is found in P-granules, prominent RNA-containing organelles present during development where a shell of gel-like MEG-3 granules forms around, and thereby locally stabilizes a liquid core of phase-separated PGL-3, which by itself is intrinsically labile (39).

Regulation of biological condensates

The formation of LLPS condensates is critically regulated by the concentration of its constituents. In a biological context, this is exemplified by P-bodies and stress granules, whose formation depends on the availability of non-translated, ribosome-free mRNA (42). However, changing protein or RNA levels within a cell, either by new synthesis or degradation, is slow and energetically costly. It is therefore of importance that besides changing the component concentration, condensation can also be controlled by changing the strength or valency of interactions, or more generally, by shifting the critical saturation threshold. In cells, this can be triggered by changes in temperature, crowding and also pH, which can occur during heat shock, nutrient starvation or changes in growth state. These altered conditions can directly induce changes in protein conformation to expose interaction sites and to promote the formation of

condensates (43-45). Thus, in particular stress situations the altered cellular environment is 'sensed' to shift the condensation threshold and propensity -without altering component concentrations- in order to respond and survive stress situations (46,47).

Furthermore, several energy-consuming, enzymatic processes modulate condensation. Post-translational modifications can change the physical and chemical properties of amino-acids, and thus have the potential to alter interaction strengths (48-51). For example, arginine methylation and phosphorylation systems have been described that prevent or promote condensation, or acutely dissolve existing condensates, thus acting as on-off switches for membraneless organelles. This includes protein kinase A (PKA), which prevents formation of P-bodies when nutrients are abundant by directly phosphorylating the P-body regulator Pat1 (37,52), and the DYRK3 kinase triggers the disassembly of several membraneless organelles during mitosis (53,54). In addition, post-translational modification could also be used to fine-tune cellular condensates, and recruitment of kinases was postulated to control and limit the size of individual condensates (6).

Another critical set of regulators of cellular condensation include protein chaperones and RNA helicases. These enzymes are prominently associated with different membraneless organelles, in particular larger condensates like the nucleolus or stress granules (32,55,56). In general, the main task of these enzymes is to constantly remodel protein-protein, protein-RNA and RNA-RNA interactions, and to shield exposed surfaces (57-60). Heat Shock Proteins such as Hsp70, Hsp90 and Hsp60 (GroEL in bacteria, CCT in eukaryotes) remodel proteins through their ATPase cycle and thus promote folding or disaggregation of condensate components (61,62). Thus, they can ultimately change interaction patterns and efficiently modulate the critical concentration threshold that is required for condensation. As a consequence, members of these protein families were shown to be important regulators of both assembly and disassembly of membraneless organelles. In addition, they can keep condensates in a liquid and reversible state (32,60,62-66), and chaperone malfunction, for example during ageing, has been linked to aggregation diseases and neurodegeneration (28,67).

How do condensates at equilibrium and non-equilibrium execute function beyond mere assembly?

Phase separation has now been intimately linked to many important cellular processes, but the holy grail remains to understand the function(s) of biological condensates: do they merely act as glue, sticking together molecules in order to build larger assemblies, or can they accomplish more than that?

In the following, we discuss several types of condensate functions (Figure 3). Some of them, like the spatially localized enrichment of macromolecules in a membraneless organelle, could -in principle- occur at thermodynamic equilibrium. At thermodynamic equilibrium, components can continuously exchange between condensates and the soluble phase, yet by definition, the net fluxes remain zero. However, as discussed, cells employ a whole set of energy-consuming enzymes that keep condensates away from

reaching equilibrium (7,68). For one thing, this can regulate condensate dynamics or prevent the ripening or solidification of condensates avoiding that their components enter into a non-reactive state with no exchange of molecules with the surrounding medium.

However, energy input procures more than the regulation of condensate assembly or reversibility. The injection of energy into condensates by enzymatic processes offsets condensates from equilibrium, yet, at the same time, condensates steadily try to return to their equilibrium state. A crucial consequence is that condensates now exist in a *non-equilibrium steady-state* that is intrinsically different from an equilibrium one: it can generate actual net fluxes, which can be exploited to drive downstream processes. Thus, energy-fuelled condensates can perform biochemical work, transmit directional information, or even power the translocation of macromolecules in a spatial and temporal manner.

Buffering of concentration changes

In a simple model, phase-separating molecules have a concentration-dependent solubility limit. Condensates form if the concentration exceeds this solubility limit, and grow until the concentration of the soluble pool is sufficiently reduced (4,7). If the total concentration of a component further increases, a new equilibrium is rapidly reached yet the concentration of the soluble phase remains constant. It is therefore 'buffered' against concentration changes. In consequence, cells could use phase separation to dampen concentration fluctuations and keep the concentration of soluble components constant. However, it has been recently demonstrated that biological, multi-component systems are more complex, and that saturation concentrations and partitioning coefficients depend on the cellular context and interaction networks (36,69,70). Nevertheless, buffering by LLPS could counteract variability or stochasticity in various steps of gene expression reducing the 'noise' in cellular protein levels. Indeed -and despite the fact that phase separation processes in cells are generally kept away from equilibrium-, it was recently demonstrated that membraneless compartments can effectively reduce protein concentration noise in a living system (71).

Concentration hubs as highly selective reaction centres

Once formed, condensates act as concentration hubs that sequester both the phase-separating component and other interacting proteins or nucleic acids, away from a homogenous distribution. However, condensates are biochemically selective: they enrich for some proteins while excluding others (19,36,72,73). This has major biological consequences: first, concentrating enzymatic activities and their substrates can influence their enzymatic activity that, by the law of mass action, depends on the concentrations of both enzyme and substrate (74-80).

Second, formation of condensates can control protein or RNA localization in a spatial and temporal manner. As an example, for the diverse set of RNA condensates this could mean that if a transcript enters a specific condensate, it will be selectively modified by enzymes or RNA-binding proteins enriched therein. Condensate formation can therefore speed up and alter RNA modification rates, RNA unfolding, and the remodeling of mRNA-protein (mRNP) complexes. Conversely, selective sequestration into condensates

can also prevent that enzymes and substrates see each other and thus slow down or even completely inhibit enzymatic reactions (81). For instance, mRNAs could be hidden or stored away in cytoplasmic RNP granules preventing their access to the translation machinery (82-84). However, given the constant exchange between the phase-separated condensate and the soluble phase at steady state, complete inhibition will require low exchange rates and high partition coefficients in a condensate.

Phase separation is further important for the formation and function of super-enhancers, where highly dynamic nano-condensates of transcription factors and the mediator complex coalesce on specific DNA sequence elements, and subsequently sequester the transcription machinery to promote transcription (85-92). Thus, phase separation on super enhancers provides an elegant explanation for the highly co-operative processes that are required to bring together many protein factors in order to allow for a highly selective transcriptional activation of developmentally important enhancer and promoter DNA elements.

Signalling cascades and gradients

Formation of condensates requires multivalent interactions at component concentrations above a certain threshold. If either the concentration or the interaction strength of primary condensate components is actively and selectively enhanced or diminished, e.g., by post-translational modifications such as phosphorylation, condensates can form or disassemble in a spatially and temporally controlled manner (7,14,68). This can result in spatial condensate gradients or other non-homogenous distributions of condensates that however need to be actively maintained and replenished by energy-consuming processes (93). Cells have adopted this concept to amplify or transmit information.

For example, P-granules in *C. elegans* are not evenly distributed, and they concentrate at the posterior end of the one-cell embryo, which is essential for proper embryonic development. This spatial gradient is achieved by the permanent, phosphorylation-dependent granule dissolution occurring at the anterior pole of the embryo (94-96). Another example is the proposed channelling of nascent mRNA transcripts through a series of condensates formed from transcription initiation or elongation factors, and RNA polymerase II itself. Here, changes in the phosphorylation pattern of RNA polymerase II promote transfer between initiation and elongation condensates, which ultimately enhances and drives directionality of the process (97-99).

Like other concentration hubs, such actively induced condensates can have emergent properties as exemplified by cellular signaling cascades: local phosphorylation-induced phase separation of upstream signalling factors enriches the downstream signaling components beyond a critical threshold to form condensates initiating a signaling cascade (100). Examples include PAR-mediated clustering of factors for early DNA damage response (31,101,102), signaling factor clustering to create asymmetric daughter cells in bacteria (103), clustering of T-cell receptors for nucleation of actin polymerisation (104), or condensation

of centrosome components for microtubule nucleation (19,105). The observed stimulation of signaling activity in such hubs can exceed what is expected merely based on the increase in local concentration, probably by prolonging residence and interaction times in condensates (106).

ATPases drive and orchestrate substrate flux through membraneless organelles

While the flow of information is rather abstract, condensates can also orchestrate the directional flux of actual molecules if they receive input from energy-consuming enzymes. For example, mRNAs are sent through a series of condensates, from transcription and splicing to export through the nuclear pore, and ultimately storage or decay in P-bodies and stress granules (107-109). Along their life, mRNA molecules are chaperoned by diverse proteins. In particular, the family of DEAD-box ATPases (DDXs) is not only a chaperone or passive companion, but rather acts as a global regulator of mRNA flux through these condensates. Many DDXs possess low complexity sequences that allow them to undergo phase-separation in the ATP- and RNA-bound form, which results in RNA accumulation in the condensate. DDX ATPase activity triggers condensate dissolution and release of the RNA substrate, that can then transit to another granule. In addition, DDXs can control condensates and RNA accumulation *in trans* by remodeling phase-separated structures (56,66). Mutations that abrogate DDX ATPase activity prevent release of the RNA, and thus halt or diminish the flux of RNA, resulting in increased or even ectopic RNA accumulation and condensate formation (56,63,83,110-112). Since mRNAs and mRNPs can be specifically modified by condensate-enriched enzymes or RNA-binding proteins, the mRNA particle leaving a condensate will be distinct from the one that entered. Such active processes keep the condensates away from equilibrium and can generate directional and irreversible fluxes. Thus, energy-consuming enzymes like DDXs not only regulate condensate formation and turnover – they have the potential to orchestrate and regulate the flux of genetic information in a spatial and temporal manner.

Concluding remarks

Slowly, we are beginning to realize how pervasively phase separation is hard-coded in the cellular proteome. A large number of proteins - up to 75%, depending on the species - contain intrinsically disordered sequences. While not all intrinsically disordered protein domains function in condensation, many of them contain sequences of low amino acid complexity and thus have the potential to promote liquid-liquid phase separation. Equally surprisingly, cells keep the majority of their proteins slightly above their intrinsic solubility threshold (16,113). Cells thus operate at the solubility limit, which could explain why upon energy depletion or during stress, the cytoplasm can convert from a fluid state to a more gel-like or solid state (114-117). This transformation is likely influenced by phase separation of some components of the proteome carrying aggregation-prone sequences that are normally kept soluble by energy-consuming processes. While this cytoplasmic transformation might be important for cells to enter a quiescent state or to remain dormant it also poses a potential risk, since phase-separated condensates can mature over time into irreversible, potentially toxic aggregates akin to aggregates observed in age-dependent aggregation diseases. For example, *in vitro* reconstituted condensates of purified proteins like FUS, implicated

in ALS, quickly 'age' into more solid, aggregated forms, whereas FUS solidification is not observed in young and healthy cells (15,25,31).

The implications are two-fold: first, given their potentially detriment, phase separations must provide huge benefits to the cell, otherwise the bulk of low complexity sequences would have been lost over evolution. And second, kinetically unstable condensates pose a major challenge, and cells constantly counteract spontaneous condensation, and once formed, prevent condensate-maturation and ageing to avoid conversion into irreversible aggregates. To keep these processes in check, evolution has created energy-consuming solutions and machineries.

Biomolecular condensates are found in both pro- and eukaryotes, and the cellular usage of LLPS as an organizational principle is thus likely evolutionarily ancient. It is tempting to speculate that molecular condensates appeared even before the beginning of life to allow for the selective concentration of specific macromolecules in the primordial soup (3-5). Turning such condensates into a living system would then necessitate the addition of non-equilibrium drivers, that could utilize free energy from the environment to create ordered structures and ultimately evolve an ability to self-replicate. In this regard, the nucleolus might be the closest remnant of such an ancestral condensate efficiently producing ribosomes around the genetic material coding for ribosomal RNA, which in an early RNA world might have acquired the ability to self-replicate even prior to the evolution of membrane-bounded cells.

Throughout the tree of life biomolecular condensates are widely used in a large number of biological processes, probably because they are an elegant combination of efficiency and flexibility. Interactions within condensates often rely on short, unstructured domains, that unlike complex three-dimensional protein folds can easily change and evolve. Thus, biological condensates might have the ability to readily add new factors or enzymes to their inventory and thereby attain new functionality. Furthermore, membrane-less organelles, in contrast to their membrane-enclosed counterparts, do not need to evolve complex transport machineries to take up new components and they are highly dynamic and can form *de novo*. This enormous flexibility might outweigh the constant energetic demands to keep condensates away from equilibrium and to prevent potential deleterious aggregation. Their dynamics and ease of regulation makes them also particularly suitable to rapidly respond to changes in the environment and to orchestrate changes in metabolism or gene expression during times of stress. Why would cells otherwise want to keep some sticky glue in their belly?

SUMMARY POINTS

1. Liquid-liquid phase separation is an equilibrium phenomenon which excludes a net flux of macromolecules between condensates and the surrounding environment.
2. Reaching the lowest free energy at equilibrium is potentially detrimental since it can lead to solidification and aggregation of condensates.
3. Cells have evolved energy-consuming mechanisms to keep condensates in a non-equilibrium steady-state. These mechanisms include phosphorylation cycles, protein chaperones and RNA helicases.
4. Beyond keeping condensates in a liquid-like state, non-equilibrium drivers enable condensates to perform work. Cells make use of this to control biochemical reactions and the flux of substrates and information in a spatial and temporal manner.

ACKNOWLEDGEMENTS

This work was supported by a grant from the Swiss National Fond (SNF 31003A_159731) to KW. The authors thank Vamshidhar Gade and Janka Zsok for critical reading of the manuscript.

AUTHOR CONTRIBUTIONS

M.H., K.W., and P.R. wrote the manuscript, with critical input from S.H.. S.H. designed and created the figures.

DECLARATION OF INTERESTS

The authors declare no conflict of interests.

FIGURE LEGENDS

Figure 1: Valency: molecules with at least three interactions can form phase-separated networks.

Figure 2: Liquid-liquid phase separation is a thermodynamic equilibrium process. In cells, energy input keeps phase separated condensates away from equilibrium, which enables them to perform work and prevents their potentially detrimental solidification.

Figure 3: Examples for equilibrium and non-equilibrium functions of biological condensates. For details, see text.

REFERENCES

1. Shin Y, Brangwynne CP. Liquid phase condensation in cell physiology and disease. *Science*. American Association for the Advancement of Science; 2017 Sep 21;357(6357):eaaf4382–13.
2. Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of cellular biochemistry. *Nature Research*; 2017 Feb 22.
3. Hansma HG. Better than Membranes at the Origin of Life? *Life (Basel)*. Multidisciplinary Digital Publishing Institute; 2017 Jun 20;7(2):28.
4. Hyman AA, Weber CA, Jülicher F. Liquid-liquid phase separation in biology. *Annu Rev Cell Dev Biol*. Annual Reviews; 2014;30(1):39–58.
5. Yoshizawa T, Nozawa R-S, Jia TZ, Saio T, Mori E. Biological phase separation: cell biology meets biophysics. *Biophys Rev*. Springer Berlin Heidelberg; 2020 Mar 18;5(80):495–21.
6. Söding J, Zwicker D, Sohrabi-Jahromi S, Boehning M, Kirschbaum J. Mechanisms for Active Regulation of Biomolecular Condensates. *Trends Cell Biol*. 2020 Jan;30(1):4–14.
7. Berry J, Brangwynne C, Haataja MP. Physical Principles of Intracellular Organization via Active and Passive Phase Transitions. *Rep Prog Phys*. IOP Publishing; 2018 Jan 9.
8. Alberti S, Dormann D. Liquid-Liquid Phase Separation in Disease. *Annu Rev Genet*. Annual Reviews; 2019 Aug 20;53(1):annurev-genet-112618-043527-194.
9. Banani SF, Rice AM, Peeples WB, Lin Y, Jain S, Parker R, et al. Compositional Control of Phase-Separated Cellular Bodies. *Cell*. 2016 Jun 29.
10. Sanders DW, Kedersha N, Lee DSW, Strom AR, Drake V, Riback JA, et al. Competing Protein-RNA Interaction Networks Control Multiphase Intracellular Organization. *Cell*. Elsevier; 2020 Apr 16;181(2):306–28.
11. Boeynaems S, Alberti S, Fawzi NL, Mittag T, Polymenidou M, Rousseau F, et al. Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol*. 2018 Jun;28(6):420–35.
12. Tuinier R., Vliegthart GA., Lekkerkerker HNW. Depletion interaction between spheres immersed in a solution of ideal polymer chains. *J. Chem. Phys*. 2000 Dec 15;113 (23):10768-75.
13. Alberti S. Phase separation in biology. *Curr Biol*. 2017 Oct 23;27(20):R1097–102.
14. Weber CA, Lee CF, Jülicher F. Droplet ripening in concentration gradients. *New Journal of Physics*. IOP Publishing; 2017 May 14;19(5):0–0.
15. Alberti S. The wisdom of crowds: regulating cell function through condensed states of living matter. *J Cell Sci*. 2017 Sep 1;130(17):2789–96.
16. Halfmann R. A glass menagerie of low complexity sequences. *Curr Opin Struct Biol*. 2016 Jun;38:18–25.
17. Kato M, McKnight SL. Cross- β Polymerization of Low Complexity Sequence Domains. *Cold Spring Harb Perspect Biol*. 2017 Mar 1;9(3):a023598.
18. Martin EW, Holehouse AS, Peran I, Farag M, Incicco JJ, Bremer A, et al. Valence and patterning of aromatic residues determine the phase behavior of prion-like domains. *Science*. American Association for the Advancement of Science; 2020 Feb 7;367(6478):694–9.

19. Woodruff JB, Ferreira Gomes B, Widlund PO, Mahamid J, Honigsmann A, Hyman AA. The Centrosome Is a Selective Condensate that Nucleates Microtubules by Concentrating Tubulin. *Cell*. 2017 Jun 1;169(6):1066–1077.e10.
20. Mittasch M, Tran VM, Rios MU, Fritsch AW, Enos SJ, Ferreira Gomes B, et al. Regulated changes in material properties underlie centrosome disassembly during mitotic exit. *J Cell Biol*. 2020 Apr 6;219(4):647.
21. Kistler KE, Trcek T, Hurd TR, Chen R, Liang F-X, Sall J, et al. Phase transitioned nuclear Oskar promotes cell division of Drosophila primordial germ cells. *eLife*. eLife Sciences Publications Limited; 2018 Sep 27;7:65.
22. Krishnakumar P, Riemer S, Perera R, Lingner T, Goloborodko A, Khalifa H, et al. Functional equivalence of germ plasm organizers. Moens C, editor. *PLoS Genet*. Public Library of Science; 2018 Nov;14(11):e1007696.
23. Nair SJ, Yang L, Meluzzi D, Oh S, Yang F, Friedman MJ, et al. Phase separation of ligand-activated enhancers licenses cooperative chromosomal enhancer assembly. *Nat Struct Mol Biol*. Nature Publishing Group; 2019 Mar;26(3):193–203.
24. Strom AR, Emelyanov AV, Mir M, Fyodorov DV, Darzacq X, Karpen GH. Phase separation drives heterochromatin domain formation. Nature Publishing Group. Nature Publishing Group; 2017 Jun 21;:1–10.
25. Woodruff JB, Hyman AA, Boke E. Organization and Function of Non-dynamic Biomolecular Condensates. *Trends in Biochemical Sciences*. 2018 Feb;43(2):81–94.
26. Saarikangas J, Barral Y. Protein aggregation as a mechanism of adaptive cellular responses. *Curr Genet*. Springer Berlin Heidelberg; 2016 Nov;62(4):711–24.
27. Aguzzi A, Altmeyer M. Phase Separation: Linking Cellular Compartmentalization to Disease. *Trends Cell Biol*. 2016 Jul;26(7):547–58.
28. Alberti S, Hyman AA. Are aberrant phase transitions a driver of cellular aging? *Bioessays*. 2016 Oct;38(10):959–68.
29. Ray S, Singh N, Pandey S, Kumar R, Gadhe L, Datta D, et al. Liquid-liquid phase separation and liquid-to-solid transition mediate α -synuclein amyloid fibril containing hydrogel formation. *bioRxiv*. Cold Spring Harbor Laboratory; 2019 Apr 26;20:619858.
30. Conicella AE, Zerbe GH, Mittal J, Fawzi NL. ALS Mutations Disrupt Phase Separation Mediated by α -Helical Structure in the TDP-43 Low-Complexity C-Terminal Domain. *Structure*. 2016 Sep 6;24(9):1537–49.
31. Patel A, Lee HO, Jawerth L, Maharana S, Jahnel M, Hein MY, et al. A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease Mutation. *Cell*. 2015 Aug 27;162(5):1066–77.
32. Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A, Parker R. ATPase-Modulated Stress Granules Contain a Diverse Proteome and Substructure. *Cell*. Elsevier Inc; 2016 Jan 19;:1–35.
33. Khong A, Jain S, Matheny T, Wheeler JR, Parker R. Isolation of mammalian stress granule cores for RNA-Seq analysis. *Methods*. 2018 Mar 15;137:49–54.
34. Fazal FM, Han S, Parker KR, Kaewsapsak P, Xu J, Boettiger AN, et al. Atlas of Subcellular RNA Localization Revealed by APEX-Seq. *Cell*. 2019 Jul 11;178(2):473–490.e26.
35. Padròn A, Iwasaki S, Ingolia NT. Proximity RNA labeling by APEX-Seq Reveals the Organization of Translation Initiation Complexes and Repressive RNA Granules. *bioRxiv*. Cold Spring Harbor Laboratory; 2018 Oct 25;15:454066.

36. Riback JA, Zhu L, Ferrolino MC, Tolbert M, Mitrea DM, Sanders DW, et al. Composition-dependent thermodynamics of intracellular phase separation. *Nature Publishing Group. Springer US*; 2020 Apr 30;:1–17.
37. Sachdev R, Hondele M, Linsenmeier M, Vallotton P, Mugler CF, Arosio P, et al. Pat1 promotes processing body assembly by enhancing the phase separation of the DEAD-box ATPase Dhh1 and RNA. *eLife. eLife Sciences Publications Limited*; 2019 Jan 16;8.
38. Mugler CF, Hondele M, Heinrich S, Sachdev R, Vallotton P, Koek AY, et al. ATPase activity of the DEAD-box protein Dhh1 controls processing body formation. *eLife. eLife Sciences Publications Limited*; 2016 Oct 3;5:4154.
39. Putnam A, Cassani M, Smith J, Seydoux G. A gel phase promotes condensation of liquid P granules in *Caenorhabditis elegans* embryos. *Nat Struct Mol Biol. Nature Publishing Group*; 2019 Mar;26(3):220–6.
40. Taylor NO, Wei M-T, Stone HA, Brangwynne CP. Quantifying Dynamics in Phase-Separated Condensates Using Fluorescence Recovery after Photobleaching. *Biophysj. Cell Press*; 2019 Oct 1;117(7):1285–300.
41. Feric M, Vaidya N, Harmon TS, Mitrea DM, Zhu L, Richardson TM, et al. Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell. 2016 Jun 16*;165(7):1686–97.
42. Balagopal V, Parker R. Polysomes, P bodies and stress granules: states and fates of eukaryotic mRNAs. *Curr Opin Cell Biol. 2009 Jun*;21(3):403–8.
43. Riback JA, Katanski CD, Kear-Scott JL, Pilipenko EV, Rojek AE, Sosnick TR, et al. Stress-Triggered Phase Separation Is an Adaptive, Evolutionarily Tuned Response. *Cell. 2017 Mar 9*;168(6):1028–1040.e19.
44. Iserman C, Altamirano CD, Jegers C, Friedrich U, Zarin T, Fritsch AW, et al. Condensation of Ded1p Promotes a Translational Switch from Housekeeping to Stress Protein Production. *Cell. Elsevier Inc*; 2020 May 1;:1–34.
45. Franzmann TM, Jahnel M, Pozniakovskiy A, Mahamid J, Holehouse AS, Nüske E, et al. Phase separation of a yeast prion protein promotes cellular fitness. *Science. 2018 Jan 5*;359(6371).
46. Franzmann TM, Alberti S. Protein Phase Separation as a Stress Survival Strategy. *Cold Spring Harb Perspect Biol. Cold Spring Harbor Lab*; 2019 Jan 7;:a034058.
47. Yoo H, Triandafillou C, Drummond DA. Cellular sensing by phase separation: Using the process, not just the products. *J Biol Chem. American Society for Biochemistry and Molecular Biology*; 2019 May 3;294(18):7151–9.
48. Berchtold D, Battich N, Pelkmans L. A Systems-Level Study Reveals Regulators of Membrane-less Organelles in Human Cells. *Mol Cell. 2018 Dec 20*;72(6):1035–5.
49. Snead WT, Gladfelter AS. The Control Centers of Biomolecular Phase Separation: How Membrane Surfaces, PTMs, and Active Processes Regulate Condensation. *Mol Cell. 2019 Oct 17*;76(2):295–305.
50. Bratek Skicki A, Pancsa R, Meszaros B, Van Lindt J, Tompa P. A guide to regulation of the formation of biomolecular condensates. *FEBS J. John Wiley & Sons, Ltd*; 2020 Mar 14;48:D360–12.
51. Hofweber M, Dormann D. Friend or foe-Post-translational modifications as regulators of phase separation and RNP granule dynamics. *J Biol Chem. American Society for Biochemistry and Molecular Biology*; 2019 May 3;294(18):7137–50.
52. Ramachandran V, Shah KH, Herman PK. The cAMP-dependent protein kinase signaling pathway is a key regulator of P body foci formation. *2011 Sep 16*;43(6):973–81.

53. Wippich F, Bodenmiller B, Trajkovska MG, Wanka S, Aebersold R, Pelkmans L. Dual specificity kinase DYRK3 couples stress granule condensation/dissolution to mTORC1 signaling. *Cell*. 2013 Feb 14;152(4):791–805.
54. Rai AK, Chen J-X, Selbach M, Pelkmans L. Kinase-controlled phase transition of membraneless organelles in mitosis. *Nature Publishing Group. Nature Publishing Group*; 2018 Jul;559(7713):211–6.
55. Bański P, Kodiha M, Stochaj U. Chaperones and multitasking proteins in the nucleolus: networking together for survival? *Trends in Biochemical Sciences. Elsevier Current Trends*; 2010 Jul 1;35(7):361–7.
56. Hondele M, Sachdev R, Heinrich S, Wang J, Vallotton P, Fontoura BMA, et al. DEAD-box ATPases are global regulators of phase-separated organelles. *Nature Publishing Group. Nature Publishing Group*; 2019 Aug 21;357:eaa4382–5.
57. Mateju D, Franzmann TM, Patel A, Kopach A, Boczek EE, Maharana S, et al. An aberrant phase transition of stress granules triggered by misfolded protein and prevented by chaperone function. *EMBO J. EMBO Press*; 2017 Apr 4;e201695957–19.
58. Qamar S, Wang G, Randle SJ, Ruggeri FS, Varela JA, Lin JQ, et al. FUS Phase Separation Is Modulated by a Molecular Chaperone and Methylation of Arginine Cation- π Interactions. *Cell*. 2018 Apr 19;173(3):720–734.e15.
59. Liu Z, Zhang S, Gu J, Tong Y, Li Y, Gui X, et al. Hsp27 chaperones FUS phase separation under the modulation of stress-induced phosphorylation. *Nat Struct Mol Biol. Springer US*; 2020 Mar 20;:1–24.
60. Tauber D, Tauber G, Parker R. Mechanisms and Regulation of RNA Condensation in RNP Granule Formation. *Trends in Biochemical Sciences*. 2020 May 11.
61. Assenza S, Sassi AS, Kellner R, Schuler B, De Los Rios P, Barducci A. Efficient conversion of chemical energy into mechanical work by Hsp70 chaperones. *eLife. eLife Sciences Publications Limited*; 2019 Dec 17;8:223.
62. Goloubinoff P, Sassi AS, Fauvet B, Barducci A, De Los Rios P. Chaperones convert the energy from ATP into the nonequilibrium stabilization of native proteins. *Nature Chemical Biology. Nature Publishing Group*; 2018 Apr;14(4):388–95.
63. Hubstenberger A, Noble SL, Cameron C, Evans TC. Translation repressors, an RNA helicase, and developmental cues control RNP phase transitions during early development. *Dev Cell*. 2013 Oct 28;27(2):161–73.
64. Elbaum-Garfinkle S, Kim Y, Szczepaniak K, Chen CC-H, Eckmann CR, Myong S, et al. The disordered P granule protein LAF-1 drives phase separation into droplets with tunable viscosity and dynamics. *Proc Natl Acad Sci USA. National Acad Sciences*; 2015 Jun 9;112(23):7189–94.
65. Chakrabarti S, Hyeon C, Ye X, Lorimer GH, Thirumalai D. Molecular chaperones maximize the native state yield on biological times by driving substrates out of equilibrium. *Proc Natl Acad Sci USA. National Academy of Sciences*; 2017 Dec 19;114(51):E10919–27.
66. Tauber D, Tauber G, Khong A, Van Treeck B, Pelletier J, Parker R. Modulation of RNA Condensation by the DEAD-Box Protein eIF4A. *Cell. Elsevier Inc*; 2020 Jan 7;:1–33.
67. Calderwood SK, Murshid A, Prince T. The shock of aging: molecular chaperones and the heat shock response in longevity and aging--a mini-review. *Gerontology. Karger Publishers*; 2009;55(5):550–8.
68. Weber CA, Zwicker D, Jülicher F, Lee CF. Physics of active emulsions. *Rep Prog Phys. IOP Publishing*; 2019 Apr 30;82(6):064601–41.


69. Choi J-M, Dar F, Pappu RV. LASSI: A lattice model for simulating phase transitions of multivalent proteins. Keskin O, editor. *PLoS Comput Biol*. Public Library of Science; 2019 Oct;15(10):e1007028.
70. Riback JA, Brangwynne CP. Can phase separation buffer cellular noise? *Science*. American Association for the Advancement of Science; 2020 Jan 24;367(6476):364–5.
71. Klosin A, Oltsch F, Harmon T, Honigmann A, Jülicher F, Hyman AA, et al. Phase separation provides a mechanism to reduce noise in cells. *Science*. American Association for the Advancement of Science; 2020 Jan 24;367(6476):464–8.
72. Nott TJ, Craggs TD, Baldwin AJ. Membraneless organelles can melt nucleic acid duplexes and act as biomolecular filters. *Nat Chem*. 2016 Jun;8(6):569–75.
73. Schmidt HB, Görlich D. Transport Selectivity of Nuclear Pores, Phase Separation, and Membraneless Organelles. *Trends in Biochemical Sciences*. 2016 Jan;41(1):46–61.
74. Al-Husini N, Tomares DT, Pfaffenberger ZJ, Muthunayake NS, Samad MA, Zuo T, et al. BR-Bodies Provide Selectively Permeable Condensates that Stimulate mRNA Decay and Prevent Release of Decay Intermediates. *Mol Cell*. Elsevier Inc; 2020 Apr 13;:1–22.
75. Sheu-Gruttadauria J, MacRae IJ. Phase Transitions in the Assembly and Function of Human miRISC. *Cell*. 2018 Mar 16.
76. Strulson CA, Molden RC, Keating CD, Bevilacqua PC. RNA catalysis through compartmentalization. *Nat Chem*. Nature Research; 2012 Nov;4(11):941–6.
77. Reinkemeier CD, Girona GE, Lemke EA. Designer membraneless organelles enable codon reassignment of selected mRNAs in eukaryotes. *Science*. 2019 Mar 29;363(6434):eaaw2644.
78. Gallego LD, Schneider M, Mittal C, Romanauska A, Gudino Carrillo RM, Schubert T, et al. Phase separation directs ubiquitination of gene-body nucleosomes. Nature Publishing Group. Nature Publishing Group; 2020 Mar;579(7800):592–7.
79. Du M, Chen ZJ. DNA-induced liquid phase condensation of cGAS activates innate immune signaling. *Science*. 2018 Aug 17;361(6403):704–9.
80. Fang X, Wang L, Ishikawa R, Li Y, Fiedler M, Liu F, et al. Arabidopsis FLL2 promotes liquid-liquid phase separation of polyadenylation complexes. Nature Publishing Group. Nature Publishing Group; 2019 May 1;18:285.
81. Prouteau M, Loewith R. Regulation of Cellular Metabolism through Phase Separation of Enzymes. *Biomolecules*. Multidisciplinary Digital Publishing Institute; 2018 Dec 3;8(4):160.
82. Wang C, Schmich F, Srivatsa S, Weidner J, Beerenwinkel N, Spang A. Context-dependent deposition and regulation of mRNAs in P-bodies. *eLife*. eLife Sciences Publications Limited; 2018 Jan 3;7:166.
83. Hooper C, Hilliker A. Packing them up and dusting them off: RNA helicases and mRNA storage. *Biochim Biophys Acta*. 2013 Aug;1829(8):824–34.
84. Ivanov P, Kedersha N, Anderson P. Stress Granules and Processing Bodies in Translational Control. *Cold Spring Harb Perspect Biol*. Cold Spring Harbor Lab; 2018 Aug 6;:a032813.
85. Cisse II, Izeddin I, Causse SZ, Boudarene L, Senecal A, Muresan L, et al. Real-time dynamics of RNA polymerase II clustering in live human cells. *Science*. American Association for the Advancement of Science; 2013 Aug 9;341(6146):664–7.

86. Kwon I, Kato M, Xiang S, Wu L, Theodoropoulos P, Mirzaei H, et al. Phosphorylation-regulated binding of RNA polymerase II to fibrous polymers of low-complexity domains. *Cell*. 2013 Nov 21;155(5):1049–60.
87. Cho W-K, Spille J-H, Hecht M, Lee C, Li C, Grube V, et al. Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science*. American Association for the Advancement of Science; 2018 Jun 21;:eaar4199.
88. Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, et al. Coactivator condensation at super-enhancers links phase separation and gene control. *Science*. American Association for the Advancement of Science; 2018 Jun 21;19:eaar3958.
89. Boehning M, Dugast-Darzacq C, Rankovic M, Hansen AS, Yu T, Marie-Nelly H, et al. RNA polymerase II clustering through carboxy-terminal domain phase separation. *Nat Struct Mol Biol*. Nature Publishing Group; 2018 Aug 20;18(Suppl 6):285–840.
90. Boija A, Klein IA, Sabari BR, Dall'Agnese A, Coffey EL, Zamudio AV, et al. Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell*. 2018 Dec 13;175(7):1842–1855.e16.
91. Shrinivas K, Sabari BR, Coffey EL, Klein IA, Boija A, Zamudio AV, et al. Enhancer Features that Drive Formation of Transcriptional Condensates. *Mol Cell*. 2019 Aug 8;75(3):549–561.e7.
92. Guo C, Che Z, Yue J, Xie P, Hao S, Xie W, et al. ENL initiates multivalent phase separation of the super elongation complex (SEC) in controlling rapid transcriptional activation. *Sci Adv*. American Association for the Advancement of Science; 2020 Apr;6(14):eaay4858.
93. Lee CF, Brangwynne CP, Gharakhani J, Hyman AA, Jülicher F. Spatial organization of the cell cytoplasm by position-dependent phase separation. *Phys Rev Lett*. American Physical Society; 2013 Aug 23;111(8):088101.
94. Brangwynne CP, Eckmann CR, Courson DS, Rybarska A, Hoege C, Gharakhani J, et al. Germline P granules are liquid droplets that localize by controlled dissolution/condensation. *Science*. American Association for the Advancement of Science; 2009 Jun 26;324(5935):1729–32.
95. Saha S, Weber CA, Nusch M, Adame-Arana O, Hoege C, Hein MY, et al. Polar Positioning of Phase-Separated Liquid Compartments in Cells Regulated by an mRNA Competition Mechanism. *Cell*. Elsevier Inc; 2016 Aug 31;:1–30.
96. Smith J, Calidas D, Schmidt H, Lu T, Rasoloson D, Seydoux G. Spatial patterning of P granules by RNA-induced phase separation of the intrinsically-disordered protein MEG-3. *eLife*. eLife Sciences Publications Limited; 2016 Dec 3;5:803.
97. Guo YE, Manteiga JC, Henninger JE, Sabari BR, Dall'Agnese A, Hannett NM, et al. Pol II phosphorylation regulates a switch between transcriptional and splicing condensates. *Nature Publishing Group*. Nature Publishing Group; 2019 Aug 7;13(7770):720–6.
98. Lu H, Yu D, Hansen AS, Ganguly S, Liu R, Heckert A, et al. Phase-separation mechanism for C-terminal hyperphosphorylation of RNA polymerase II. *Nature Publishing Group*. Nature Publishing Group; 2018 Jun 1;558(7709):318–23.
99. Portz B, Shorter J. Switching Condensates: The CTD Code Goes Liquid. *Trends in Biochemical Sciences*. 2020 Jan;45(1):1–3.
100. Li P, Banjade S, Cheng H-C, Kim S, Chen B, Guo L, et al. Phase transitions in the assembly of multivalent signalling proteins. *Nature*. Nature Research; 2012 Mar 15;483(7389):336–40.
101. Altmeyer M, Neelsen KJ, Teloni F, Pozdnyakova I, Pellegrino S, Grøfte M, et al. Liquid demixing of intrinsically disordered proteins is seeded by poly(ADP-ribose). *Nat Commun*. Nature Publishing Group; 2015 Aug 19;6(1):8088–12.


102. Leung AKL. Poly(ADP-ribose): A Dynamic Trigger for Biomolecular Condensate Formation. *Trends Cell Biol.* 2020 May;30(5):370–83.
103. Lasker K, Diezmann L, Zhou X, Ahrens DG, Mann TH, Moerner WE, et al. Selective sequestration of signalling proteins in a membraneless organelle reinforces the spatial regulation of asymmetry in *Caulobacter crescentus*. *Nat Microbiol.* Springer US; 2020 Jan 8;;1–26.
104. Su X, Ditlev JA, Hui E, Xing W, Banjade S, Okrut J, et al. Phase separation of signaling molecules promotes T cell receptor signal transduction. *Science.* American Association for the Advancement of Science; 2016 Apr 29;352(6285):595–9.
105. Woodruff JB, Wueseke O, Viscardi V, Mahamid J, Ochoa SD, Bunkenborg J, et al. Centrosomes. Regulated assembly of a supramolecular centrosome scaffold in vitro. *Science.* American Association for the Advancement of Science; 2015 May 15;348(6236):808–12.
106. Case LB, Zhang X, Ditlev JA, Rosen MK. Stoichiometry controls activity of phase-separated clusters of actin signaling proteins. *Science.* 2019 Mar 8;363(6431):1093–7.
107. Hnisz D, Shrinivas K, Young RA, Chakraborty AK, Sharp PA. A Phase Separation Model for Transcriptional Control. *Cell.* 2017 Mar 23;169(1):13–23.
108. McKnight SL. Passing Through. *Trends in Biochemical Sciences.* 2019 Nov;44(11):899–901.
109. Ishov AM, Gurumurthy A, Bungert J. Coordination of transcription, processing, and export of highly expressed RNAs by distinct biomolecular condensates. *Emerg Top Life Sci.* 2020 Apr 27;57(8):2462.
110. Hilliker A, Gao Z, Jankowsky E, Parker R. The DEAD-box protein Ded1 modulates translation by the formation and resolution of an eIF4F-mRNA complex. 2011 Sep 16;43(6):962–72.
111. Xiol J, Spinelli P, Laussmann MA, Homolka D, Yang Z, Cora E, et al. RNA clamping by Vasa assembles a piRNA amplifier complex on transposon transcripts. *Cell.* 2014 Jun 19;157(7):1698–711.
112. Valentin-Vega YA, Wang Y-D, Parker M, Patmore DM, Kanagaraj A, Moore J, et al. Cancer-associated DDX3X mutations drive stress granule assembly and impair global translation. *Sci Rep.* Nature Publishing Group; 2016 May 16;6(1):25996.
113. Vecchi G, Sormanni P, Mannini B, Vandelli A, Tartaglia GG, Dobson CM, et al. Proteome-wide observation of the phenomenon of life on the edge of solubility. *Proc Natl Acad Sci USA.* National Academy of Sciences; 2019 Dec 31;116(24):12444–12451.
114. Parry BR, Surovtsev IV, Cabeen MT, O'Hern CS, Dufresne ER, Jacobs-Wagner C. The bacterial cytoplasm has glass-like properties and is fluidized by metabolic activity. *Cell.* 2014 Jan 16;156(1-2):183–94.
115. Joyner RP, Tang JH, Helenius J, Dultz E, Brune C, Holt LJ, et al. A glucose-starvation response regulates the diffusion of macromolecules. *eLife.* eLife Sciences Publications Limited; 2016;5:833.
116. Munder MC, Midtvedt D, Franzmann T, Nüske E, Otto O, Herbig M, et al. A pH-driven transition of the cytoplasm from a fluid- to a solid-like state promotes entry into dormancy. *eLife.* eLife Sciences Publications Limited; 2016;5:e09347.
117. Heimlicher MB, Bächler M, Liu M, Ibeneche-Nnewiwe C, Florin E-L, Hoenger A, et al. Reversible solidification of fission yeast cytoplasm after prolonged nutrient starvation. *J Cell Sci.* The Company of Biologists Ltd; 2019 Nov 8;132(21):jcs231688.

Valency


$v = 1$ Dimers




$v = 2$ Chains



$v = 3$ Networks



MIXED LIQUID




LCD: low complexity domain

liquid-liquid phase separation



CONDENSATE




reversible

liquid to solid






AGGREGATION




irreversible


MAINTAIN NON-EQUILIBRIUM




Buffering



Selective concentration hubs



Signaling cascades and gradients



ATPase-driven substrate flux

