

Is abdominal radiography still adapted for the screening of illegal intracorporeal containers (“body-packing”)? A comparative study with low-dose CT

**Pierre-Alexandre Poletti, MD^{1,2},
Laurent Canel³,
Christoph D Becker, MD¹,
Hans Wolff, MD⁴,
Bernice Elger, MD⁴,
Eric Lock³,
François Sarasin, MD²,
Monica S Bonfanti³,
Elise Dupuis Lozeron, MS⁵,
Thomas Perneger, MD PhD⁵,
Alexandra Platon, MD¹**

Department of Radiology, University Hospital of Geneva, Switzerland (1)

Emergency Center, University Hospital of Geneva, Switzerland (2)

Geneva State Police (3)

Department of Forensic Medicine, University Hospital of Geneva, Switzerland(4)

Division of Clinical Epidemiology, University Hospital of Geneva, Geneva, Switzerland (5)

Address Correspondence & Reprints

Pierre-A. Poletti, MD
Department of Radiology,
University Hospital of Geneva
4 rue Gabrielle Perret-Gentil
1211 Genève-14 Switzerland
Phone: 4122/ 372 81 59
Fax: 4122/ 372 70 72
e-mail: pierre-alexandre.poletti@hcuge.ch

This paper was presented at RSNA 2011 meeting

This paper was supported by a grant of the Academic Council of the Department of Radiology, University Hospital of Geneva, Switzerland

Original research

Word counts: 2898



**Is abdominal radiography still adapted for the screening of
illegal intra-corporeal containers ("body-packing")? A
comparative study with low-dose CT**

Journal:	<i>Radiology</i>
Manuscript ID:	RAD-11-2767.R1
Manuscript Type:	Meeting Paper 11
Manuscript Categorization Terms:	Adults < 1. SUBJECT MATTER, CT < 2. MODALITIES/TECHNIQUES, Conventional Radiography < 2. MODALITIES/TECHNIQUES, Abdomen/GI < 4. AREAS/SYSTEMS, Small Bowel < 5. STRUCTURES, Large Bowel < 5. STRUCTURES, Screening < 7. METHODOLOGY, Comparative Studies < 7. METHODOLOGY, Forensics < 6. TOPICS, Foreign Bodies < 6. TOPICS

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Is abdominal radiography still adapted for the screening of illegal intra-corporeal containers (“body-packing”)? A comparative study with low-dose CT

Original research

Advances in knowledge:

- 1) Abdominal radiography (AR) achieves a limited sensitivity (77%) for the screening of illegal intra-corporeal containers when compared to low-dose CT
- 2) Illegal intra-corporeal packets are difficult to detect by AR when they are in small number (< 12)
- 3) The sensitivity of AR for detection of illegal intra-corporeal containers is lower (50%) when they appear iso-dense to the bowel content at low-dose CT than when they appear denser (89%).

Implications for patient care:

Performing low-dose CT instead of AR will improve the detection of illegal intra-corporeal packets, without increasing the radiation dose.

Summary statement:

The use of low-dose CT may constitute a reasonable alternative to abdominal radiography to improve the detection of illegal intra-abdominal packets

1
2
3
4
5
6
7
8
9
10 **Abstract**

11
12 **Purpose:** To evaluate the diagnostic performance of the abdominal radiography (AR) to a low-dose
13 CT (LDCT) in the detection of illegal intra-corporeal containers (“packets”) using LDCT as a
14 reference standard.
15
16
17

18
19 **Materials and Methods:** This study was approved by the institutional ethical review board ; a
20 written informed consent was required (CER 06-023).
21
22

23
24 330 consecutive persons, suspected of having ingested drug packets, underwent a supine AR. The
25 presence or absence of packets at AR were reported and compared to the result of LDCT ,
26 considered reference standard. The density and the number of packets (≤ 12 or above) at LDCT were
27 recorded and analyzed to determine if they may influence the AR interpretation.
28
29
30

31
32 **Results:** Packets were detected at LDCT in 53 (16%) suspects. The sensitivity of AR for depiction
33 of packets was 77% (41/53), the specificity 96% (267/277). The packets appeared iso-dense to the
34 bowel contents on LDCT in sixteen (30%) of the 53 positive cases. Nineteen (36%) of the 53
35 positive LDCT displayed fewer than 12 packets. Iso-dense packets on LDCT and having a low
36 number of packets (≤ 12) were both significantly associated with false negative AR exams.
37
38
39
40
41
42

43
44 **Conclusion:** AR is mainly limited by a low sensitivity when compared to LDCT for the screening of
45 persons suspected of carrying drug packets. LDCT constitutes an efficient imaging alternative to
46
47
48
49 AR.

Introduction

The management of persons suspected of having swallowed drug containers is a complex procedure, which usually integrates the results of specific investigations (1, 2). Once a person has been suspected of carrying intra-corporeal drug containers, he (or she) is usually referred to a medical center to undergo a radiological examination. Usually, this examination consists in an abdominal radiography (AR), which is used to confirm or infirm the prior suspicion and, thus, to determine whether or not a stool analysis will be required (3-5). However, the value of the AR for the screening for intracorporeal containers has never been prospectively evaluated in a large series of suspects, using the CT as reference standard. Indeed, in spite of the fact that the CT has been reported to be the most accurate imaging method for displaying intra-abdominal containers (6-11), the dose of radiation typically delivered using this technique is a major limitation in its systematic use to screen persons suspected of conveying illegal intra-corporeal drug containers. This concern can now be overcome by the use of low-dose CT protocols, which deliver a dose of radiation close to that of an AR. These low-dose CT protocols have been reported to be accurate for the screening of well defined medical conditions, such as renal colic (12) and appendicitis (13-15) and have also been recently reported a useful in demonstrating the presence of an intracorporeal drug packet (16, 17).

The purpose of this study is to evaluate the diagnostic performance of the abdominal radiography (AR) to a low-dose CT (LDCT) in the detection of illegal intra-corporeal containers ("packets") using low-dose CT as a reference standard.

Materials and Methods

This research was approved by the Research Ethics Committee of our institution (CER 06-023) ; written informed consent was obtained. During the study period (July 2007 to July 2010), all consecutive adult persons suspected of having ingested drug packets within the State territory (n=338) were systematically brought to our emergency department, at any time of the day or the night, instead of being dispatched in various medical institutions to undergo an AR.

Immediately after admission in our emergency radiology unit, suspects underwent a supine AR. The AR was immediately interpreted by the radiologist on call (a fellow or a senior resident). The prior training for a fellow (respectively a senior resident) was 5 to 6 years of general radiology (respectively 3 to 5 years), including at least 6 months of conventional radiology and 2 years of body CT. Twenty-two fellows or senior residents did participate to the AR and LDCT reading during the study period. The radiologist on call had to report on a standardized electronic form whether AR was considered positive or negative for the presence of packets. An AR was considered positive for the presence of packet(s), when at least one of the following criteria was present (18): 1) one or multiple well defined densities in the stomach, small bowel and/or colon, not suggestive of alimentary content 2) a “double condom sign” (Fig. 1), defined as a definite crescent of air surrounding an ovoid density (3), 3) a smooth and uniformly shaped oblong structure (sometimes referred to as the “tic-tac” sign (18)) 4) the “parallelism sign”, defined as “rigid packages aligning parallel to each other in the bowel lumen”(18). The radiologist was also asked to report the degree of confidence in the AR interpretation, for the presence or absence of packets, on a scale from 1 to 4 (1= minimal confidence in the diagnosis, 2 = moderate confidence, 3= good confidence, 4= excellent confidence). The suspect’s BMI was also recorded.

Once the AR was interpreted, the suspect underwent an abdominal LDCT which was also immediately interpreted as positive or negative for the presence of packets by the same radiologist. The interpretation of LDCT was also reported on an electronic form. The radiologist had to indicate

1 whether the number of foreign bodies depicted on LDCT was less or equal to 12 (Fig. 2) or greater.

2
3
4 The visual aspect of the packets at LDCT was reported as iso-dense (Fig. 3) or hyper-dense with
5
6 regard to the intestinal content.

7
8 A brief report, just indicating the presence or absence of packets at both AR and LDCT, was given to
9
10 the police or boarder guard authorities.

11 12 13 14 15 **Reference standards**

16
17 LDCT was considered reference standard. When LDCT was negative, no further examination was
18
19 performed and the suspect left the hospital. When LDCT was positive, the suspect was hospitalized in
20
21 a dedicated ward of our institution and kept under surveillance for stool analysis. All collected packets
22
23 were handed over to a dedicated laboratory for chemical characterization of the content.

24 25 26 27 28 **Exclusion criteria**

29
30 Persons who refused to participate in the study, those under 18 year of age and pregnant women, were
31
32 excluded from the current survey. They usually underwent a stool analysis. A pregnancy test was
33
34 systematically obtained in every woman of childbearing age.

35 36 37 38 39 **Technical imaging parameters**

40
41 ARs were performed with the patient in the supine position using an X-Ray Philips Optimus 65 unit
42
43 with automatic exposure control (Philips Medical Systems, Best, the Netherlands).

44
45 LDCT were performed with a 16-row Philips MX 8000 (Philips Medical Systems, Best, the
46
47 Netherlands), from lung bases to pelvis, without administration of oral or rectal contrast material,
48
49 using the following parameters: 16x1.5 mm collimation, pitch 1.25, gantry rotation period 0.5
50
51 second, tube potential 120 kV, tube charge per gantry rotation 30 mAs ($75 \text{ mA} \times 0.5 \text{ s} / 1.25 = 30$
52
53 mAs), reconstruction slice thickness 3.0 mm.

Effective dose calculation

1. AR

For a field of 35 x 43 cm, the entrance doses delivered by the AR were 6.0 mGy (66 kV and 50 mA) to 12 mGy (73 kV and 120 mA), with an average of 9 ± 3.0 mGy. The effective doses (E) were computed using ODS 60 software(19); for women, $E = 2.0 \pm 0.7$ mSv and for men, $E = 1.3 \pm 0.4$ mSv.

2. LDCT

The dose delivered by LDCT was estimated using the ImPACT CT patient dosimetry calculator (20), using the default nCTDIw of 7.0 mGy/100 mAs proposed at 120 kV (value compatible with our measurements within 10%). The following results were provided by the program: DLP (women) = 84 ± 10.5 mGy cm; E (women) = 1.7 ± 0.2 mSv. DLP (men) = 84 ± 10.5 mGy cm; E (men) = 1.2 ± 0.1 mSv.

Statistical analysis

AR were compared to LDCT used as reference standard to estimate the sensitivity, the specificity, the positive predictive value (PPV) and the negative predictive value (NPV) for depiction of foreign bodies in the whole study population (n=330). The 95% confidence intervals of these statistics were estimated with the Clopper-Pearson method.

The radiologist's confidence in the AR analysis was considered as an ordinal variable consisting in a 8 level scale, from negative AR with a level of confidence of 4 (level 1) to a positive AR with confidence level of 4 (level 8). A ROC curve was obtained. The optimal cut-off on the level of confidence of the radiologist was determined by maximising Youden's index (sensitivity + specificity - 1)(21). Likelihood ratios (LR) were used to assess how informative were the levels of confidence given by the radiologists.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Considering the positive cases at LDCT only, a logistic regression model was obtained to test the influence of the conveyers' BMI (< 25 or ≥ 25), of the density and of the quantity of the packets on their detection on the AR. Furthermore, sensitivities were computed on the strata defined by the density (iso-dense versus hyper-dense) and the quantity (below 12 versus above 12) of the packets. The significance level was fixed to 5% (two-tailed). Differences between two groups were tested with t-test for continuous variables and with Fisher's exact test for proportions. All analyses were performed using R for Windows (version 2.13.0)

Results

Population and packets characteristics

Three hundred thirty-eight suspects were brought in our institution during the study period. Eight were excluded from the study because they refused to participate (n=6) or because they were pregnant (n=2). Thus, 330 suspects were included in our protocol, 296 (90%) men and 34 (10%) women, with a mean age of 32 years (range: 18 to 55). BMI was reported under 18.5 (underweight) in 22 (7%) suspects, between 18.5 and 25 (normal range) in 258 (78%), between 25 and 30 (overweight) in 44 (13%), and over 30 (obesity) in 6 (2%). Packets were detected by LDCT, and found at stool analysis, in 53 (16%) of the 330 suspects.

In 50 (94%) of the 53 true positive cases, packets content consisted of cocaine-hydrochlorate powder (also containing various cutting agents such as phenacetin), weighting between 7 and 25 grams each (mean 10.6 grams). In three persons, packets contained rolls of banknotes, wrapped in cellophane bags. Packets were located in the bowel or rectum (n=51) or in the vagina (n=2, 8 packets of drug in one, banknotes in the other). Twelve (23%) of the 53 drug conveyers carried between 1 and 6 packets, 7 (13%) between 7 and 12 packets, and 34 (64%) had more than 12 packets.

AR versus LDCT

1 The presence of packets was suspected by the radiologists at AR in 51 (15%) of the 330 suspects.
2
3
4 When compared to LDCT, 41 AR were true positive, 10 false positive, 267 true negative, 12 false
5
6 negative. The sensitivity, specificity, PPV and NPV of AR are reported in Table 1.
7
8
9

10 **Confidence of the radiologist in his/her AR interpretation**

11
12 The degree of confidence in the interpretation was equal to 1 in 17 (5%) of the 330 suspects, 2 in 86
13
14 (26%), 3 in 164 (50%) and 4 in 63 (19%). The ROC curve associated with confidence in the AR
15
16 analysis considered as an ordinal 8-level variable showed good discrimination (Fig. 4), as the area
17
18 under the ROC curve was 0.95. The optimal sensitivity and specificity (85% and 94% respectively)
19
20 are obtained at the cut-off point of 4, when a negative result with a level of confidence of 1 is
21
22 considered as a “positive” result. This cut point leads to a PPV of 74% (45/61) and to a NPV of
23
24 97% (261/269).
25
26
27

28 Table 2 shows the likelihood ratios with regard to the degree of confidence in the AR interpretation.
29
30 In this chart, level 1 and level 2 of confidence have been merged as low confidence, level 3 and 4 as
31
32 high confidence. Under these conditions, a strong LR (greater than 10 or lower than 1/10) was only
33
34 achieved, for positive or negative AR, when the confidence was high.
35
36
37
38
39

40 **Influence of the conveyers' BMI, of the density and of the quantity of the packets on the AR** 41 **interpretation.**

42
43 Among the 53 persons who carried packets, 43 had a BMI below 25 and 10 above 25 (including one
44
45 above 30). The packets' content was reported iso-dense compared to the bowel content on the
46
47 LDCT in 16 (30%) of the 53 positive cases, and hyper-dense in 37 (70%). Nineteen conveyers
48
49 carried 12 or less packets, 34 more than 12. Results of univariate logistic regression (Table 3)
50
51 showed a non-significant association between BMI >25 and a correct positive identification, but the
52
53 packets' density and quantity were strongly and significantly associated with true positive status.
54
55
56
57
58
59
60

1 The multivariate model confirmed these findings: packets were more difficult to detect when they
2 were iso-dense and/or fewer than 12.
3

4
5 The sensitivity of AR for depiction of packets with regard to their number and their density found at
6 LDCT is reported in Table 4. The highest sensitivity (92%, 24/26) of AR was achieved in the
7 presence of multiple (>12) packets of high density. The lowest sensitivity (25%, 2/8) was found in
8 the presence of a small number of packets (\leq 12) of a low density.
9
10
11
12
13
14
15
16

17 **Discussion**

18
19
20 This study aimed to assess the value of AR for the screening of the body-packers, when compared
21 to LDCT. Although some reports have already stressed the limitations of the AR in this setting (2,
22 6, 7, 22, 23), the diagnostic performance of the AR for identification of illegal packets remained
23 uncertain, due to the absence of a systematic reference standard in the prior reports. Indeed, no prior
24 series systematically included a CT examination or a stool analysis in every suspect after the AR
25 was reported negative. The lack of a straightforward reference standard explains the wide range of
26 sensitivities (from 40% to 100%) that have been previously reported for the detection of intra-
27 corporeal containers by AR (3, 7, 22, 24). In our prospective study, AR achieved an overall 77%
28 (41/53) sensitivity and 96% (267/277) specificity for depiction of packets, when compared to
29 LDCT, systematically obtained in every suspect. Our results show that this technique is mainly
30 limited by a high percentage (23%, 12/53) of false negative cases. Hence, the diagnostic value of
31 AR for the screening of drug conveyers is probably overestimated, and raises questions about its
32 exact role and its limitations in this application. Beside, using AR as sole screening test would
33 have led to falsely consider 4% (10/277) of innocent suspects as positive.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Our data showed that the value of AR is closely related to the confidence of the radiologist in
52 his/her interpretation. High levels of confidence (3 or 4) were associated with high likelihood ratios,
53 while low levels (1 or 2) were not. This observation suggests that an AR cannot be relied on for the
54 detection of packets when the radiologist is not confident in his/her interpretation, which
55
56
57
58
59
60

1 corresponds, in the current series, to 31% (104/330) of our study group. In such situations, an AR
2 should ideally be completed by a LDCT. However, a CT-scanner is not always available after the
3
4 should ideally be completed by a LDCT. However, a CT-scanner is not always available after the
5
6 AR has been performed, especially when the screening is not performed in a medical system (but,
7
8 for instance, within a remote airport). Beside, the additional cost of an unenhanced CT examination
9
10 (about 3 to 4 times the price of an AR) often precludes its systematic use in this setting. When the
11
12 screening of body-packers must be based on AR alone, our results (ROC curve) showed that the
13
14 optimal ratio for sensitivity (85%) and specificity (94%) is obtained by considering a negative AR
15
16 with a lower level of confidence (1 of 4) as positive result (cut-off point). Doing so will still lead to
17
18 a 15% rate of false negative and a 6% rate of false positive AR. Our results did not show any
19
20 relationship between the suspects' BMI and the AR results. However, the group of overweight
21
22 body-packers (BMI ≥ 25) only included 1 obese person (BMI > 30); it is therefore impossible to
23
24 draw any conclusion with regard to this specific sub-group.
25
26

27
28 The last objective of the study was to retrospectively determine whether the apparent density of the
29
30 packets at LDCT and/or their quantity might influence their detectability at AR. Our results showed
31
32 that the density and the number of the packets at LDCT were significantly correlated to the rate of
33
34 false negative readings, at both univariate and multivariate analysis. The association between the
35
36 radiological attenuation of various drugs and their possible presentations (powder, stones, tablets,
37
38 pills etc..) has already been reported in extra-corporeal analyses (5).
39
40

41
42 The fact that the sensitivity of the AR was only 50% (8/16) in suspects carrying iso-dense packets,
43
44 when compared to 89% (33/37) in those carrying dense containers, suggests that the increased X-
45
46 ray attenuation constitutes a major radiological sign for their detection at AR. Furthermore the
47
48 sensitivity dropped to 25% (2/8) when packets were both iso-dense and in small quantity (≤ 12).
49

50
51 With the improvement in the packets manufacturing, it would not be surprising if a majority of
52
53 intra-corporeal containers became undetectable at AR in a short time span. This supposition is
54
55 bolstered by recent reports of incidental seizure of liquid or mushy forms of intra-abdominal
56
57
58
59
60

1 concealed cocaine packets, that remained absolutely undetectable at AR, even at retrospective
2
3 analysis(25).
4

5
6 Our study had a number of limitations. In the current series, packets consisted exclusively in large
7
8 “finger like” containers from 7 to 25 grams each. Therefore, our data can certainly not be
9
10 transposed to a population of smugglers using smaller drug packets, sometimes referred to as
11
12 “body-stuffers” or “mini-packers” (10). In the latter situation, it is possible that the AR
13
14 interpretation would have led to a higher rate of false negative interpretations.
15

16
17 Similarly, in the current study, intra-corporeal containers contained only cocaine-hydrochlorate
18
19 powder (along with cutting agents), and banknotes in three cases, which is linked to the local trends
20
21 in drug trafficking. Whether our results can be extrapolated to other packets content (such as heroin
22
23 or liquid cocaine) remains an open question.
24

25
26 Finally, the methodology of the current series was based on the postulate that LDCT is a reference
27
28 standard for detection of intra-corporeal containers. However, no prior study has yet evaluated a
29
30 negative LDCT (or even a standard CT) with a systematic stool analysis. Since the sensitivity of
31
32 LDCT for detection of intra-abdominal packets is unknown, it can therefore not be completely
33
34 excluded that some intra-corporeal containers may have been missed by both AR and LDCT.
35

36
37 Nevertheless, our study results demonstrated the high specificity of LDCT in this setting; indeed, all
38
39 positive LDCTs of our study population were confirmed by stool analysis without any report of
40
41 false positive cases.
42

43
44 In conclusion, the current study shows that the detection of illegal intra-abdominal containers by
45
46 AR is related to their X-Ray attenuation and to their quantity. The interpretation of AR should take
47
48 into consideration the level of confidence of the radiologist in his/her interpretation to optimize both
49
50 sensitivity and specificity. The use of LDCT may constitute a reasonable alternative to AR to
51
52 improve the detection of illegal intra-abdominal packets
53
54

55 56 57 **References** 58 59 60

1. Gherardi R, Marc B, Alberti X, Baud F, Diamant-Berger O. A cocaine body packer with normal abdominal plain radiograms. Value of drug detection in urine and contrast study of the bowel. *Am J Forensic Med Pathol.* 1990;11(2):154-7.
2. Gherardi RK, Baud FJ, Leporc P, Marc B, Dupeyron JP, Diamant-Berger O. Detection of drugs in the urine of body-packers. *Lancet.* 1988;1(8594):1076-8.
3. Beerman R, Nunez D, Jr., Wetli CV. Radiographic evaluation of the cocaine smuggler. *Gastrointest Radiol.* 1986;11(4):351-4.
4. Algra PR, Brogdon BG, Marugg RC. Role of radiology in a national initiative to interdict drug smuggling: the Dutch experience. *AJR Am J Roentgenol.* 2007;189(2):331-6.
5. Hergan K, Kofler K, Oser W. Drug smuggling by body packing: what radiologists should know about it. *Eur Radiol.* 2004;14(4):736-42.
6. Kersschot EA, Beaucourt LE, Degryse HR, De Schepper AM. Roentgenographical detection of cocaine smuggling in the alimentary tract. *Rofo.* 1985;142(3):295-8.
7. Karhunen PJ, Suoranta H, Penttila A, Pitkaranta P. Pitfalls in the diagnosis of drug smuggler's abdomen. *J Forensic Sci.* 1991;36(2):397-402.
8. Sengupta A, Page P. Window manipulation in diagnosis of body packing using computed tomography. *Emerg Radiol.* 2008;15(3):203-5.
9. Freisleder A, Bautz W, Schmidt V. [Body packing: the value of modern imaging procedures in the detection of intracorporeal transport media]. *Arch Kriminol.* 1988;182(5-6):143-53.
10. Schmidt S, Hugli O, Rizzo E, et al. Detection of ingested cocaine-filled packets--diagnostic value of unenhanced CT. *Eur J Radiol.* 2008;67(1):133-8.
11. Yang RM, Li L, Feng J, et al. Heroin body packing: clearly discerning drug packets using CT. *South Med J.* 2009;102(5):470-5.
12. Poletti PA, Platon A, Rutschmann OT, Schmidlin FR, Iselin CE, Becker CD. Low-dose versus standard-dose CT protocol in patients with clinically suspected renal colic. *AJR Am J Roentgenol.* 2007;188(4):927-33.
13. Keyzer C, Tack D, de Maertelaer V, Bohy P, Gevenois PA, Van Gansbeke D. Acute appendicitis: comparison of low-dose and standard-dose unenhanced multi-detector row CT. *Radiology.* 2004;232(1):164-72.
14. Platon A, Jlassi H, Rutschmann OT, et al. Evaluation of a low-dose CT protocol with oral contrast for assessment of acute appendicitis. *Eur Radiol.* 2009;19(2):446-54.
15. Seo H, Lee KH, Kim HJ, et al. Diagnosis of acute appendicitis with sliding slab ray-sum interpretation of low-dose unenhanced CT and standard-dose i.v. contrast-enhanced CT scans. *AJR Am J Roentgenol.* 2009;193(1):96-105.
16. Maurer MH, Niehues SM, Schnapauff D, et al. Low-dose computed tomography to detect body-packing in an animal model. *Eur J Radiol.* 78(2):302-6.
17. Beauverd Y, Poletti PA, Wolff H, Ris F, Dumonceau JM, Elger BS. A body-packer with a cocaine bag stuck in the stomach. *World J Radiol.* 3(6):155-8.
18. Niewiarowski S, Gogbashian A, Afaq A, Kantor R, Win Z. Abdominal X-ray signs of intra-intestinal drug smuggling. *J Forensic Leg Med.* 17(4):198-202.
19. Sermovaa A, Rannikko S, Nikitin V, et al. A topographically and anatomically unified phantom model for organ dose determination in radiation hygiene. Report STUK-A87, Finnish Centre for Radiation and Nuclear Safety, Helsinki. 1989.
20. ImPACT. Imaging performance assessment of CT scanners: a medical devices agency evaluation group. London UK www.impactscan.org/ctdosimetry.htm. 2006.
21. Fluss R, Faraggi D, Reiser B. Estimation of the Youden Index and its associated cutoff point. *Biom J.* 2005;47(4):458-72.
22. McCarron MM, Wood JD. The cocaine 'body packer' syndrome. Diagnosis and treatment. *Jama.* 1983;250(11):1417-20.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
23. Traub SJ, Hoffman RS, Nelson LS. False-positive abdominal radiography in a body packer resulting from intraabdominal calcifications. *Am J Emerg Med.* 2003;21(7):607-8.
24. Bogusz MJ, Althoff H, Erkens M, Maier RD, Hofmann R. Internally concealed cocaine: analytical and diagnostic aspects. *J Forensic Sci.* 1995;40(5):811-5.
25. Mourachko R. Oral communication. Council of Europe, annual airport group meeting, Strasbourg. 2011.

Table 1 Evaluation of abdominal radiography for detection of body-packs, using low-dose CT as reference standard

	Estimated Value	95 % Confidence Interval
Sensitivity	41/53 = 0.77	0.64 – 0.88
Specificity	267/277 = 0.96	0.93 – 0.98
Positive predictive value	41/51 = 0.80	0.67 – 0.90
Negative predictive value	267/279 = 0.96	0.93 – 0.98

Table 2 Results of the abdominal radiography (AR) interpretation according to the confidence of the radiologist in his/her interpretation, when using low-dose CT (LDCT) as reference standard

	Positive LDCT	Negative LDCT	Likelihood Ratio ⁽³⁾
Negative AR High confidence ⁽¹⁾	2 (3.8%)	187 (67.5%)	0.06
Negative AR Low confidence ⁽²⁾	10 (18.9%)	80 (28.9%)	0.65
Positive AR Low confidence	5 (9.4%)	8 (2.9%)	3.27
Positive AR High confidence	36 (67.9%)	2 (0.7%)	94.08
Total	53 (100%)	277 (100%)	

(1) High confidence: reported 3 or 4 on a 4 level scale (2) Low confidence : reported 1 or 2 on a 4 level scale (3) A strong likelihood ratio is considered when greater than 10 or below 1/10

Table 3 Association between the suspect's body mass index (BMI), the quantity and the density of the packets and their detection at abdominal radiography when compared to low-dose CT (reference standard)

		Univariate		Multivariate	
		Odds Ratio (95% CI)	p-value	Odds Ratio (95% CI)	p-value
BMI	< 25 ⁽¹⁾	1			
	≥ 25	3.1 (0.4-27.3)	0.31		
Density	Iso-dense ⁽¹⁾	1		1	
	Hyper-dense	8.3 (2.0-34.4)	0.004	7.5 (1.6-33.7)	0.009
Quantity	≤ 12 ⁽¹⁾	1		1	
	> 12	5.5 (1.4-21.8)	0.016	4.8 (1.1-21.9)	0.042

(1) Reference level for comparison

Table 4 Sensitivity of abdominal radiography for depiction of intracorporeal packets with regard to their quantity and density at low-dose CT

		N	Sensitivity
Density			
	Iso density	8/16	0.50 (0.25-0.75)
	High density	33/37	0.89 (0.75-0.97)
Quantity			
	≤12 packets	11/19	0.58 (0.33-0.80)
	>12 packets	30/34	0.89 (0.73-0.97)
Both variables			
	Iso density ≤12 packets	2/8	0.25 (0.03-0.65)
	Iso density >12 packets	6/8	0.75 (0.35-0.97)
	High density ≤12 packets	9/11	0.82 (0.48-0.98)
	High density >12 packets	24/26	0.92 (0.75-0.99)

Number between parentheses are 95% CI

FIGURE CAPTIONS**Figure 1**

True positive abdominal radiography (AR) in a 30 year-old man suspected of conveying drug packets.

Admission AR (a) shows multiple well defined smooth and uniformly shaped oblong densities (asterisks) spread throughout the abdomen, not suggestive of alimentary content, with peripheral crescent of air or “double condom sign” (arrow), consistent with drug packets. The radiologist on call did report this AR as positive, with a very high confidence (4/4) in his diagnosis. Axial (b) low-dose CT images shows a large quantity (>12) of intra-intestinal packets, hyper-dense to the surrounding bowel content. Eighty containers stuffed with cocaine were found at stool analysis (c).

Figure 2

False negative abdominal radiography (AR) in a 26 year-old man, suspected of conveying drug packets.

Admission AR (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis.

Axial (b) and sagittal MPR reformatted (c) low-dose CT images, show a small quantity (≤ 12) of packets within the rectum, hyper-dense to the surrounding bowel content (arrow). Six containers stuffed with cocaine were found at stool analysis (d).

Figure 3

False negative abdominal radiography (AR) in a 54 year-old man, suspected of conveying drug packets.

Admission AR (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis.

Axial (b) low-dose CT images, shows a large quantity (>12) of packets within the stomach (arrowheads) and the small bowel (arrow), iso-dense to the surrounding gastric and bowel content. Three dozen of containers stuffed with cocaine were found at stool analysis.

Figure 4

ROC curve displaying the sensitivity and specificity of the abdominal radiography (AR) interpretation for the presence of intra-corporeal packets, with regard to the level of confidence of the radiologist in his/her interpretation

Points 1 to 8 correspond to the sensitivity and specificity for depiction of packets if the AR were considered positive at these threshold levels:

Point 1: corresponds to an AR reported as negative by the radiologist, with a high level of confidence (4 of 4)

Point 2: AR reported as negative, level of confidence of 3

Point 3: AR reported as negative, level of confidence of 2

Point 4: AR reported as negative, level of confidence of 1

Point 5: AR reported as positive, level of confidence of 1

Point 6: AR reported as positive, level of confidence of 2

Point 7: AR reported as positive, level of confidence of 3

Point 8: AR reported as positive, level of confidence of 4

The optimal sensitivity and specificity (85% and 94% respectively) are obtained at the cut-off point of 4, when a negative result with a level of confidence of 1 is considered as a positive result.




Figure 1a True positive abdominal plain film (APF) in a 30 year-old man suspected of conveying drug packets.

Admission APF (a) shows multiple well defined smooth and uniformly shaped oblong densities (asterisks) spread throughout the abdomen, not suggestive of alimentary content, with peripheral crescent of air or "double condom sign" (arrow), consistent with drug packets. The radiologist on call did report this

APF as positive, with a very high confidence (4/4) in his diagnosis. Axial (b) low-dose CT images shows a large quantity (>12) of intra-intestinal packets, hyper-dense to the surrounding bowel content. Eighty containers stuffed with cocaine were found at stool analysis (c).

103x116mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1b True positive abdominal plain film (APF) in a 30 year-old man suspected of conveying drug packets.

Admission APF (a) shows multiple well defined smooth and uniformly shaped oblong densities (asterisks) spread throughout the abdomen, not suggestive of alimentary content, with peripheral crescent of air or "double condom sign" (arrow), consistent with drug packets. The radiologist on call did report this APF as positive, with a very high confidence (4/4) in his diagnosis. Axial (b) low-dose CT images shows a large quantity (>12) of intra-intestinal packets, hyper-dense to the surrounding bowel content. Eighty containers stuffed with cocaine were found at stool analysis (c).

117x119mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 1c True positive abdominal plain film (APF) in a 30 year-old man suspected of conveying drug packets.

Admission APF (a) shows multiple well defined smooth and uniformly shaped oblong densities (asterisks) spread throughout the abdomen, not suggestive of alimentary content, with peripheral crescent of air or "double condom sign" (arrow), consistent with drug packets. The radiologist on call did report this

APF as positive, with a very high confidence (4/4) in his diagnosis. Axial (b) low-dose CT images shows a large quantity (>12) of intra-intestinal packets, hyper-dense to the surrounding bowel content. Eighty containers stuffed with cocaine were found at stool analysis (c).

90x33mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 2a False negative abdominal plain film (APF) in a 26 year-old man, suspected of conveying drug packets.

Admission APF (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis. Axial (b) and sagittal MPR reformatted (c) low-dose CT images, show a small quantity (< 12) of packets within the rectum, hyper-dense to the surrounding bowel content (arrow). Six containers stuffed with cocaine were found at stool analysis (d).

119x150mm (300 x 300 DPI)



Figure 2b False negative abdominal plain film (APF) in a 26 year-old man, suspected of conveying drug packets.

Admission APF (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis.

Axial (b) and sagittal MPR reformatted (c) low-dose CT images, show a small quantity (< 12) of packets within the rectum, hyper-dense to the surrounding bowel content (arrow). Six containers stuffed with cocaine were found at stool analysis (d).

119x82mm (300 x 300 DPI)




Figure 2c False negative abdominal plain film (APF) in a 26 year-old man, suspected of conveying drug packets.

Admission APF (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis. Axial (b) and sagittal MPR reformatted (c) low-dose CT images, show a small quantity (< 12) of packets within the rectum, hyper-dense to the surrounding bowel content (arrow). Six containers stuffed with cocaine were found at stool analysis (d).

73x119mm (300 x 300 DPI)



Figure 2d False negative abdominal plain film (APF) in a 26 year-old man, suspected of conveying drug packets.

Admission APF (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis. Axial (b) and sagittal MPR reformatted (c) low-dose CT images, show a small quantity (< 12) of packets within the rectum, hyper-dense to the surrounding bowel content (arrow). Six containers stuffed with cocaine were found at stool analysis (d).

106x48mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 3a False negative abdominal plain film (APF) in a 54 year-old man, suspected of conveying drug packets.

Admission APF (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis. Axial (b) low-dose CT images, shows a large quantity (>12) of packets within the stomach (arrowheads) and the small bowel (arrow), iso-dense to the surrounding gastric and bowel content. Three dozen of containers stuffed with cocaine were found at stool analysis.

91x119mm (300 x 300 DPI)



Figure 3b False negative abdominal radiography (AR) in a 54 year-old man, suspected of conveying drug packets.

Admission AR (a) was initially reported as negative by the radiologist on call, with a low degree of confidence (1/4) in his diagnosis. Axial (b) low-dose CT images, shows a large quantity (>12) of packets within the stomach (arrowheads) and the small bowel (arrow), iso-dense to the surrounding gastric and bowel content. Three dozen of containers stuffed with cocaine were found at stool analysis.

119x93mm (300 x 300 DPI)




Figure 4 ROC curve displaying the sensitivity and specificity of the abdominal radiography (AR) interpretation for the presence of intra-corporeal packets, with regard to the level of confidence of the radiologist in his/her interpretation
 Points 1 to 8 correspond to the sensitivity and specificity for depiction of packets if the AR were considered positive at these threshold levels:
 Point 1: corresponds to an AR reported as negative by the radiologist, with a high level of confidence (4 of 4)
 Point 2: AR reported as negative, level of confidence of 3
 Point 3: AR reported as negative, level of confidence of 2
 Point 4: AR reported as negative, level of confidence of 1
 Point 5: AR reported as positive, level of confidence of 1
 Point 6: AR reported as positive, level of confidence of 2
 Point 7: AR reported as positive, level of confidence of 3
 Point 8: AR reported as positive, level of confidence of 4
 The optimal sensitivity and specificity (85% and 94% respectively) are obtained at the cut-off point of 4, when a negative result with a level of confidence of 1 is considered as a positive result.

58x42mm (600 x 600 DPI)