

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

### Electron Transfer across *ortho*-Phenylene Wires

Journal:	<i>The Journal of Physical Chemistry</i>
Manuscript ID	jp-2018-11236a.R1
Manuscript Type:	Article
Date Submitted by the Author:	15-Dec-2018
Complete List of Authors:	Malzkuhn, Sabine; Universitat Basel, Department of Chemistry Guo, Xingwei; Universitat Basel, Department of Chemistry Häussinger, Daniel; Universitat Basel, Department of Chemistry Wenger, Oliver; Universitat Basel, Department of Chemistry

SCHOLARONE™  
Manuscripts

1  
2  
3 Electron Transfer across *ortho*-Phenylene Wires  
4  
5  
6  
7

8 Sabine Malzkuhn, Xingwei Guo, Daniel Häussinger, and Oliver S. Wenger\*  
9

10  
11  
12 Department of Chemistry, University of Basel, St. Johannis-Ring 19, 4056 Basel, Switzerland  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Abstract:**

Photoinduced electron transfer across rigid rod-like oligo-*para*-phenylenes has been thoroughly investigated in the past, but their *ortho*-connected counterparts are yet entirely unexplored in this regard. We report on three molecular dyads comprised of a triarylamine donor and a Ru(bpy)<sub>3</sub><sup>2+</sup> (bpy =2,2'-bipyridine) acceptor connected covalently by 2 to 6 *ortho*-phenylene units. Pulsed excitation of the Ru(II) sensitizer at 532 nm leads to the rapid formation of oxidized triarylamine and reduced ruthenium complex via intramolecular electron transfer. The subsequent thermal reverse charge-shift reaction to reinstate the electronic ground-state occurs on a timescale of 120 – 220 ns in de-aerated CH<sub>3</sub>CN at 25 °C. The conformational flexibility of the *ortho*-phenylene bridges causes multi-exponential transient absorption kinetics for the photoinduced forward process, but the thermal reverse reaction produces single-exponential transient absorption decays. The key finding is that the flexible *ortho*-phenylene bridges permit rapid formation of photoproducts storing ca. 1.7 eV of energy with lifetimes on the order of hundreds of nanoseconds, similar to what is possible with rigid rod-like donor-acceptor compounds. Thus, the conformational flexibility of the *ortho*-phenylenes represents no disadvantage with regard to the photoproduct lifetimes, and this is relevant in the greater context of light-to-chemical energy conversion.


## Introduction

Long-range electron transfers play key roles in many biological and artificial systems, and neither photosynthesis nor respiration would function without them.<sup>1</sup> With artificial photosynthesis as a long-term perspective, many prior investigations concentrated on obtaining long-lived electron-hole pairs in donor-bridge-acceptor compounds,<sup>2-10</sup> whereas other studies explored the distance or driving-force dependence of electron transfer rates ( $k_{ET}$ ).<sup>11-19</sup> Rigid rod-like molecular bridges are usually preferred, because they yield systems with well-defined donor-acceptor distances and restricted conformational degrees of freedom, simplifying the analysis of kinetic data. Oligo-*para*-phenylenes are a prototype class of relatively rigid spacers and as such have received considerable attention,<sup>20-36</sup> but to our knowledge there have been no prior studies of photoinduced long-range electron transfer across oligo-*ortho*-phenylenes.

Recently, much progress has been made in the controlled synthesis of monodisperse *ortho*-phenylene oligomers,<sup>37-45</sup> and under favorable conditions helical folding was observed as a consequence of arene-arene interactions.<sup>40,46-50</sup> Furthermore, oligo-*ortho*-phenylenes were found to exhibit remarkably long effective conjugation lengths.<sup>37,51</sup> Against this background and given the role of the respective *para*-isomers as prototype molecular wires it seemed attractive to explore long-range electron transfer in oligo-*ortho*-phenylenes.


## Results and Discussion

We synthesized a series of compounds comprised of a triarylamine (TAA) donor, variable-length *ortho*-phenylene bridges, and a Ru(bpy)<sub>3</sub><sup>2+</sup> (bpy = 2,2'-bipyridine) acceptor (Figure 1), relying on



**Figure 1.** *Para*-phenylene (a) and *ortho*-phenylene (b) bridged donor-acceptor (D-A) dyads, along with the molecular structure of the specific donor-bridge-acceptor compounds synthesized and investigated in this work (c).

recently reported methodologies for modular bridge elongation (see Supporting Information (SI) pages S4-S24).<sup>37</sup> The TAA oxidation and  $\text{Ru}(\text{bpy})_3^{2+}$  reduction potentials are essentially invariant with increasing bridge length (SI pages S25-S27), and on the basis of the cyclic voltammograms of our dyads we estimate a reaction free energy of ca. -0.4 eV for electron transfer from TAA to  $^3\text{MLCT}$ -excited  $\text{Ru}(\text{bpy})_3^{2+}$  in  $\text{CH}_3\text{CN}$  for all 3 compounds (SI page S28-S29). Expectedly, the common  $^3\text{MLCT}$  luminescence detectable in the  $\text{Ru}(\text{bpy})_3^{2+}$  parent complex is quenched when this photosensitizer unit is integrated into our donor-bridge-acceptor compounds (SI page S30). The photoproducts forming as a result of non-radiative  $^3\text{MLCT}$  deactivation can readily be identified as  $\text{TAA}^+$  and  $\text{Ru}(\text{bpy})_3^+$  based on transient absorption spectroscopy (Figure 2a) combined with spectro-electrochemistry (Figure 2b). Following excitation of 20  $\mu\text{M}$  solutions of the dyads from Figure 1c in deaerated  $\text{CH}_3\text{CN}$  at 532 nm using laser pulses of ca. 10 ns duration,



**Figure 2.** (a) Transient absorption spectra recorded immediately after excitation of the 3 compounds from Figure 1c at 532 nm with laser pulses of ca. 10 ns duration. The spectra were time-integrated over an interval of 200 ns. (b) Spectro-electrochemical UV-Vis difference spectra obtained after selective oxidation of the TAA unit (at a potential of 0.8 V vs. SCE, orange trace) or selective reduction of the  $\text{Ru}(\text{bpy})_3^{2+}$  acceptor (at a potential of -1.2 V vs. SCE, green trace) in the compound with  $n = 1$ . The solvent was de-aerated  $\text{CH}_3\text{CN}$  at 22 °C, the sample concentrations were 20  $\mu\text{M}$  for transient absorption and 205  $\mu\text{M}$  with 0.1 M  $\text{TBAPF}_6$  for spectro-electrochemistry.

transient absorption bands with maxima at 370, 510, and 750 nm appear in all 3 compounds (Figure 2a). Selective oxidation of the TAA unit at a potential of 0.8 V vs. SCE shows that the formation of  $\text{TAA}^+$  causes bands at 370 and 750 nm (orange trace in Figure 2b), whilst reduction of  $\text{Ru}(\text{bpy})_3^{2+}$  to  $\text{Ru}(\text{bpy})_3^+$  at -1.2 V vs. SCE leads to absorptions with maxima at 330 and 510


nm (green trace in Figure 2b). The black trace in Figure 2b is a 1:1 superposition of the TAA<sup>+</sup> and Ru(bpy)<sub>3</sub><sup>+</sup> difference spectra, and this linear combination is in excellent agreement with the transient absorption data in Figure 2a. On this short timescale (< 200 ns) at such low sample concentration (20 μM), the TAA<sup>+</sup> and Ru(bpy)<sub>3</sub><sup>+</sup> photoproducts can only form via intramolecular electron transfer, as confirmed by the kinetic studies below.

**Table 1.** Time constants for intramolecular photoinduced electron transfer from TAA to <sup>3</sup>MLCT-excited Ru(bpy)<sub>3</sub><sup>2+</sup> (τ<sub>f1</sub>, τ<sub>f2</sub>) and thermal reverse electron transfer from Ru(bpy)<sub>3</sub><sup>+</sup> to TAA<sup>+</sup> (τ<sub>b</sub>) in the compounds from Figure 1c.<sup>a</sup>

n	τ <sub>f1</sub> / ps	τ <sub>f2</sub> / ps	τ <sub>b</sub> / ns
1	110 (51%)	870 (49%)	120
2	220 (43%)	1600 (57%)	170
3	320 (62%)	6200 (38%)	220

<sup>a</sup> In de-aerated CH<sub>3</sub>CN at room temperature. Determined from the data in Figure 3 and Figure S10. Relative importance given in parentheses.


Picosecond transient absorption measurements reveal that the TAA<sup>+</sup> signal at 750 nm rises in bi-exponential fashion with time constants ranging from 110 to 6200 ps (Figure 3, τ<sub>f1</sub> and τ<sub>f2</sub> values in Table 1). In related donor-acceptor dyads with oligo-*p*-xylene bridges we previously observed strictly mono-exponential transient absorption kinetics.<sup>33,52-55</sup> Given the conformational flexibility of our oligo-*ortho*-phenylene bridges, it seems plausible that the bi-exponential nature of the transients in Figure 2 is due to the presence of slowly interconverting conformers. This interpretation is supported by variable-temperature NMR experiments, which demonstrate that



**Figure 3.** Temporal evolution of the transient absorption signal at 750 nm (red traces) of the three dyads following excitation at 532 nm with laser pulses of ca. 30 ps duration. Measurements were performed in de-aerated  $\text{CH}_3\text{CN}$  at 22 °C. Black curves are bi-exponential fits to the data, yielding the time constants reported in Table 1 ( $\tau_{f1}$  and  $\tau_{f2}$  values).

for the dyad with  $n = 2$  there is a barrier of ca. 60 kJ/mol for interconversion between two different main sets of conformers (SI pages S35-S37). Assuming first-order reaction kinetics for interconversion over a barrier of 60 kJ/mol height, a half-life for conformational exchange of  $\sim 3$

ms can be estimated with the Eyring equation (SI page S37). This is considerably longer than the timescale of the photoinduced electron transfer events in our dyads, and consequently the respective conformational exchange cannot occur during the intramolecular charge transfer processes studied herein. In the transient absorption studies, both sets of conformers are excited and both of them subsequently exhibit different electron transfer kinetics, leading to the observable bi-exponential transient absorption kinetics (Figure 3).



**Figure 4.** Key torsion angles in an *ortho*-phenylene hexamer.<sup>47</sup> A-type conformation:  $\phi_i = -55^\circ$  (“folded”); B-type conformation:  $\phi_i = +133^\circ$  (“unfolded”). The key conformers in Figure 5 and Table 2 are “A” or “B” for  $n = 1$ , “AA” or “BB” for  $n = 2$ , and “AAA” or “BBB” for  $n = 3$ .


In order to identify the key conformers of our dyads, we performed DFT calculations using Gaussian 09. The diffusion-corrected B97-D functional, which has previously been used to study the folding of *ortho*-phenylenes,<sup>47</sup> was used in combination with the 6-31G(d) basis set (LANL2DZ ECP for Ru atom) for structure optimization together with the IEFPCM/UFF model for  $\text{CH}_3\text{CN}$  solvent. There is considerable conformational flexibility in oligo-*ortho*-phenylene systems, and a complete conformational analysis is beyond our scope. As noted earlier, in a simple *ortho*-phenylene hexamer the backbone conformation is dictated by three key dihedral angles ( $\phi_2$ ,  $\phi_3$ ,  $\phi_4$ ) as illustrated in Figure 4.<sup>47</sup> Within a single *ortho*-phenylene strand each one of

**Table 2.** Relative energies ( $E_{\text{rel}}$ ) of the individual conformers from Figure 5 and calculated distances ( $d_{\text{N-Ru}}$ ) between the triarylamine N-atom and the Ru-center in the initial ground state (prior to electron transfer) and the photoproduct triplet state (formed as a result of intramolecular electron transfer; relevant structures shown in Figure S16).<sup>a</sup>

n	ground state				photoproduct triplet state			
	unfolded		folded		unfolded		folded	
	$E_{\text{rel}}$ / kJ/mol	$d_{\text{N-Ru}}$ / Å						
1	0 (B)	11.65	-36 (A)	7.29	0 (B)	13.30	-15 (A)	7.36
2	0 (BB)	9.25	-38 (AA)	8.18	0 (BB)	9.60	-42 (AA)	7.93
3	0 (BBB)	7.64	-17 (AAA)	15.91	0 (BBB)	7.36	-36 (AAA)	15.74

<sup>a</sup> Conformer types in notation used in main text indicated in parentheses. N-Ru distances measured through space.

these torsion angles ( $\phi_2$ ,  $\phi_3$ ,  $\phi_4$ ) can adopt one of two equilibrium values, referred to earlier as “A”-state (for  $\phi_i = -55^\circ$ ) or “B”-state (for  $\phi_i = +133^\circ$ ).<sup>47</sup> Our dyad with  $n = 3$  contains this hexameric structure as a molecular bridge, and in our calculations we concentrated on its AAA and BBB conformers. In principle, there are four additional conformers (ABA, BAB, AAB, and ABB) for this particular backbone, but they were not considered because they lead to excessive steric strain in our dyads with covalently attached triarylamine and  $\text{Ru}(\text{bpy})_3^{2+}$  units. By analogy, for the two shorter dyads only the AA and BB conformers (for  $n = 2$ ), as well as the A and B conformers (for  $n = 1$ ) were considered. In the AAA and AA conformers (Figure 5f, d) there are



**Figure 5.** Key ground state conformer geometries of the *ortho*-phenylene bridged dyads: a)  $n = 1$ , unfolded B-form; b)  $n = 1$ , folded A-form; c)  $n = 2$ , unfolded BB-form; d)  $n = 2$ , folded AA-form; e)  $n = 3$ , unfolded BBB-form; f)  $n = 3$ , folded AAA-form. See SI page S38-S42 for details.

arene-arene interactions leading to the folded structures reported earlier for oligo-*ortho*-phenylenes, whereas the BBB and BB conformers (Figure 5e, c) can be considered unfolded structures without such arene-arene interactions. These calculations were performed for the dyads

10

1  
2  
3 in their electronic ground state (Figure 5) as well as in their photoproduct states with oxidized  
4 triarylamine and reduced ruthenium photosensitizer (Figure S16). The AAA, AA, and A  
5 conformers are the most stable structures according to these calculations, lower in energy than the  
6  
7  
8 conformers are the most stable structures according to these calculations, lower in energy than the  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

in their electronic ground state (Figure 5) as well as in their photoproduct states with oxidized triarylamine and reduced ruthenium photosensitizer (Figure S16). The AAA, AA, and A conformers are the most stable structures according to these calculations, lower in energy than the BBB, BB, and B conformers by 15 to 42 kJ/mol (Table 2). These two different types of conformers could in fact correspond to the two sets of conformers observed in the variable-temperature NMR experiments.

The distance between the triarylamine N-atom and the Ru atom is a useful proxy for the electron transfer distance in our dyads, and that distance (measured through space) is markedly different for different conformers of a given compound. For example, in the dyad with  $n = 3$  the calculated N-Ru distance in the AAA conformer is 15.9 Å but in the BBB conformer it is only 7.6 Å (Table 2, Figure 5e, f). Such a marked difference in donor-acceptor distance can readily account for the bi-exponential transient absorption kinetics in Figure 3. However, it should be kept in mind that minor conformational changes are easily possible on the timescale of our electron transfer events, and it is not possible to attribute a certain measurable time constant to any specific computed conformer structure,<sup>56</sup> in particular because the observable rate constants could in fact correspond to a combination of rate constants for electron transfer and conformational changes.<sup>57,58</sup> Thus, our calculations primarily provide a qualitative rationale for the multi-exponential transient absorption kinetics, and they illustrate the difficulty of assessing a distance-dependence for electron transfer in the dyads from Figure 1c. Different conformers of the same compound can exhibit very different donor-acceptor distances, depending on whether the conformers are more U-shaped with donor and acceptor pointing into roughly the same direction (Figure 5e,  $d_{\text{N-Ru}} \approx 8$  Å) or whether they are more Z-shaped with donor and acceptor pointing towards opposite directions (Figure 5f,  $d_{\text{N-Ru}} \approx 16$  Å).

1  
2  
3 Interestingly, the thermal reverse charge-shift from  $\text{Ru}(\text{bpy})_3^+$  to  $\text{TAA}^+$  gives rise to single-  
4 exponential transient absorption kinetics (SI page S32). The time constants for intramolecular  
5 backward electron transfer from  $\text{Ru}(\text{bpy})_3^+$  to  $\text{TAA}^+$  ( $\tau_b$ , Table 1) were determined from the  
6 decays of the transient absorption signals at 370, 510 and 750 nm (Figure S10). In all three dyads,  
7 the  $\text{Ru}(\text{bpy})_3^+ / \text{TAA}^+$  photoproduct state decays much more slowly than it forms, and the  
8 lifetimes  $\tau_b$  vary only over the small range between 120 and 220 ns despite the very different  
9 lengths of the three dyads. As noted above, based on variable-temperature NMR studies it is to be  
10 expected that two sets of slowly interconverting ( $\sim 3$  ms) conformers are initially excited,  
11 manifesting in bi-exponential kinetics for the initial photoinduced electron transfer from TAA to  
12  $^3\text{MLCT}$ -excited  $\text{Ru}(\text{bpy})_3^{2+}$  (Figure 3). At first glance, the observation of mono-exponential  
13 kinetics for the ensuing thermal reverse charge-shift therefore looks peculiar. However, it seems  
14 plausible that in one of the two key conformer sets of each dyad the  $\text{Ru}(\text{bpy})_3^+ / \text{TAA}^+$   
15 photoproduct disappears more rapidly than it is formed due to close donor-acceptor contacts,  
16 making its detection impossible for these specific conformers. Indeed, some of the calculated  
17 conformers for the  $\text{Ru}(\text{bpy})_3^+ / \text{TAA}^+$  photoproduct state exhibit N-Ru distances below 8 Å  
18 (Table 2), essentially corresponding to van-der-Waals contact distance for which very rapid  
19 reverse charge-shift is expected. However, there are other low-energy conformers for which the  
20 N-Ru distance is up to ca. 16 Å (Table 2), compatible with electron transfer kinetics on the 120 –  
21 220 ns timescale.

22  
23 In *para*-phenylene-bridged donor-acceptor systems, charges usually transfer along the  $\sigma$ -bonded  
24 backbone benefiting from extended  $\pi$ -conjugation, whereas through-space or through-solvent  
25 pathways play no significant role due to their linear rigid rod-like structure.<sup>20,22,29,33,36,52-54,59,60</sup> By  
26 contrast, in our conformationally flexible *ortho*-phenylene dyads through-bond, through-space or  
27

1  
2  
3 through-solvent pathways are in principle all viable,<sup>61,62</sup> and their relative contributions to the  
4 overall electron transfer kinetics cannot be disentangled from the available experimental and  
5 computational data. Consequently, the true distance dependence for electron transfer in our dyads  
6 cannot be determined, despite the variation of the number of phenylene units from 2 to 6 (Figure  
7 1c).  
8  
9

10  
11  
12  
13  
14 As noted in the introduction, oligo-*ortho*-phenylenes can form folded structures with arene-arene  
15 interactions,<sup>40,46-50</sup> and the question whether photoinduced electron transfer could be observed in  
16 such folded structures seemed interesting, particularly because electron transfer across artificial  
17 foldamers is yet relatively little explored.<sup>63-65</sup> The DFT calculations performed for the compounds  
18 from Figure 1c indeed suggest that conformers with phenylene-phenylene interactions (AAA and  
19 AA conformers, Figure 5f, d) are significantly more stable than conformers lacking such  
20 interactions (BBB and BB conformers, Table 2). For instance, for the dyad with  $n = 3$  the folded  
21 AAA-conformer is 36 kJ/mol more stable than the unfolded BBB-conformer in the photoproduct  
22 triplet state, and the calculated N-Ru distance of 15.7 Å is compatible with a time constant of 220  
23 ns for thermal charge-shift between  $\text{Ru}(\text{bpy})_3^+$  and  $\text{TAA}^+$  (Table 2). In passing we note that the  
24 additional electron on the reduced ruthenium complex should be largely localized on the electron-  
25 deficient ester-functionalized bpy ligand rather than the metal center, yet the N-Ru distance  
26 remains a useful proxy for the electron transfer distance. In case charge transfer across arene-  
27 arene interactions in foldamer structures would indeed play a significant role, then one might  
28 expect electron transfer rates to increase with decreasing temperature, due to increasing  
29 stabilization of the arene-arene contacts.<sup>66,67</sup> However, opposite behavior is observed (SI page  
30 S34), leading to ordinary Arrhenius-plots providing activation energies between 48 and 98 meV,  
31 in line with prior studies of *para*-phenylene based dyads that cannot fold.<sup>33,54</sup> Thus, it seems that  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

1  
2  
3 if folded structures are indeed involved (as suggested by the computational results, see above),  
4 then the dominant charge transfer pathway is unlikely to involve the types of arene-arene  
5 interactions that cause the folding.  
6  
7  
8

9  
10 Regarding charge transfer along the  $\sigma$ -bonded backbone of our *ortho*-phenylene bridged dyads,  
11 the presence of methoxy-substituents deserves some attention. These electron-donating  
12 substituents lower the potential corresponding to bridge oxidation markedly, and consequently  
13 several prior studies on methoxy-decorated *para*-phenylene-bridged systems reported evidence  
14 for hole hopping.<sup>68-71</sup> We therefore synthesized a series of oligo-*ortho*-phenylene compounds  
15 without appended Ru(bpy)<sub>3</sub><sup>2+</sup> and TAA units and determined the potentials for their oxidation  
16 using cyclic voltammetry (SI page S26-S27). In a bridge model compound made from only two  
17 methoxy-substituted phenylene units, the first oxidation wave occurs at 1.70 V vs. SCE, whereas  
18 in longer congeners made from four and six methoxy-decorated *ortho*-phenylenes the first  
19 oxidation waves peak at 1.23 and 1.25 V vs. SCE, respectively. In the dyads from Figure 1c,  
20 TAA<sup>+</sup> reduction takes place at 0.60 V vs. SCE (SI pages S25-S27), and consequently hole  
21 transfer from TAA<sup>+</sup> to the bridge would be energetically uphill by 0.63 – 1.10 eV since the hole  
22 hopping mechanism implies temporary oxidation of the bridge.<sup>72,73</sup> Electron hopping is similarly  
23 unlikely, because the reduction potential of our bridges is more negative than -2.0 V vs. SCE,  
24 whereas Ru(bpy)<sub>3</sub><sup>+</sup> is oxidized at -1.2 V vs. SCE (SI pages S25-S26). Thus, we conclude that  
25 hopping mechanisms are unlikely to play an important role for the thermal charge-shift reaction  
26 in our dyads. The situation might be different in *para*-connected analogues,<sup>59</sup> but their  
27 investigation is beyond our scope here. In a direct comparison with *para*-connected isomers, the  
28 distance dependence of quantum yields for formation of charge-separated states would be an  
29 interesting aspect.<sup>74</sup>  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Conclusions

Conformationally flexible *ortho*-phenylene bridges between a triarylamine donor and a Ru(bpy)<sub>3</sub><sup>2+</sup> acceptor permit rapid (<10 ns) photoinduced electron transfer even when up to six phenylene bridging units are present. The resulting photoproduct state comprised of oxidized triarylamine and reduced ruthenium complex stores ca. 1.7 eV of energy, and its lifetime is only weakly dependent on the number of phenylene bridging units, ranging from 120 ns for the compound with only two phenylene bridge units to 220 ns for the dyad with six phenylenes in de-aerated CH<sub>3</sub>CN at room temperature. Given the conformational flexibility of these dyads enabling through-space and through-solvent pathways in addition to the through-bond pathways known from oligo-*para*-phenylenes, these photoproduct lifetimes are remarkably long. This shows that rigid rod-like molecular structures are not a key requirement for obtaining long-lived electron-hole separation.

Whilst folded conformers with arene-arene interactions seem to be formed in the donor-bridge-acceptor compounds studied herein, no direct evidence for charge transfer pathways involving the arene-arene interactions leading to foldamer structures could be found.

The distance dependence of charge (and energy) transfer across oligo-*para*-phenylenes has been investigated thoroughly in the past,<sup>22,23,25,26,33,52-54,60,70,73,75-79</sup> but similarly thorough insight into the distance dependence of electron transfer across oligo-*ortho*-phenylenes cannot be obtained from our study due to the many conformational degrees of freedom available in our compounds. Future studies might therefore focus on oligo-*ortho*-naphthalene connected systems with less conformational flexibility,<sup>80-83</sup> in which it might be more readily possible to enforce through-

1  
2  
3 bond electron transfer. Photoinduced electron transfer across naphthalene-bridged systems has  
4  
5 been explored relatively little until now.<sup>84,85</sup>  
6  
7  
8  
9

## 10 **Supporting Information Description**

11  
12  
13  
14 Detailed synthetic protocols and characterization data, description of equipment and methods,  
15  
16 supplementary electrochemical, spectroscopic, and computational data.  
17  
18  
19  
20

## 21 **Acknowledgments**

22  
23  
24  
25  
26 Funding by the Swiss National Science Foundation through grant numbers 200021\_178760/1 and  
27  
28 206021\_157687 is gratefully acknowledged.  
29  
30  
31  
32

## 33 **References**

- 34  
35  
36  
37 (1) Winkler, J. R.; Gray, H. B. Long-Range Electron Tunneling. *J. Am. Chem. Soc.*  
38 **2014**, *136*, 2930-2939.  
39  
40  
41 (2) Arrigo, A.; Santoro, A.; Puntoriero, F.; Lainé, P. P.; Campagna, S. Photoinduced  
42  
43 Electron Transfer in Donor-Bridge-Acceptor Assemblies: The Case of Os(II)-Bis(terpyridine)-  
44  
45 (Bi)pyridinium Dyads. *Coord. Chem. Rev.* **2015**, *304*, 109-116.  
46  
47  
48 (3) Lim, G. N.; Obondi, C. O.; D'Souza, F. A High-Energy Charge-Separated State of  
49  
50 1.70 eV from a High-Potential Donor-Acceptor Dyad: A Catalyst for Energy-Demanding  
51  
52 Photochemical Reactions. *Angew. Chem. Int. Ed.* **2016**, *55*, 11517-11521.  
53  
54  
55  
56  
57  
58  
59  
60

(4) Schroot, R.; Schlotthauer, T.; Dietzek, B.; Jäger, M.; Schubert, U. S. Extending Long-lived Charge Separation Between Donor and Acceptor Blocks in Novel Copolymer Architectures Featuring a Sensitizer Core. *Chem. Eur. J.* **2017**, *23*, 16484-16490.

(5) Yamamoto, M.; Föhlinger, J.; Petersson, J.; Hammarström, L.; Imahori, H. A Ruthenium Complex-Porphyrin-Fullerene-Linked Molecular Pentad as an Integrative Photosynthetic Model. *Angew. Chem. Int. Ed.* **2017**, *56*, 3329-3333.

(6) Mendes Marinho, S.; Ha-Thi, M.-H.; Pham, V.-T.; Quaranta, A.; Pino, T.; Lefumeux, C.; Chamailé, T.; Leibl, W.; Aukauloo, A. Time-Resolved Interception of Multi-Charge Accumulation in a Sensitizer-Acceptor Dyad. *Angew. Chem. Int. Ed.* **2017**, *56*, 15936–15940.

(7) Delor, M.; Archer, S. A.; Keane, T.; Meijer, A.; Sazanovich, I. V.; Greetham, G. M.; Towrie, M.; Weinstein, J. A. Directing the Path of Light-induced Electron Transfer at a Molecular Fork Using Vibrational Excitation. *Nat. Chem.* **2017**, *9*, 1099-1104.

(8) Pannwitz, A.; Wenger, O. S. Photoinduced Electron Transfer Coupled to Donor Deprotonation and Acceptor Protonation in a Molecular Triad Mimicking Photosystem II. *J. Am. Chem. Soc.* **2017**, *139*, 13308-13311.

(9) Majewski, M. B.; de Tacconi, N. R.; MacDonnell, F. M.; Wolf, M. O. Long-Lived, Directional Photoinduced Charge Separation in Ru<sup>II</sup> Complexes Bearing Laminate Polypyridyl Ligands. *Chem.-Eur. J.* **2013**, *19*, 8331-8341.

(10) Willa, A. R.; Philip, A. M.; Bhat, V.; Hariharan, M. Persistent Charge-Separated States in Self-Assembled Twisted Nonsymmetric Donor-Acceptor Triads. *J. Phys. Chem. C* **2017**, *121*, 4765-4777.

1  
2  
3 (11) Lloveras, V.; Vidal-Gancedo, J.; Figueira-Duarte, T. M.; Nierengarten, J. F.;  
4  
5 Novoa, J. J.; Mota, F.; Ventosa, N.; Rovira, C.; Veciana, J. Tunneling versus Hopping in Mixed-  
6  
7 Valence Oligo-p-phenylenevinylene Polychlorinated Bis(triphenylmethyl) Radical Anions. *J.*  
8  
9 *Am. Chem. Soc.* **2011**, *133*, 5818-5833.

10  
11 (12) Sukegawa, J.; Schubert, C.; Zhu, X. Z.; Tsuji, H.; Guldi, D. M.; Nakamura, E.  
12  
13 Electron Transfer Through Rigid Organic Molecular Wires Enhanced by Electronic and Electron-  
14  
15 Vibration Coupling. *Nat. Chem.* **2014**, *6*, 899-905.

16  
17 (13) Linton, K. E.; Fox, M. A.; Palsson, L. O.; Bryce, M. R. Oligo(*p*-  
18  
19 Phenyleneethynylene) (OPE) Molecular Wires: Synthesis and Length Dependence of  
20  
21 Photoinduced Charge Transfer in OPEs with Triarylamine and Diaryloxadiazole End Groups.  
22  
23 *Chem.-Eur. J.* **2015**, *21*, 3997-4007.

24  
25 (14) Waskasi, M. M.; Kodis, G.; Moore, A. L.; Moore, T. A.; Gust, D.; Matyushov, D.  
26  
27 V. Marcus Bell-Shaped Electron Transfer Kinetics Observed in an Arrhenius Plot. *J. Am. Chem.*  
28  
29 *Soc.* **2016**, *138*, 9251-9257.

30  
31 (15) Krzeszewski, M.; Espinoza, E. M.; Cervinka, C.; Derr, J. B.; Clark, J. A.;  
32  
33 Borchardt, D.; Beran, G. J. O.; Gryko, D. T.; Vullev, V. I. Dipole Effects on Electron Transfer  
34  
35 are Enormous. *Angew. Chem. Int. Ed.* **2018**, doi: 10.1002/anie.201802637.

36  
37 (16) Yonemoto, E. H.; Saupe, G. B.; Schmechl, R. H.; Hubig, S. M.; Riley, R. L.;  
38  
39 Iverson, B. L.; Mallouk, T. E. Electron-Transfer Reactions of Ruthenium Trisbipyridyl-Viologen  
40  
41 Donor-Acceptor Molecules - Comparison of the Distance Dependence of Electron-Transfer Rates  
42  
43 in the Normal and Marcus Inverted Regions. *J. Am. Chem. Soc.* **1994**, *116*, 4786-4795.

44  
45 (17) Barbara, P. F.; Meyer, T. J.; Ratner, M. A. Contemporary Issues in Electron  
46  
47 Transfer Research. *J. Phys. Chem.* **1996**, *100*, 13148-13168.

1  
2  
3 (18) Rosspeintner, A.; Angulo, G.; Vauthey, E. Bimolecular Photoinduced Electron  
4  
5 Transfer Beyond the Diffusion Limit: The Rehm-Weller Experiment Revisited with Femtosecond  
6  
7 Time Resolution. *J. Am. Chem. Soc.* **2014**, *136*, 2026-2032.

8  
9  
10 (19) Wiberg, J.; Guo, L. J.; Pettersson, K.; Nilsson, D.; Ljungdahl, T.; Mårtensson, J.;  
11  
12 Albinsson, B. Charge Recombination versus Charge Separation in Donor-Bridge-Acceptor  
13  
14 Systems. *J. Am. Chem. Soc.* **2007**, *129*, 155-163.

15  
16  
17 (20) Helms, A.; Heiler, D.; McLendon, G. Electron-Transfer in Bis-Porphyrin Donor-  
18  
19 Acceptor Compounds with Polyphenylene Spacers Shows a Weak Distance Dependence. *J. Am.*  
20  
21 *Chem. Soc.* **1992**, *114*, 6227-6238.

22  
23  
24 (21) Rosokha, S. V.; Sun, D.-L.; Kochi, J. K. Conformation, Distance, and  
25  
26 Connectivity Effects on Intramolecular Electron Transfer between Phenylene-Bridged Aromatic  
27  
28 Redox Centers. *J. Phys. Chem. A* **2002**, *106*, 2283-2292.

29  
30  
31 (22) Weiss, E. A.; Ahrens, M. J.; Sinks, L. E.; Gusev, A. V.; Ratner, M. A.;  
32  
33 Wasielewski, M. R. Making a Molecular Wire: Charge and Spin Transport through Para-  
34  
35 Phenylene Oligomers. *J. Am. Chem. Soc.* **2004**, *126*, 5577-5584.

36  
37  
38 (23) Welter, S.; Salluce, N.; Belser, P.; Groeneveld, M.; De Cola, L. Photoinduced  
39  
40 Electronic Energy Transfer in Modular, Conjugated, Dinuclear Ru(II) / Os(II) Complexes. *Coord.*  
41  
42 *Chem. Rev.* **2005**, *249*, 1360-1371.

43  
44  
45 (24) Barlow, S.; Risko, C.; Chung, S. J.; Tucker, N. M.; Coropceanu, V.; Jones, S. C.;  
46  
47 Levi, Z.; Brédas, J. L.; Marder, S. R. Intervalence Transitions in the Mixed-Valence Monocations  
48  
49 of Bis(triarylamines) Linked with Vinylene and Phenylene-Vinylene Bridges. *J. Am. Chem. Soc.*  
50  
51 **2005**, *127*, 16900-16911.

1  
2  
3 (25) Indelli, M. T.; Chiorboli, C.; Flamigni, L.; De Cola, L.; Scandola, F. Photoinduced  
4 Electron Transfer across Oligo-*p*-phenylene Bridges. Distance and Conformational Effects in  
5 Ru(II)-Rh(III) Dyads. *Inorg. Chem.* **2007**, *46*, 5630-5641.  
6  
7

8  
9  
10 (26) Wenger, O. S. Photoinduced Electron and Energy Transfer in Phenylene  
11 Oligomers. *Chem. Soc. Rev.* **2011**, *40*, 3538-3550.  
12  
13

14 (27) Warren, J. J.; Menzeleev, A. R.; Kretchmer, J. S.; Miller, T. F.; Gray, H. B.;  
15 Mayer, J. M. Long-Range Proton-Coupled Electron-Transfer Reactions of Bis(imidazole) Iron  
16 Tetraphenylporphyrins Linked to Benzoates. *J. Phys. Chem. Lett.* **2013**, *4*, 519-523.  
17  
18

19 (28) Gorczak, N.; Renaud, N.; Tarkuc, S.; Houtepen, A. J.; Eelkema, R.; Siebbeles, L.  
20 D. A.; Grozema, F. C. Charge Transfer versus Molecular Conductance: Molecular Orbital  
21 Symmetry Turns Quantum Interference Rules Upside Down. *Chem. Sci.* **2015**, *6*, 4196-4206.  
22  
23

24 (29) Shen, J. J.; Zhong, Y. W. Long-Range Ruthenium-Amine Electronic  
25 Communication through the *para*-Oligophenylene Wire. *Sci Rep* **2015**, *5*, doi:  
26 10.1038/srep13835.  
27  
28

29 (30) Stasiw, D. E.; Zhang, J. Y.; Wang, G. B.; Dangi, R.; Stein, B. W.; Shultz, D. A.;  
30 Kirk, M. L.; Wojtas, L.; Sommer, R. D. Determining the Conformational Landscape of *s* and *p*  
31 Coupling Using *para*-Phenylene and "Aviram-Ratner" Bridges. *J. Am. Chem. Soc.* **2015**, *137*,  
32 9222-9225.  
33  
34

35 (31) Klein, J. H.; Schmidt, D.; Steiner, U. E.; Lambert, C. Complete Monitoring of  
36 Coherent and Incoherent Spin Flip Domains in the Recombination of Charge-Separated States of  
37 Donor-Iridium Complex-Acceptor Triads. *J. Am. Chem. Soc.* **2015**, *137*, 11011-11021.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

(32) Hu, K.; Blair, A. D.; Piechota, E. J.; Schauer, P. A.; Sampaio, R. N.; Parlane, F. G. L.; Meyer, G. J.; Berlinguette, C. P. Kinetic Pathway for Interfacial Electron Transfer from a Semiconductor to a Molecule. *Nat. Chem.* **2016**, *8*, 853-859.

(33) Kuss-Petermann, M.; Wenger, O. S. Electron Transfer Rate Maxima at Large Donor-Acceptor Distances. *J. Am. Chem. Soc.* **2016**, *138*, 1349-1358.

(34) Xu, Y.; Wang, B.; Kaur, R.; Minameyer, M. B.; Bothe, M.; Drewello, T.; Guldi, D. M.; von Delius, M. A Supramolecular [10]CPP Junction Enables Efficient Electron Transfer in Modular Porphyrin-[10]CPP Fullerene Complexes. *Angew. Chem. Int. Ed.* **2018**, doi: 10.1002/anie.201802443.

(35) Bauer, C.; Teuscher, J.; Brauer, J. C.; Punzi, A.; Marchioro, A.; Ghadiri, E.; De Jonghe, J.; Wielopolski, M.; Banerji, N.; Moser, J. E. Dynamics and Mechanisms of Interfacial Photoinduced Electron Transfer Processes of Third Generation Photovoltaics and Photocatalysis. *Chimia* **2011**, *65*, 704-709.

(36) Benniston, A. C.; Harriman, A. Charge on the Move: How Electron-Transfer Dynamics Depend on Molecular Conformation. *Chem. Soc. Rev.* **2006**, *35*, 169-179.

(37) He, J. A.; Crase, J. L.; Wadumethrige, S. H.; Thakur, K.; Dai, L.; Zou, S. Z.; Rathore, R.; Hartley, C. S. Ortho-Phenylenes: Unusual Conjugated Oligomers with a Surprisingly Long Effective Conjugation Length. *J. Am. Chem. Soc.* **2010**, *132*, 13848-13857.

(38) Ohta, E.; Sato, H.; Ando, S.; Kosaka, A.; Fukushima, T.; Hashizume, D.; Yamasaki, M.; Hasegawa, K.; Muraoka, A.; Ushiyama, H.; Yamashita, K.; Aida, T. Redox-Responsive Molecular Helices with Highly Condensed p-Clouds. *Nat. Chem.* **2011**, *3*, 68-73.

(39) Mathew, S. M.; Hartley, C. S. Parent o-Phenylene Oligomers: Synthesis, Conformational Behavior, and Characterization. *Macromolecules* **2011**, *44*, 8425-8432.

1  
2  
3 (40) Kajitani, T.; Suna, Y.; Kosaka, A.; Osawa, T.; Fujikawa, S.; Takata, M.;  
4  
5 Fukushima, T.; Aida, T. o-Phenylene Octamers as Surface Modifiers for Homeotropic Columnar  
6  
7 Ordering of Discotic Liquid Crystals. *J. Am. Chem. Soc.* **2013**, *135*, 14564-14567.  
8

9  
10 (41) Koyanagi, M.; Eichenauer, N.; Ihara, H.; Yamamoto, T.; Suginome, M.  
11  
12 Anthranilamide-masked o-Iodoarylboronic Acids as Coupling Modules for Iterative Synthesis of  
13  
14 ortho-Linked Oligoarenes. *Chem. Lett.* **2013**, *42*, 541-543.  
15

16  
17 (42) Ito, S. G.; Takahashi, K.; Nozaki, K. Formal Aryne Polymerization: Use of [2.2.1]  
18  
19 Oxabicyclic Alkenes as Aryne Equivalents. *J. Am. Chem. Soc.* **2014**, *136*, 7547-7550.  
20

21 (43) Mizukoshi, Y.; Mikami, K.; Uchiyama, M. Aryne Polymerization Enabling  
22  
23 Straightforward Synthesis of Elusive Poly(ortho-arylene)s. *J. Am. Chem. Soc.* **2015**, *137*, 74-77.  
24

25  
26 (44) Lehnherr, D.; Chen, C.; Pedramrazi, Z.; DeBlase, C. R.; Alzola, J. M.; Keresztes,  
27  
28 I.; Lobkovsky, E. B.; Crommie, M. F.; Dichtel, W. R. Sequence-Defined Oligo(ortho-arylene)  
29  
30 Foldamers Derived from the Benzannulation of Ortho(arylene ethynylene)s. *Chem. Sci.* **2016**, *7*,  
31  
32 6357-6364.  
33

34  
35 (45) Hammer, B. A. G.; Müllen, K. Dimensional Evolution of Polyphenylenes:  
36  
37 Expanding in All Directions. *Chem. Rev.* **2016**, *116*, 2103-2140.  
38

39  
40 (46) Ando, S.; Ohta, E.; Kosaka, A.; Hashizume, D.; Koshino, H.; Fukushima, T.;  
41  
42 Aida, T. Remarkable Effects of Terminal Groups and Solvents on Helical Folding of o-Phenylene  
43  
44 Oligomers. *J. Am. Chem. Soc.* **2012**, *134*, 11084-11087.  
45

46  
47 (47) Mathew, S. M.; Engle, J. T.; Ziegler, C. J.; Hartley, C. S. The Role of Arene-  
48  
49 Arene Interactions in the Folding of ortho-Phenylenes. *J. Am. Chem. Soc.* **2013**, *135*, 6714-6722.  
50

1  
2  
3 (48) Mathew, S.; Crandall, L. A.; Ziegler, C. J.; Hartley, C. S. Enhanced Helical  
4 Folding of ortho-Phenylenes through the Control of Aromatic Stacking Interactions. *J. Am.*  
5 *Chem. Soc.* **2014**, *136*, 16666-16675.

6  
7  
8  
9  
10 (49) Hartley, C. S. Folding of ortho-Phenylenes. *Acc. Chem. Res.* **2016**, *49*, 646-654.

11  
12 (50) Vemuri, G. N.; Chu, M.; Dong, H.; Spinello, B. J.; Hartley, C. S. Solvent Effects  
13 on the Folding of o-Phenylene Oligomers. *Org. Biomol. Chem.* **2017**, *15*, 845-851.

14  
15  
16 (51) Hartley, C. S. Excited-State Behavior of ortho-Phenylenes. *J. Org. Chem.* **2011**,  
17 *76*, 9188-9191.

18  
19  
20 (52) Hanss, D.; Wenger, O. S. Electron Tunneling Through Oligo-*p*-xylene Bridges.  
21 *Inorg. Chem.* **2008**, *47*, 9081-9084.

22  
23  
24 (53) Hanss, D.; Wenger, O. S. Tunneling Barrier Effects on Photoinduced Charge  
25 Transfer through Covalent Rigid Rod-like Bridges. *Inorg. Chem.* **2009**, *48*, 671-680.

26  
27  
28 (54) Kuss-Petermann, M.; Wenger, O. S. Increasing Electron-Transfer Rates with  
29 Increasing Donor-Acceptor Distance. *Angew. Chem. Int. Ed.* **2016**, *55*, 815-819.

30  
31 (55) Neumann, S.; Wenger, O. S. Fundamentally Different Distance Dependences of  
32 Electron-Transfer Rates for Low and High Driving Forces. *Inorg. Chem.* **2019**, doi:  
33 10.1021/acs.inorgchem.8b02973.

34  
35  
36 (56) Skourtis, S. S.; Waldeck, D. H.; Beratan, D. N. Fluctuations in Biological and  
37 Bioinspired Electron-Transfer Reactions. *Annu. Rev. Phys. Chem.* **2010**, *61*, 461-485.

38  
39  
40 (57) Druzhinin, S. I.; Ernsting, N. P.; Kovalenko, S. A.; Lustres, L. W.; Senyushkina,  
41 T. A.; Zachariasse, K. A. Dynamics of Ultrafast Intramolecular Charge Transfer with 4-  
42 (Dimethylamino)benzotrile in Acetonitrile. *J. Phys. Chem. A* **2006**, *110*, 2955-2969.

1  
2  
3 (58) Druzhinin, S. I.; Galievsky, V. A.; Demeter, A.; Kovalenko, S. A.; Senyushkina,  
4 T.; Dubbaka, S. R.; Knoche, P.; Mayer, P.; Grosse, C.; Stalke, D.; Zachariasse, K. A. Two-State  
5 Intramolecular Charge Transfer (ICT) with 3,5-Dimethyl-4(dimethylamino)benzotrile (MMD)  
6 and Its Meta-Isomer mMMD. Ground State Amino Twist Not Essential for ICT. *J. Phys. Chem.*  
7 *A* **2015**, *119*, 11820-11836.  
8  
9

10  
11  
12 (59) Weiss, E. A.; Tauber, M. J.; Kelley, R. F.; Ahrens, M. J.; Ratner, M. A.;  
13 Wasielewski, M. R. Conformationally Gated Switching between Superexchange and Hopping  
14 within Oligo-*p*-phenylene-Based Molecular Wires. *J. Am. Chem. Soc.* **2005**, *127*, 11842-11850.  
15  
16

17  
18 (60) Albinsson, B.; Eng, M. P.; Pettersson, K.; Winters, M. U. Electron and Energy  
19 Transfer in Donor-Acceptor Systems with Conjugated Molecular Bridges. *Phys. Chem. Chem.*  
20 *Phys.* **2007**, *9*, 5847-5864.  
21  
22

23  
24 (61) Graff, B. M.; Lamont, D. N.; Parker, M. F. L.; Bloom, B. P.; Schafmeister, C. E.;  
25 Waldeck, D. H. Through-Solvent Tunneling in Donor-Bridge-Acceptor Molecules Containing a  
26 Molecular Cleft. *J. Phys. Chem. A* **2016**, *120*, 6004-6013.  
27  
28

29  
30 (62) Schmidt, H. C.; Larsen, C. B.; Wenger, O. S. Electron Transfer around a  
31 Molecular Corner. *Angew. Chem. Int. Ed.* **2018**, *57*, 6696-6700.  
32  
33

34  
35 (63) Méndez-Ardoy, A.; Markandeya, N.; Li, X. S.; Tsai, Y. T.; Pecastaings, G.;  
36 Buffeteau, T.; Maurizot, V.; Muccioli, L.; Castet, F.; Huc, I.; Bassani, D. M. Multi-Dimensional  
37 Charge Transport in Supramolecular Helical Foldamer Assemblies. *Chem. Sci.* **2017**, *8*, 7251-  
38 7257.  
39  
40

41  
42 (64) Li, X. S.; Markandeya, N.; Jonusauskas, G.; McClenaghan, N. D.; Maurizot, V.;  
43 Denisov, S. A.; Huc, I. Photoinduced Electron Transfer and Hole Migration in Nanosized Helical  
44 Aromatic Oligoamide Foldamers. *J. Am. Chem. Soc.* **2016**, *138*, 13568-13578.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

(65) Wolffs, M.; Delsuc, N.; Veldman, D.; Van Anh, N.; Williams, R. M.; Meskers, S. C. J.; Janssen, R. A. J.; Huc, I.; Schenning, A. P. H. J. Helical Aromatic Oligoamide Foldamers as Organizational Scaffolds for Photoinduced Charge Transfer. *J. Am. Chem. Soc.* **2009**, *131*, 4819-4829.

(66) Fukuzumi, S.; Ohkubo, K.; Tokuda, Y.; Suenobu, T. Hydride Transfer from 9-Substituted 10-Methyl-9,10-dihydroacridines to Hydride Acceptors via Charge-transfer Complexes and Sequential Electron-Proton-Electron Transfer. A Negative Temperature Dependence of the Rates. *J. Am. Chem. Soc.* **2000**, *122*, 4286-4294.

(67) Larsen, C. B.; Wenger, O. S. Circular Photoinduced Electron Transfer in a Donor-Acceptor-Acceptor Triad. *Angew. Chem. Int. Ed.* **2018**, *57*, 841-845.

(68) Miller, S. E.; Lukas, A. S.; Marsh, E.; Bushard, P.; Wasielewski, M. R. Photoinduced Charge Separation Involving an Unusual Double Electron Transfer Mechanism in a Donor-Bridge-Acceptor Molecule. *J. Am. Chem. Soc.* **2000**, *122*, 7802-7810.

(69) Lambert, C.; Nöll, G.; Schelter, J. Bridge-Mediated Hopping or Superexchange Electron-Transfer Processes in Bis(Triarylamine) Systems. *Nat. Mater.* **2002**, *1*, 69-73.

(70) Walther, M. E.; Wenger, O. S. Tuning the Rates of Long-Range Charge Transfer across Phenylene Wires. *ChemPhysChem* **2009**, *10*, 1203-1206.

(71) Heinz, L. G.; Yushchenko, O.; Neuburger, M.; Vauthey, E.; Wenger, O. S. Tetramethoxybenzene is a Good Building Block for Molecular Wires: Insights from Photoinduced Electron Transfer. *J. Phys. Chem. A* **2015**, *119*, 5676-5684.

(72) Cordes, M.; Giese, B. Electron Transfer in Peptides and Proteins. *Chem. Soc. Rev.* **2009**, *38*, 892-901.

(73) Natali, M.; Campagna, S.; Scandola, F. Photoinduced Electron Transfer Across Molecular Bridges: Electron- and Hole-Transfer Superexchange Pathways. *Chem. Soc. Rev.* **2014**, *43*, 4005-4018.

(74) Higashino, T.; Yamada, T.; Yamamoto, M.; Furube, A.; Tkachenko, N. V.; Miura, T.; Kobori, Y.; Jono, R.; Yamashita, K.; Imahori, H. Remarkable Dependence of the Final Charge Separation Efficiency on the Donor-Acceptor Interaction in Photoinduced Electron Transfer. *Angew. Chem. Int. Ed.* **2016**, *55*, 629-633.

(75) Wenger, O. S. How Donor-Bridge-Acceptor Energetics Influence Electron Tunneling Dynamics and Their Distance Dependences. *Acc. Chem. Res.* **2011**, *44*, 25-35.

(76) Montes, V. A.; Perez-Bolivar, C.; Agarwal, N.; Shinar, J.; Anzenbacher, P. Molecular-Wire Behavior of OLED Materials: Exciton Dynamics in Multichromophoric Alq<sub>3</sub>-Oligofluorene-Pt(II) Porphyrin Triads. *J. Am. Chem. Soc.* **2006**, *128*, 12436-12438.

(77) Welter, S.; Lafolet, F.; Cecchetto, E.; Vergeer, F.; De Cola, L. Energy Transfer by a Hopping Mechanism in Dinuclear Ir<sup>III</sup> / Ru<sup>II</sup> Complexes: A Molecular Wire? *ChemPhysChem* **2005**, *6*, 2417-2427.

(78) Schlicke, B.; Belser, P.; De Cola, L.; Sabbioni, E.; Balzani, V. Photonic Wires of Nanometric Dimensions. Electronic Energy Transfer in Rigid Rodlike Ru(bpy)<sub>3</sub><sup>2+</sup>-(ph)<sub>n</sub>-Os(bpy)<sub>3</sub><sup>2+</sup> Compounds (ph=1,4-phenylene; n=3, 5, 7). *J. Am. Chem. Soc.* **1999**, *121*, 4207-4214.

(79) Eng, M. P.; Albinsson, B. Non-Exponential Distance Dependence of Bridge-Mediated Electronic Coupling. *Angew. Chem. Int. Ed.* **2006**, *45*, 5626-5629.

(80) Lotter, D.; Neuburger, M.; Rickhaus, M.; Häussinger, D.; Sparr, C. Stereoselective Arene-Forming Aldol Condensation: Synthesis of Configurationally Stable Oligo-1,2-naphthylenes. *Angew. Chem. Int. Ed.* **2016**, *55*, 2930-2933.

1  
2  
3 (81) Witzig, R. M.; Lotter, D.; Fäseke, V. C.; Sparr, C. Stereoselective Arene-Forming  
4 Aldol Condensation: Catalyst-Controlled Synthesis of Axially Chiral Compounds. *Chem. Eur. J.*  
5 **2017**, *23*, 12960-12966.  
6  
7

8  
9  
10 (82) Link, A.; Sparr, C. Stereoselective Arene Formation. *Chem. Soc. Rev.* **2018**, *47*,  
11 3804-3815.  
12  
13

14 (83) Rickhaus, M.; Mayor, M.; Juricek, M. Strain-induced Helical Chirality in  
15 Polyaromatic Systems. *Chem. Soc. Rev.* **2016**, *45*, 1542-1556.  
16  
17

18  
19 (84) Benniston, A. C.; Harriman, A.; Rewinska, D. B.; Yang, S.; Zhi, Y. G. On the  
20 Conjugation Length for Oligo(ethynyl)naphthalene)-Based Molecular Rods. *Chem. Eur. J.* **2007**,  
21 *13*, 10194-10203.  
22  
23

24  
25 (85) Benniston, A. C.; Clift, S.; Hagon, J.; Lemmetyinen, H.; Tkachenko, N. V.; Clegg,  
26 W.; Harrington, R. W. Effect on Charge Transfer and Charge Recombination by Insertion of a  
27 Naphthalene-Based Bridge in Molecular Dyads Based on Borondipyrromethene (Bodipy).  
28 *ChemPhysChem* **2012**, *13*, 3672-3681.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Table of Contents Image

